

## Introduction: naive set theory

$$\mathbb{N} = \{1, 2, 3, \dots\}$$

$$\mathbb{Z} = \{0, \pm 1, \pm 2, \dots\}$$

$$\mathbb{Z}_+ = \{0, 1, 2, \dots\}$$

$$\mathbb{Q} = \left\{ \frac{a}{b} \mid a, b \in \mathbb{Z}, b \neq 0 \right\}$$

$$\mathbb{C} = \{a + bi \mid a, b \in \mathbb{R}\}$$

$$\mathbb{C}_q = \{a + bi \mid a, b \in \mathbb{Q}\}$$

Recall: given sets  $X$  and  $Y$ , a relation from  $X$  to  $Y$  is a subset of  $X \times Y$ , where  $\times$  denotes the cartesian product of  $X$  and  $Y$ .

A relation  $f \subseteq X \times Y$  is a function from  $X$  to  $Y$  such that  $\forall x \in X, \exists! y \in Y$  such that  $(x, y) \in f$ . We write  $f(x) = y$ , and denote  $f$  as  $f : X \rightarrow Y$ .

$X$  is the **domain** of  $f$  and  $Y$  is the **codomain**. The range  $\text{ran}(f) = \{f(x) \mid x \in X\} \subseteq Y$ .

The graph of a function  $\text{Graph}(f) = \{(x, f(x)) \mid x \in X\} \subseteq X \times Y$ .

## Examples

$$\text{id}_x : X \rightarrow X, \text{id}_X(x) = x$$

This is the identity function.

The Characteristic Function: If  $A \subseteq X$

$$\mathbf{1}_A : X \rightarrow \mathbb{R}, \mathbf{1}_A(x) = \begin{cases} 1, & x \in A \\ 0, & x \notin A \end{cases}$$

## Algebra of Functions

Let  $X$  be any set, and  $(X; \mathbb{R}) = \{f : X \rightarrow \mathbb{R}\}$  represent the function space of  $X$  with codomain  $\mathbb{R}$ .

Let  $f, g \in \mathcal{F}(X; \mathbb{R})$ . Then,  $(f + g)(x) = f(x) + g(x)$ , and  $(f \cdot g)(x) = f(x) \cdot g(x)$ .

If  $t \in \mathbb{R}$ , then  $(tf)(x) = tf(x)$  (scalar multiplication). If  $g(x) \neq 0 \forall x \in X$ , then  $\left(\frac{f}{g}\right)(x) := \frac{f(x)}{g(x)}$ .

Finally, we have composition. If  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$  are functions, then  $g \circ f(x) = g(f(x))$ .

### Injective, Subjective, and Bijective

A function  $f : X \rightarrow Y$  is a **injective** map, then, if  $f(x_1) = f(x_2)$ , then  $x_1 = x_2$ . For example, the shift map  $S : \mathbb{N} \rightarrow \mathbb{N}$ ,  $S(n) = n + 1$  is injective.

Any strictly increasing function  $f : I \rightarrow \mathbb{R}$ , where  $I$  is any interval, is injective.

A function  $f$  is **surjective** if  $\forall y \in Y, \exists x \in X$  such that  $f(x) = y$ .

Consider the function  $f : \mathbb{R} \rightarrow \mathbb{R}$ ,  $f(x) = x^3 - 2x + 1$ . We can show that this function is surjective because  $\lim_{x \rightarrow \infty} f(x) = \infty$ ,  $\lim_{x \rightarrow -\infty} f(x) = -\infty$ . Due to the intermediate value theorem, we get that  $\text{ran}(f) = \mathbb{R}$ .

$f$  is **bijective** if it is injective and surjective.

## Invertibility

Let  $f : X \rightarrow Y$  be a function.  $f$  is **left-invertible** if  $\exists g : Y \rightarrow X$  such that  $g \circ f = \text{id}_X$ .  $f$  is **right-invertible** if  $\exists h : Y \rightarrow X$  such that  $f \circ h = \text{id}_Y$ .

$f$  is **invertible** if  $\exists k : Y \rightarrow X$  such that  $f \circ k = \text{id}_Y$  and  $k \circ f = \text{id}_X$ .

## Proposition

$f$  is invertible if and only if  $f$  is left and right invertible.

Forward direction: This is via the definition of invertibility.

Reverse direction: Suppose  $g$  is a left-inverse of  $f$ , and  $h$  is a right-inverse of  $f$ . Therefore,  $g \circ f = \text{id}_X$ , and  $f \circ h = \text{id}_Y$ . Observe that  $g = g \circ \text{id}_Y$ . Therefore,  $g = g \circ (f \circ h)$ . Via associativity,  $g = (g \circ f) \circ h = \text{id}_X \circ h = h$ .

## Theorem

If  $f : X \rightarrow Y$  is a function:

1.  $f$  is injective  $\Leftrightarrow f$  is left-invertible.
2.  $f$  is surjective  $\Leftrightarrow f$  is right-invertible.
3.  $f$  is bijective  $\Leftrightarrow f$  is invertible.

We will prove the first proposition in the forward direction. Suppose  $f$  is injective. Given  $y \in \text{ran}(f)$ , we know that  $\exists! x_y \in X$  such that  $f(x_y) = y$ , by the definition of injective.

Let  $g : Y \rightarrow X$ . We will define  $g$  as follows:

$$g(y) = \begin{cases} x_y & y \in \text{ran}(f) \\ x_0 & y \notin \text{ran}(f) \end{cases}$$

Where  $x_0$  is an arbitrary point in  $X$ . We can see that  $g \circ f = \text{id}_X$ .

For example, the function  $\text{Sin}(x)$  defined as  $\sin(x)$  restricted to  $[-\pi/2, \pi/2]$  has an inverse,  $\arcsin(x) : [-1, 1] \rightarrow [-\pi/2, \pi/2]$ .

## Cardinality and Finitude

Which set is “larger,”  $\{1, 2, 3\}$  or  $\{1, 2, 3, 4\}$ ?  $\mathbb{N}$  or  $\mathbb{N}_0$ ?  $\mathbb{Z}$  or  $\mathbb{Q}$ ?

In order to prove that one set is “the same size” as the other, we can create pairs. For two sets  $A$  and  $B$ , we can show that  $A$  is the same size as  $B$  by creating a function. For example, to show that  $\mathbb{N}$  and  $\mathbb{N}_0$  have the same size, we create  $s : \mathbb{N} \rightarrow \mathbb{N}_0$ ,  $s(n) = n + 1$ .

## Definition

Sets  $A$  and  $B$  have the same **cardinality** if  $\exists$  bijection  $f : A \rightarrow B$ . We write  $\text{card}(A) = \text{card}(B)$ .

## Example

Given  $a < b$  and  $c < d$ , we know that  $\text{card}([a, b]) = \text{card}([c, d])$ .

We can create a linear function from  $[a, b]$  to  $[c, d]$ , and since linear functions are bijections, we know that  $\text{card}([a, b]) = \text{card}([c, d])$ .

## Example 2

$$\text{card}((0, 1)) = \text{card}(\mathbb{R})$$

- $\tan : (-\pi/2, \pi/2) \rightarrow \mathbb{R}$  is a bijection:
  - $\tan$  is strictly increasing (and thus injective)
  - $\lim_{x \rightarrow \infty} \tan(x) = \infty$  and  $\lim_{x \rightarrow -\infty} \tan(x) = -\infty$ , and by intermediate value theorem,  $\tan$  is surjective
- $\ell : (0, 1) \rightarrow (-\pi/2, \pi/2)$  is a bijection as it is a linear function between two intervals.
- Therefore, our bijection is  $\tan \circ \ell : (0, 1) \rightarrow \mathbb{R}$ .

## Definition

A set  $F$  is **finite** if  $F$  is empty or  $\exists n \in \mathbb{N}$  such that  $\text{card}(F) = \text{card}(\{1, 2, \dots, n\})$ . A non-finite set is called infinite.

We can *enumerate*  $F$  by creating a function  $\sigma : \{1, 2, \dots, n\} \rightarrow F$ , such that  $x_j = \sigma(j)$  for  $F = \{x_1, x_2, \dots, x_n\}$ .

## Proposition

If  $m \neq n$ , then  $\text{card}\{1, 2, \dots, m\} \neq \text{card}\{1, 2, \dots, n\}$ .

WLOG, suppose  $m > n$ .

Suppose toward contradiction that  $f : \{1, 2, \dots, m\} \rightarrow \{1, 2, \dots, n\}$  is our bijection. This means there are  $m$  “pigeons” and  $n$  “holes.”

One hole,  $j$ , must contain at least two pigeons (i.e.,  $f(i) = f(k) = j$  for some  $i \neq k \in \{1, 2, \dots, m\}$ ). Since  $f$  is assumed to be injective, this is a contradiction.

## Proposition

$\mathbb{N}$  is infinite.

Suppose toward contradiction that  $\mathbb{N}$  is finite. Thus,  $\exists m \in \mathbb{N}$  such that  $f : \mathbb{N} \rightarrow \{1, 2, \dots, m\}$  is a bijection.

Consider the inclusion  $i : \{1, 2, \dots, m+1\} \rightarrow \mathbb{N}$ .  $i$  is injective.

Then,  $f \circ i : \{1, 2, \dots, m+1\} \rightarrow \{1, 2, \dots, m\}$  is an injection, but by the pigeonhole principle, this cannot be. Therefore, we have reached a contradiction.

## Proposition

If  $A$  is infinite,  $\exists i : \mathbb{N} \hookrightarrow A$ .

$\exists a_1 \in A$ , as  $A \neq \emptyset$ .

$A \setminus \{a_1\} \neq \emptyset$ , so  $\exists a_2 \in A \setminus \{a_1\}$ .

$A \setminus \{a_1, a_2\} \neq \emptyset$ , so  $\exists a_3 \in A \setminus \{a_1, a_2\}$ .

$\vdots$

We thus get a sequence  $\{a_1, a_2, \dots\}$  of distinct elements of  $A$ .

Consider  $f : \mathbb{N} \rightarrow A$ ,  $f(n) = a_n$ .  $f$  is injective as  $a_n$  are distinct.

## Example

$$\text{card}(\mathbb{Z}) = \text{card}(\mathbb{N})$$

$$f : \mathbb{Z} \rightarrow \mathbb{N}$$

$$f(m) = \begin{cases} 2m+1 & m \geq 0 \\ -2m & m < 0 \end{cases}$$

$f$  is a bijection as  $g : \mathbb{N} \rightarrow \mathbb{Z}$ ,  $g(n) = (-1)^{n+1} \lfloor \frac{n}{2} \rfloor$  is the inverse of  $f$ .

## Definition

Given any set  $X$ ,  $\mathcal{P}(X) = \{A \mid A \subseteq X\}$  is the **power set** of  $X$ .

$$2^X := \{f \mid f : X \rightarrow \{0, 1\}\}.$$

## Proposition

$$\text{card}(\mathcal{P}(X)) = \text{card}(2^X)$$

Let  $\varphi : \mathcal{P}(X) \rightarrow 2^X$ .

For  $A \subseteq X$ , put  $\varphi(A) := \mathbf{1}_A$ .

Consider  $\psi : 2^X \rightarrow \mathcal{P}(X)$ .  $\psi(f) = f^{-1}(\{1\}) = \{x \in X \mid f(x) = 1\}$ .

Then,  $\psi \circ \varphi(A) = \psi(\mathbf{1}_A) = \mathbf{1}^{-1}(\{1\}) = A$ ,

and, we claim  $\varphi(\psi(f)) = \varphi(f^{-1}(\{1\})) = \mathbf{1}_{f^{-1}(\{1\})} = f$ .

## Cantor's theorem

$$\nexists \text{ surjection } \mathbb{N} \rightarrow (0, 1)$$

Fact from calculus:  $\forall \sigma \in (0, 1)$ ,  $\sigma$  can be written uniquely as a decimal expansion.

$$\sigma = \sum_{k=1}^{\infty} \frac{\sigma_k}{10^k}$$

Where  $\sigma_k \in \{0, 1, \dots, 9\}$  and not terminating in 9s.

Suppose toward contradiction that  $\exists r : \mathbb{N} \rightarrow (0, 1)$  that is a surjection. Write  $r(n) = 0.\sigma_1(n)\sigma_2(n)\sigma_3(n)\dots$ , and  $\sigma_j(n) \in \{0, 1, \dots, 9\}$ , and not terminating in 9s.

Consider  $\tau : \mathbb{N} \rightarrow \{0, 1, \dots, 9\}$ :

$$\tau(n) = \begin{cases} 3 & \sigma_n(n) = 2 \\ 2 & \sigma_n(n) \neq 2 \end{cases}$$

Let  $\tau = 0.\tau(1)\tau(2)\tau(3)\dots$ . Since  $r$  is surjective,  $\exists m \in \mathbb{N}$  such that  $r(m) = 0.\sigma_1(m)\sigma_2(m)\dots\sigma_m(m)\dots = \tau = 0.\tau(1)\tau(2)\dots\tau(m)\dots$ .

This implies that  $\sigma_m(m) = \tau(m)$ , which is definitionally not true, which is our contradiction.

## Comparing Cardinalities

- $\text{card}(A) \leq \text{card}(B) \Rightarrow \exists f : A \hookrightarrow B$
- $\text{card}(A) < \text{card}(B) \Rightarrow \text{card}(A) \leq \text{card}(B), \text{card}(A) \neq \text{card}(B)$

For example,  $X \subseteq Y \Rightarrow \text{card}(X) \leq \text{card}(Y)$  because  $i : X \hookrightarrow Y, i(x) = x$  is an injection.

### Transitive Property

If  $\text{card}(A) \leq \text{card}(B) \leq \text{card}(C)$ , then  $\text{card}(A) \leq \text{card}(C)$ .

The composition of two injective functions is injective.

### Canonical Set Comparisons

Via the inclusion map, we know the following:

$$\text{card}(\mathbb{N}) \leq \text{card}(\mathbb{Z}) \leq \text{card}(\mathbb{Q}) \leq \text{card}(\mathbb{R})$$

### Cantor-Schröder-Bernstein

For any set  $A$ ,  $\text{card}(A) < \text{card}(\mathcal{P}(A))$ .

Let us construct a function:  $f : A \rightarrow \mathcal{P}(A)$ , where  $a \mapsto \{a\}$ .

$f$  is injective, as if  $\{a\} = \{a'\}$ ,  $a = a'$ . So,  $\text{card}(A) \leq \text{card}(\mathcal{P}(A))$ .

**Claim**  $\nexists g : A \rightarrow \mathcal{P}(A)$ ,  $g$  is surjective.

Suppose toward contradiction that such a  $g$  exists. Consider  $S : \{a \in A \mid a \notin g(a)\}$ .

Since  $g$  is onto,  $\exists a_0 \in A$  with  $g(a_0) = S$ .  $a_0 \in g(a_0) \Leftrightarrow a_0 \in S \Leftrightarrow a_0 \notin g(a_0)$ .  $\perp$

### Equivalent Propositions

- (i)  $\text{card}(A) \leq \text{card}(B)$
- (ii)  $\exists f : A \hookrightarrow B$
- (iii)  $\exists g : B \rightarrow A$ ,  $g$  surjection.

By definition, (i)  $\Leftrightarrow$  (ii).

(ii)  $\Rightarrow$  (iii) If  $f : A \hookrightarrow B$ ,  $f$  is left-invertible, and thus  $\exists g : B \rightarrow A$  with  $g \circ f = id_A$ . So,  $g$  is right-invertible, so  $g$  is surjective.

(iii)  $\Rightarrow$  (ii) If  $g : B \rightarrow A$  is surjective, then  $g$  is right-invertible, so  $\exists f : A \rightarrow B$  such that  $g \circ f = id_B$ . So,  $f$  is left-invertible, so  $f$  is injective.

### Corollary

If  $f : A \rightarrow B$  is any map,  $\text{card}(f(A)) \leq \text{card}(A)$ .

Consider  $g : A \rightarrow f(A)$ , where  $g(a) = f(a)$ . So,  $g$  is onto, so  $\exists$  an injection  $f(A) \hookrightarrow A$ .

### More Cardinality of Canonical Sets

Consider the map  $q : \mathbb{Z} \times \mathbb{N} \rightarrow \mathbb{Q}, q(m, n) = \frac{m}{n}$ . This map is *not* injective, as  $2/4 = 1/2$ . However, it is surjective, meaning  $\text{card}(\mathbb{Q}) \leq \text{card}(\mathbb{Z} \times \mathbb{N})$ .

Earlier, we showed that  $\exists h : \mathbb{Z} \leftrightarrow \mathbb{N}$ . Consider  $H : \mathbb{Z} \times \mathbb{N} \rightarrow \mathbb{N} \times \mathbb{N}$ , defined as  $H(m, n) = (h(m), n)$ .

**Claim**  $H$  is a bijection.

**Proof of Injection** If  $H(m_1, n_1) = H(m_2, n_2)$ , then  $h(m_1) = h(m_2)$ , and  $n_1 = n_2$ , and since  $h$  is bijective,  $m_1 = m_2$ , and  $n_1 = n_2$ , so  $(m_1, n_1) = (m_2, n_2)$ .

**Proof of Surjection** Let  $(k, \ell) \in \mathbb{N} \times \mathbb{N}$ . We want to find  $(m, n) \in \mathbb{Z} \times \mathbb{N}$  such that  $H(m, n) = (k, \ell)$ . Set  $n = \ell$ , and since  $h$  is surjective, set  $m \in \mathbb{Z}$  such that  $h(m) = k$ .

Therefore  $\text{card}(\mathbb{Z} \times \mathbb{N}) = \text{card}(\mathbb{N} \times \mathbb{N})$ .

We claim that  $\text{card}(\mathbb{N} \times \mathbb{N}) = \text{card}(\mathbb{N})$ . First, we need to find  $\varphi : \mathbb{N} \times \mathbb{N} \hookrightarrow \mathbb{N}$ . Consider  $\varphi(m, n) = 2^m \cdot 3^n$ . By the Fundamental Theorem of Arithmetic,  $\varphi$  is injective.

Bringing together our inequalities, we have:

$$\begin{aligned} \text{card}(\mathbb{N}) &\leq \text{card}(\mathbb{Q}) \\ &\leq \text{card}(\mathbb{Z} \times \mathbb{N}) \\ &= \text{card}(\mathbb{N} \times \mathbb{N}) \\ &\leq \text{card}(\mathbb{N}) \end{aligned}$$



## Cardinality Rules

- (i)  $\text{card}(A) \leq \text{card}(A)$  (Reflexivity)
- (ii)  $\text{card}(A) \leq \text{card}(B) \leq \text{card}(C) \Rightarrow \text{card}(A) \leq \text{card}(C)$  (Transitivity)
- (iii)  $\text{card}(A) \leq \text{card}(B)$  and  $\text{card}(B) \leq \text{card}(A) \Rightarrow \text{card}(A) = \text{card}(B)$  (Cantor-Schröder-Bernstein)
- (iv) Either  $\text{card}(A) \leq \text{card}(B)$  or  $\text{card}(B) \leq \text{card}(A)$ .

**Proof of (iii)** We have injections  $f : A \hookrightarrow B$  and  $g : B \hookrightarrow A$ .

Let  $A_0 \setminus \text{ran}(g)$ . Let  $A_1 = g \circ f(A_0)$ . Note that  $A_0 \cap A_1 = \emptyset$ . Let  $A_2 = g \circ f(A_1)$ . Note that  $A_0 \cap A_2 = \emptyset$ .

**Claim** We claim  $A_1 \cap A_2 = \emptyset$ . If  $\exists z \in A_1 \cap A_2$ , then  $z = g(f(x_0))$  for some  $x_0 \in A_0$ , and  $z = g(f(x_1))$  where  $x_1 \in A_1$ . However,  $g$  and  $f$  are injective, so  $g \circ f$  is injective, so  $x_0 = x_1$ , but  $A_0 \cap A_1 = \emptyset$ .

We let  $A_n = g \circ f(A_{n-1})$  for arbitrary  $n$ , and  $A_\infty = \bigcup_{n \geq 0} A_n$ . If  $a \notin A_\infty$ , then  $a \notin A_0$ , so  $a \in \text{ran}(g)$ . Define  $h : A \rightarrow B$ .

$$h(x) = \begin{cases} f(x) & x \in A_\infty \\ y_x & x \notin A_\infty \end{cases}$$

Where  $y_x$  is the unique element in  $B$  with  $g(y_x) = x$ .

**Claim** We claim  $h$  is the desired bijection.

**Proof of Injection** Suppose  $h(x_1) = h(x_2)$ .

If  $x_1, x_2 \in A_\infty$ , then by the definition of  $H$ ,  $f(x_1) = f(x_2)$ ,  $f$  is injective, so  $x_1 = x_2$ .

Suppose  $x_1, x_2 \notin A_\infty$ . Then, by definition,  $h(x_1) = y_{x_1}$  and  $h(x_2) = y_{x_2}$ , then  $g(y_{x_1}) = g(y_{x_2})$ , so  $x_1 = x_2$ .

WLOG, suppose  $x_1 \in A_\infty$ , and  $x_2 \notin A_\infty$ .  $h(x_1) = f(x_1) = h(x_2) = y_{x_2}$ . Then,  $g(f(x_1)) \in A_\infty = g(y_{x_2}) = x_2 \notin A_\infty$ . This case is not possible.

Thus,  $h$  is injective.

**Proof of Surjection** Let  $y \in B$ . Set  $x := g(y)$ .

Suppose  $x \notin A_\infty$ . Then,  $h(x) = y_x$ , where  $y_x$  is the unique element in  $B$  with  $g(y_x) = x = g(y)$ , so  $y = y_x$ , so  $h(x) = y$ .

If  $x \in A_\infty$ . We know that  $x \notin A_0$ , as  $x \in \text{ran}(g)$ . So,  $x = g(f(z))$  for some  $z \in A_{m-1}$ . Since  $g$  is injective,  $y = f(z)$ ,  $z \in A_\infty$ . Thus,  $h(z) = f(z) = y$ .

Therefore, we have  $\text{card}(\mathbb{Q}) = \text{card}(\mathbb{N})$ .

## Countability

A set  $X$  is *countable* if  $\exists f : x \hookrightarrow \mathbb{N}$  ( $\text{card}(X) \leq \text{card}(\mathbb{N})$ ).  $\text{card}(\mathbb{N}) = \aleph_0$ . If  $X$  is countable and infinite,  $X$  is *denumerable*.

## Corollary to Cantor-Schröder-Bernstein

If  $X$  is denumerable, then  $\text{card}(X) = \aleph_0$ .

Since  $X$  is infinite,  $\exists f : \mathbb{N} \hookrightarrow X$ . Since  $X$  is countable,  $\exists g : X \hookrightarrow \mathbb{N}$ . By Cantor-Schröder-Bernstein,  $\text{card}(X) = \text{card}(\mathbb{N})$ , so  $\text{card}(X) = \aleph_0$ .

Thus, we have:

$$\text{card}(\mathbb{N}) = \text{card}(\mathbb{Z}) = \text{card}(\mathbb{Q})$$

(as shown earlier)

## Countability under Union

The countable union of countable sets is countable. If  $I$  is a countable indexing set and for each  $i \in I$ ,  $A_i$  is countable, then  $\bigcup_{i \in I} A_i$  is countable.

Since each  $A_i$  is countable,  $\exists \pi_i : \mathbb{N} \rightarrow A_i$ . Consider the function

$$\pi : I \times \mathbb{N} \rightarrow \bigcup_{i \in I} A_i$$

defined as  $\pi(i, j) = \pi_i(j)$ .

**Claim 1**  $\pi$  is a surjection.

**Proof 1** Let  $x \in \bigcup_{i \in I} A_i$ .  $\exists i_0$  such that  $x \in A_{i_0}$ . Since  $\pi_{i_0}$  is surjective,  $\exists k \in \mathbb{N}$  with  $\pi_{i_0}(k) = x$ .  $\pi_{i_0}(k) = \pi(i_0, k)$ . Therefore,  $\pi$  is surjective.

**Claim 2**  $I \times \mathbb{N}$  is countable.

**Proof 2** We know  $\exists f : I \hookrightarrow \mathbb{N}$  since  $I$  is countable. Thus,  $g : I \times \mathbb{N} \hookrightarrow \mathbb{N} \times \mathbb{N}$ ,  $(i, n) \mapsto (f(i), n)$ . Recall,  $\mathbb{N} \times \mathbb{N} \hookrightarrow \mathbb{N}$ ,  $(m, n) \mapsto 2^m \cdot 3^n$  is an injection. By composing these maps,  $I \times \mathbb{N} \hookrightarrow \mathbb{N}$ . Since  $\pi$  is onto, and  $I \times \mathbb{N}$  is countable,  $\bigcup_{i \in I} A_i$  is countable.

## Continuum Hypothesis

We saw that  $\text{card}(\mathbb{N}) < \text{card}(\mathcal{P}(\mathbb{N})) = \text{card}(2^{\mathbb{N}})$ , where  $2^{\mathbb{N}} = \{f \mid f : \mathbb{N} \rightarrow \{0, 1\}\}$ .

**Theorem**  $\text{card}(\mathbb{R}) = \text{card}(I) = \text{card}(2^{\mathbb{N}})$ , where  $I$  is any non-degenerate interval.

**Lemma 1**  $\text{card}([0, 1]) \leq \text{card}(2^{\mathbb{N}})$ .

**Proof 1** Every  $t \in [0, 1]$  has a binary expansion.

$$t = \sum_{k=1}^{\infty} \frac{\sigma_k}{2^k}$$

where  $\sigma_k \in \{0, 1\}$ .

Consider  $2^{\mathbb{N}} \xrightarrow{\varphi} [0, 1]$ , defined as  $\phi(f) = \sum_{k=1}^{\infty} \frac{f(k)}{2^k}$ . Set  $f : \mathbb{N} \rightarrow \{0, 1\}$ ,  $f(k) = \sigma_k$ .

Therefore,  $\varphi$  is surjective, so  $\exists \{0, 1\} \hookrightarrow 2^{\mathbb{N}}$ , so  $\text{card}([0, 1]) \leq 2^{\mathbb{N}}$

**Lemma 2**  $\text{card}([0, 1]) = \text{card}(\mathbb{R})$ .

**Proof 2** We have  $[0, 1] \xrightarrow{i} \mathbb{R}$  via inclusion, so  $\text{card}([0, 1]) \leq \text{card}(\mathbb{R})$ .

Also,  $\text{card}(\mathbb{R}) = \text{card}((0, 1)) \leq \text{card}([0, 1])$ , so by Cantor-Schröder-Bernstein,  $\text{card}(\mathbb{R}) = \text{card}([0, 1])$ .

**Lemma 3** Any two non-degenerate intervals  $I$  and  $J$  have the same cardinality.

**Proof 3** We can create injections  $I \hookrightarrow J$  and vice-versa.

**Lemma 4**  $\text{card}(2^{\mathbb{N}}) \leq \text{card}([0, 1])$ .

**Proof 4**  $\psi : 2^{\mathbb{N}} \rightarrow [0, 1]$ . Where  $\psi(f) = \sum_{k=1}^{\infty} \frac{f(k)}{3^k}$ .

$\psi$  is well-defined:

$$0 \leq \sum_{k=1}^{\infty} \frac{f(k)}{3^k} \leq \sum_{k=1}^{\infty} \frac{1}{3^k} \leq \frac{1}{2} \leq 1$$

We claim  $\psi$  is injective. Suppose  $f \neq g$  in  $2^{\mathbb{N}}$ . Let  $k_0 = \min\{k \mid f(k) \neq g(k)\}$ . WLOG,  $f(k_0) = 0, g(k_0) = 1$ . Let  $t_f = \sum_{k>k_0}^{\infty} \frac{f(k)}{3^k}$ ,  $t_g = \sum_{k>k_0}^{\infty} \frac{g(k)}{3^k}$ .

Therefore,  $\psi(f) = \sum_{k=1}^{k_0-1} \frac{f(k)}{3^k} + 0 + t_f$ , and  $\psi(g) = \sum_{k=1}^{k_0-1} \frac{1}{3^k} + t_g$ .

Suppose toward contradiction  $\psi(f) = \psi(g)$ . Then,  $t_f = \frac{1}{3^{k_0}} + t_g$ , or  $t_f - t_g = \frac{1}{3^{k_0}}$ .

$$\begin{aligned} |t_f - t_g| &= \left| \sum_{k>k_0} \frac{f(k)}{3^k} - \sum_{k>k_0} \frac{g(k)}{3^k} \right| \\ &\leq \sum_{k>k_0} \frac{|f(k) - g(k)|}{3^k} \\ &\leq \sum_{k>k_0} \frac{1}{3^k} \\ &= \frac{(1/3)^{k_0+1}}{1 - (1/3)} \\ &= \frac{1}{2} \cdot \frac{1}{3^{k_0}} \end{aligned}$$

$\perp$

We have thus shown:

$$\text{card}(\mathbb{R}) = \text{card}([0, 1]) = \text{card}(2^{\mathbb{N}})$$

We know that

$$\aleph_0 = \text{card}(\mathbb{N}) = \text{card}(\mathbb{Q}) = \text{card}(\mathbb{Z}) < 2^{\aleph_0} = \text{card}(2^{\mathbb{N}}) = \text{card}(\mathbb{R}) = \text{card}(I)$$

However, the existence of an infinity with cardinality strictly greater than  $\aleph_0$  and strictly less than  $2^{\aleph_0}$  is an axiom (i.e., it can be an assumption or not).

## Ordering

Let  $X$  be a non-empty set. A relation on  $X$  is a subset of  $X \times X$ .

- $R$  is *reflexive* if  $\forall x \in X, (x, x) \in R$ .
- $R$  is *transitive* if  $(x, y), (y, z) \in R \rightarrow (x, z) \in R$ .
- If  $R$  is *antisymmetric*  $(x, y), (y, x) \in R \rightarrow x = y$ .

If  $R$  is reflexive, transitive, and antisymmetric, then  $R$  is an *ordering* of  $X$ .

If  $R$  is an ordering of  $X$ , then we write:

$$(x, y) \in R \Leftrightarrow xRy \Leftrightarrow x \leq_R y$$

- $x \leq_R x \forall x \in X$
- $x \leq_R y, y \leq_R z \rightarrow x \leq_R z$
- $x \leq_R y, y \leq_R x \rightarrow x = y$

Additionally,  $x <_R y$  means  $x \leq_R y$  and  $x \neq y$ .

Algebraic ordering of  $\mathbb{N}_0$ 

$$n \leq_a m \Leftrightarrow \exists k \in \mathbb{N}_0 \text{ such that } n + k = m$$

 $\mathbb{N}$  ordered via division

$$n \leq_D m \Leftrightarrow n \mid m$$

Under this definition, it is false that  $2 \leq_D 5$ , but it is true that  $4 \leq_D 20$ .

**Inclusion** Let  $S$  be any set, and let  $X = \mathcal{P}(S)$ . For  $A, B \in \mathcal{P}(S)$ , we define  $A \leq_i B \Leftrightarrow A \subseteq B$ .

**Containment** With  $X$  defined as above,  $A \leq_c B \Leftrightarrow A \supseteq B$ .

For  $\mathcal{F}(X, \mathbb{R}) = \{f \mid f : X \rightarrow \mathbb{R}\}$ , we can define  $f \leq g \Leftrightarrow f(x) \leq g(x) \forall x \in X$ .

## Types of Orderings

- An ordering  $\leq$  of  $X$  is *total* or *linear* if  $\forall x, y \in X, x \leq y$  or  $y \leq x$ .
- An ordering is *directed* if  $\forall x, y \in X \exists z \in X$  such that  $x \leq z$  and  $y \leq z$ .

If  $X$  is a totally ordered set,  $X$  is directed.

For example, all the following orderings are directed but not total:

$$(\mathbb{N}_0, \leq_D), (\mathcal{P}(S), \leq_i), (\mathcal{P}(S), \leq_c)$$

## Upper/Lower Bounds

- (i) Let  $(X, \leq)$  be an ordered set,  $A \subseteq X$ .  $A$  is bounded above if  $\exists v \in X$  with  $a \leq v \forall a \in A$ . Such a  $v$  is an upper bound.

- (ii)  $A$  is bounded below if  $\exists \ell \in X$  such that  $a \geq \ell \forall a \in A$ . Such a  $w$  is a lower bound.
- (iii) If  $v$  is an upper bound of  $A$  and  $v \in A$ , then  $v$  is the greatest element of  $A$ , or  $\max(A) = v$ .
- (iv) If  $\ell$  is a lower bound for  $A$  and  $\ell \in A$ , then  $\ell$  is the least element of  $A$ , or  $\min(A) = \ell$ .
- (v) If  $u$  is an upper bound for  $A$ , and  $u \leq v$  for all other upper bounds  $v$  of  $A$ , then  $u$  is the *least upper bound* of  $A$ , or  $\sup(A) = u$  (for *supremum*).
- (vi) If  $\ell$  is a lower bound for  $A$ , and  $\ell \leq g$  for all other lower bounds  $g$  of  $A$ , then  $\ell$  is the *greatest lower bound* of  $A$ , or  $\inf(A) = \ell$  (for *infimum*).
- (vii) If  $A$  is bounded above and below, then  $A$  is bounded.

### Well-Ordering Principle

With  $(\mathbb{N}, \leq_a)$ , every nonempty  $A \subseteq \mathbb{N}$  has a least element.

### Examples

#### Example 1

For  $A \subseteq (\mathbb{N}, \leq_a)$ ,  $A = \{2, 3, \dots, 12\}$ , we have the following:

**Bounded Above?** Yes.

**Upper Bounds** 12, 13, 14, ...

**Greatest Element** 12

#### Example 2

For  $A \subseteq (\mathbb{N}, \leq_D)$ ,  $A = \{2, 3, \dots, 10\}$

**Bounded Above?** Yes.

**Upper Bounds** 10!

**Greatest Element?** No.

**Supremum**  $2^3 \cdot 3^2 \cdot 5 \cdot 7$

**Bounded Below?** Yes.

**Lower Bound** 1

**Least Element?** No.

**Infimum** 1

**Example 3**

For  $\mathcal{A} \subseteq (\mathcal{P}(S), \leq_i)$ ,  $A = \{A_i\}_{i \in I} \subseteq \mathcal{P}(S)$ .

**Supremum**  $\bigcup_{i \in I} A_i$

**Infimum**  $\bigcap_{i \in I} A_i$

**Complete Sets**

An ordered set  $(X, \leq)$  is *complete* if for all  $A \subseteq X$  bounded,  $\inf(A)$  and  $\sup(A)$  exist.

For example,  $\mathbb{Q}$  is *not* complete, as there is not a largest rational number less than  $\sqrt{2}$ , for example.

**Ordering of  $\mathbb{Z}$** 

$$n \leq_a m \Leftrightarrow \exists k \in \mathbb{N}_0, n + k = m$$

This defines a total and complete ordering.

Define  $\mathbb{Z}^+ = \{m \in \mathbb{Z} \mid 0 \leq_a m\}$

**Properties of  $\mathbb{Z}^+$** 

- (i)  $m, n \in \mathbb{Z} \Rightarrow m + n \in \mathbb{Z}^+, m \cdot n \in \mathbb{Z}^+$
- (ii)  $m \in \mathbb{Z}$ , then  $m \in \mathbb{Z}^+$  or  $-m \in \mathbb{Z}^+$
- (iii)  $m, -m \in \mathbb{Z}^+$ , then  $m = 0$
- (iv)  $m \leq_a n \Leftrightarrow n - m \in \mathbb{Z}^+$

**Ordering of  $\mathbb{Z}$ ,  $\mathbb{Q}$ , and  $\mathbb{R}$** 

Recall the ordering of  $\mathbb{Z}$ :

$$n \leq_a m \stackrel{\text{def}}{\Leftrightarrow} \exists k \in \mathbb{N}_0 \text{ with } n + k = m$$

**Claim**  $\leq_a$  is an ordering of  $\mathbb{Z}$

We claim that  $\mathbb{Z}^+ = \{m \in \mathbb{Z} \mid 0 \leq_a m\}$ . Thus,  $\mathbb{Z}^+ = \mathbb{N}_0$ .

**Properties of  $\mathbb{Z}^+$** 

- (i)  $m, n \in \mathbb{Z} \Rightarrow m + n \in \mathbb{Z}^+, m \cdot n \in \mathbb{Z}^+$
- (ii)  $m \in \mathbb{Z}$ , then  $m \in \mathbb{Z}^+$  or  $-m \in \mathbb{Z}^+$
- (iii)  $m, -m \in \mathbb{Z}^+$ , then  $m = 0$
- (iv)  $m \leq_a n \Leftrightarrow n - m \in \mathbb{Z}^+$

Other Properties of  $\mathbb{Z}$ 

- (1)  $n \leq_a m \Leftrightarrow m - n \in \mathbb{Z}^+$
- (2)  $m \leq_a n$  and  $p \leq_a q \Rightarrow m + p \leq_a n + q$
- (3)  $m \leq_a n$  and  $p \in \mathbb{Z}^+ \Rightarrow pm \leq_a pn$
- (4)  $m \leq_a n \Rightarrow -m \geq_a n$
- (5)  $\leq_a$  is total.
- (6) If  $a_a > -$ , and  $ab_a \geq 0$ , then  $b_a > 0$
- (7) If  $a > 0$  and  $ab_a \geq ac$ , then  $b \geq c$ .

**Proof of (3):**

$m \leq_a n \Rightarrow \exists k \in \mathbb{N}_0$  with  $m + k = n$ .  
 $\Rightarrow pm + pk = pn$   
 $pk \in \mathbb{N}_0$  by the properties of  $\mathbb{Z}^+$ . So,  $pm \leq_a pn$

**Proof of (5):**

Let  $m, n \in \mathbb{Z}$ . Consider  $m - n$ .  
 By (ii),  $m - n \in \mathbb{Z}^+$  or  $-(m - n) \in \mathbb{Z}^+$ . Thus,  $m - n = k$  for some  $k \in \mathbb{Z}^+$ , or  $-(m - n) = \ell$  for some  $\ell \in \mathbb{Z}^+$ .  
 Thus,  $n \leq_a m$  in the first case, or  $m \leq_a n$  in the second case.

We now want an ordering on  $\mathbb{Q}$ .

### Creating the Rationals

Recall that  $Q = \mathbb{Z} \times \mathbb{Z}^* = \{(a, b) \mid a \in \mathbb{Z}, b \in \mathbb{Z}, b \neq 0\}$ . Consider the equivalence relation:

$$(a, b) \sim (c, d) \stackrel{\text{def}}{\iff} ad = bc$$

We will let  $\mathbb{Q} = \{[(a, b)] \mid (a, b) \in Q\}$  be the set of all equivalence classes in  $Q$ . We write:

$$[(a, b)] = \frac{a}{b}$$

We define addition as follows:

$$\frac{a}{b} + \frac{c}{d} = \frac{ad + bc}{bd}$$

We must check that addition is well-defined:  $\frac{a'}{b'} = \frac{a}{b}$  and  $\frac{c'}{d'} = \frac{c}{d}$ , then  $\frac{a'd' + c'b'}{b'd'} = \frac{ad + bc}{bd}$ .

We define multiplication as follows:

$$\frac{a}{b} \cdot \frac{c}{d} = \frac{ac}{bd}$$

These operations make  $\mathbb{Q}$  a **field**:

### Fields

A ring is a nonempty set  $R$  equipped with two binary operations:

- $+: R \times R \rightarrow R, (a, b) \mapsto a + b$  ("addition")
- $\cdot: R \times R \rightarrow R, (a, b) \mapsto a \cdot b$  ("multiplication")

such that the following hold:

- (1)  $(a + b) + c = a + (b + c)$
- (2)  $\exists z \in R$  such that  $a + z = a = z + a \forall a \in R$ ; there is at most one such  $z$ . Set  $z = 0_R$ .
- (3)  $\forall a \in R, \exists b \in R$  such that  $a + b = 0_R = b + a$ ; there is at most one such  $b$ . Set  $b = -a$ .
- (4)  $\forall a, b \in R, a + b = b + a$ .
- (5)  $(a \cdot b) \cdot c = a \cdot (b \cdot c)$
- (6)  $a \cdot (b + c) = a \cdot b + a \cdot c, (a + b) \cdot c = a \cdot c + b \cdot c$

The above six rules define a ring. If  $(R, +, \cdot)$  satisfies  $ab = ba$ ,  $R$  is a commutative ring.

If there exists  $u \in R$  such that  $ua = au = a \forall a \in R$ ,  $R$  is a unital ring; there is at most one unit. Set  $u = 1_R$ .

An integral domain is a unital, commutative ring such that  $ab = 0 \Rightarrow a = 0 \vee b = 0$ . For example,  $\mathbb{Z}$  is an integral domain. However,  $c(\mathbb{R}) = \{f : \mathbb{R} \rightarrow \mathbb{R} \mid f \text{ continuous}\}$  is a unital, commutative ring, but there exist two functions such that  $f, g \neq \mathbf{0}$ , but  $f \cdot g = \mathbf{0}$ .

A field is a unital, commutative ring such that every element has a multiplicative inverse.

$$\forall a \in R, a \neq 0_R, \exists b \in R, \text{ with } ab = 1_R$$

There is only one such  $b$ . Set  $b = a^{-1}$ .

### Proof that $\mathbb{Q}$ is a Field:

$$\left(\frac{a}{b}\right)^{-1} = \frac{b}{a}$$



Ordering of  $\mathbb{Q}$ 

$$\frac{a}{b} \leq_a \frac{c}{d} \Leftrightarrow ad \leq_a bc \in \mathbb{Z}$$

Prove that this ordering is well-defined.

## Order Embedding

$\leq$  is a well-defined total ordering of  $\mathbb{Q}$ , and  $j : \mathbb{Z} \hookrightarrow \mathbb{Q}$ ,  $j(n) = \frac{n}{1}$  is an order embedding.

$$j(n) \leq j(m) \Leftrightarrow n \leq_a m \in \mathbb{Z}$$

Properties of  $\mathbb{Q}^+$ 

$$\mathbb{Q}^+ = \{q \in \mathbb{Q} \mid q \geq 0_{\mathbb{Q}}\}$$

- (i)  $q_1, q_2 \in \mathbb{Q}^+ \Rightarrow q_1 + q_2 \in \mathbb{Q}^+, q_1 q_2 \in \mathbb{Q}^+$
- (ii)  $q \in \mathbb{Q} \Rightarrow q \in \mathbb{Q}^+ \vee -q \in \mathbb{Q}^+$
- (iii)  $\pm q \in \mathbb{Q}^+, q = 0$
- (iv)  $x \leq y, !u \leq v \Rightarrow x + u \leq y + v$
- (v)  $x \leq y, 0 \leq z \Rightarrow zx \leq zy$

Ordering of  $\mathbb{R}$ , cont'd

An **ordered field** is a field  $F$  equipped with a total ordering  $\leq_F$  such that:

- (i) if  $s \leq_F t$ , and  $x \leq_F y$ , then  $s + x \leq_F t + y$
- (ii) if  $s \leq_F t$  and  $0 \leq_F z$ , then  $zs \leq_F zt$

For example,  $\mathbb{Q}$  with its ordering is an ordered field.

**Proposition 1:** If  $(F, \leq_F)$  is an ordered field, we define  $F^+ = \{x \in F, x_F \geq 0\}$  with the following properties:

- (1)  $x, y \in F^+ \Rightarrow x + y \in F^+, xy \in F^+$
- (2)  $x \in F \Rightarrow x \in F^+, -x \in F^+$
- (3)  $\pm x \in F^+ \Rightarrow x = 0_F$

## Proofs

- (1) Let  $x, y \in F^+$ . Then,  $x \geq 0$  and  $y \geq 0$ , so by property (i) of an ordered field,  $x + y \geq 0 + 0 = 0$ , so  $x + y \in F^+$ . Additionally, we have  $x \cdot y \geq x \cdot 0 = 0$ , so  $xy \in F^+$ .
- (2) Let  $x \in F$ . Since the ordering on  $F$  is total,  $x \geq 0$  or  $0 \geq x$ . In the first case,  $x \in F^+$ . In the second case, we add  $-x$  to both sides, so by (i),  $-x \geq 0$ , so  $-x \in F^+$ .
- (3) We have  $x \geq 0$  and  $-x \geq 0$ . So  $x \geq 0$  and  $x + (-x) \geq x + 0$ , so  $x \geq 0$  and  $0 \geq x$ . So,  $x = 0$  by antisymmetry.

**Note:**  $x \leq_F y \Leftrightarrow y - x \in F^+$ .

**Proposition 2:** Let  $F$  be an ordered field. Then, the following is true:

- (1)  $\forall a \in F, a^2 \in F^+$
- (2)  $0, 1 \in F^+$
- (3) If  $n \in \mathbb{N}$ ,  $n \cdot 1_F = \underbrace{1_F + 1_F + \cdots + 1_F}_{n \text{ times}}$
- (4) If  $x \in F^+$ , and  $x \neq 0$ , then  $x^{-1} \in F^+$
- (5) If  $xy > 0$ , then  $x, y \in F^+$ , or  $-x, -y \in F^+$
- (6) If  $0 < x \leq y$ , then  $0 < y^{-1} \leq x^{-1}$
- (7) If  $x \leq y$ , then  $-y \leq -x$
- (8)  $x \geq 1 \Rightarrow x^2 \geq x \geq 1$ , and  $0 \leq x \leq 1 \Rightarrow 0 \leq x^2 \leq x \leq 1$ .

#### Proofs

- (1) Let  $a \in F$ . Then,  $a \in F^+$  or  $-a \in F^+$ .  
CASE 1 If  $a \in F^+$ , then by the previous proposition,  $a^2 \in F^+$ .  
CASE 2 If  $-a \in F^+$ , then by the previous proposition,  $(-a)(-a) = a^2 \in F^+$ .
- (2)  $0 \geq 0$ , so  $0 \in F^+$ .  $1 = 1 \cdot 1 = 1^2 \in F^+$  by the previous result.
- (3)  $n \cdot 1_F = \underbrace{1_F + 1_F + \cdots + 1_F}_{n \text{ times}} \in F^+$  by the previous proposition.
- (4) Let  $x \neq 0, x \in F^+$ . Suppose toward contradiction that  $x^{-1} \notin F^+$ , then  $-x^{-1} \in F^+$ . Thus,  $x \cdot (-x^{-1}) \in F^+$ , so  $-1 \in F^+$ , but  $1 \in F^+$ , so  $1 = 0$ .  $\perp$
- (5) Let  $xy > 0$ , meaning  $xy \in F^+$ . Suppose toward contradiction that  $x > 0$  and  $y < 0$ . So,  $x > 0$  and  $-y > 0$ , so  $(x)(-y) > 0$ , so  $-(xy) \in F^+$ , so  $xy = 0$ .  $\perp$
- (6) Let  $0 < x \leq y$ . We know  $x^{-1} \in F^+$ , so  $x^{-1}x \leq x^{-1}y$ . So  $1 \leq x^{-1}y$ . We also know  $y \in F^+$ , so  $y^{-1} \in F^+$ . So,  $1 \cdot y^{-1} \leq x^{-1} \cdot y \cdot y^{-1}$ .
- (7) Let  $x \leq y$ . Then,  $0 \leq y - x$ , so  $-y \leq -x$ .
- (8) Let  $x \geq 1$ . Then,  $x \cdot x \geq 1 \cdot x \geq 1$ .

#### Order Axiom

$\mathbb{R}$  is an ordered field. The injection  $\mathbb{Q} \hookrightarrow \mathbb{R}$ ,  $i(q) = q$  is an order embedding.

#### Rational Orderings

**Proposition 1:** If  $a \leq b$ , then  $a \leq \frac{1}{2}(a + b) \leq b$

**Proof**

$2a = a + a \leq a + b \leq b + b$ , all by property (i) of an ordered field.

Therefore,  $2a \leq a + b \leq 2b$ . Since  $2 = 1 + 1 \in \mathbb{R}^+$ ,  $2^{-1} \in \mathbb{R}^+$ , so  $(2a)/2 \leq \frac{1}{2}(a + b) \leq (2b)/2$ , so  $a \leq \frac{1}{2}(a + b) \leq b$ .

**Proposition 2:** If  $a \geq 0$  and  $(\forall \varepsilon > 0), a \leq \varepsilon$ .

**Proof**

If  $a \geq 0$  and  $a \neq 0$ , then  $a > 0$ . So, we have that  $\frac{1}{2}a < a$ . Let  $\varepsilon = \frac{1}{2}a$ . We also have that  $a \leq \varepsilon = \frac{1}{2}a < a$ , so  $a < a$ .  $\perp$

**Arithmetic and Geometric Means**

Given  $a_1, a_2, \dots, a_n \in \mathbb{R}^+$ :

**Arithmetic Mean**

$$= \frac{\sum_{i=1}^n a_i}{n}$$

**Geometric Mean**

$$= \sqrt[n]{a_1 a_2 \cdots a_n}$$

**Arithmetic Mean-Geometric Mean Inequality**

Let  $a, b \geq 0$ .

$$(ab)^{1/2} \leq \frac{1}{2}(a + b)$$

If  $x, y \geq 0$ ,  $x \leq y \Leftrightarrow x^2 \leq y^2$ .

$$0 \leq x \cdot x \leq x \cdot y \leq y \cdot y \quad \text{by property (ii) of ordered fields}$$

Therefore,

$$(ab)^{1/2} \leq \frac{1}{2}(a + b)$$

$$ab \leq \frac{1}{4}(a^2 + 2ab + b^2)$$

$$4ab \leq a^2 + 2ab + b^2$$

$$0 \leq a^2 - 2ab + b^2$$

$$0 \leq (a - b)^2 \quad \text{by definition}$$

**Challenge:** Prove for  $m$ .

**Remark:** The harmonic mean is defined as:

$$\frac{n}{\sum_{i=1}^n \frac{1}{a_i}}$$

### Bernoulli's Inequality

If  $x \geq -1$ , then  $(1+x)^n \geq 1+nx$ , for any  $n \in \mathbb{N}_0$

By induction, we know that for  $n = 0$  and  $n = 1$ , this holds.

Assume the inequality holds for some  $m \geq 1$ .

$$\begin{aligned} (1+x)^{m+1} &= (1+x)^m(1+x) \\ &\geq (1+mx)(1+x) && \text{by the inductive hypothesis} \\ &= 1+x+mx+mx^2 \\ &= 1+(m+1)x+mx^2 \\ &\geq 1+(m+1)x \end{aligned}$$

### Cauchy's Inequality

Let  $a_1, \dots, a_n, b_1, \dots, b_n \in \mathbb{R}$ . Then

$$\left| \sum_{j=1}^n a_j b_j \right| \leq \left( \sum_{j=1}^n a_j^2 \right)^{1/2} \left( \sum_{j=1}^n b_j^2 \right)^{1/2}$$

In linear algebra language, this is equivalent to  $\vec{v} \cdot \vec{w} \leq \|\vec{v}\| \cdot \|\vec{w}\|$ .

Consider  $f : \mathbb{R} \Rightarrow \mathbb{R}$

$$f(x) = \sum_{i=1}^n (a_i - b_i x)^2$$

We know that  $f(x) \geq 0$  for all  $x \in \mathbb{R}$

$$\begin{aligned} &= \sum_{i=1}^n (a_i^2 - 2a_i b_i x + b_i^2 x^2) \\ &= \left( \sum_{j=1}^n b_j^2 \right) x^2 + \left( \sum_{j=1}^n -2a_j b_j \right) x + \sum_{j=1}^n a_j^2 \\ &= Ax^2 + Bx + C \end{aligned}$$

Therefore,  $\Delta = B^2 - 4AC \leq 0 \Rightarrow B^2 \leq 4AC$

$$\left(-2 \sum_{j=1}^n a_j b_j\right)^2 \leq 4 \left(\sum_{j=1}^n a_j\right) \left(\sum_{j=1}^n b_j\right)$$

$$\left|\sum_{j=1}^n a_j b_j\right| = \left(\sum_{j=1}^n a_j\right)^{1/2} \left(\sum_{j=1}^n b_j\right)^{1/2}$$

As we know from linear algebra, the way we get equality is when  $\vec{v} = c\vec{w}$ , or that  $a_j = cb_j \forall j$  for some  $c \in \mathbb{R}$ .

### Triangle Inequality

Given  $a_1, \dots, a_n, b_1, \dots, b_n \in \mathbb{R}$

$$\left(\sum_{j=1}^n (a_j + b_j)^2\right)^{1/2} \leq \left(\sum_{j=1}^n a_j^2\right)^{1/2} + \left(\sum_{j=1}^n b_j^2\right)^{1/2}$$

In linear algebra, this is equivalent to  $\|\vec{v} + \vec{w}\| \leq \|\vec{v}\| + \|\vec{w}\|$ .

$$\begin{aligned} \sum (a_j + b_j)^2 &= \sum a_j^2 + \sum 2a_j b_j + \sum b_j^2 \\ &\leq \sum a_j^2 + 2 \left(\sum a_j^2\right)^{1/2} \left(\sum b_j^2\right)^{1/2} + \sum b_j^2 && \text{by Cauchy} \\ &= \left(\left(\sum a_j^2\right)^{1/2} + \left(\sum b_j^2\right)^{1/2}\right)^2 \end{aligned}$$

we take square roots to get our end result

### Metrics and Norms on $\mathbb{R}^n$

Consider  $|\cdot| : \mathbb{R} \rightarrow \mathbb{R}$ , defined as follows:

$$|x| := \begin{cases} x, & x \in \mathbb{R}^+ \\ -x, & x \notin \mathbb{R}^+ \end{cases}$$

#### Theorems about Absolute Value:

- (i)  $|ab| = |a||b|$
- (ii)  $|a^2| = |a|^2$
- (iii)  $|-a| = |a|$
- (iv)  $|a| \in \mathbb{R}^+$
- (v)  $-|a| \leq a \leq |a|$
- (vi)  $|a| \leq \delta \Rightarrow -\delta \leq a \leq \delta$  for  $\delta > 0$
- (vii)  $|a + b| \leq |a| + |b|$ ,  $|a - b| \leq |a| + |b|$ ,  $||a| - |b|| \leq |a - b|$

## Proofs

Proof of (i)

**Case 1:** If  $a, b \in \mathbb{R}^+$ , then  $|a| = a$ , and  $|b| = b$ , and  $ab \in \mathbb{R}^+$ , so  $|ab| = ab$

**Case 2:** If  $a, b \notin \mathbb{R}^+$ , then  $|a| = -a$ , and  $|b| = -b$ . Additionally,  $(-a)(-b) = ab \in \mathbb{R}^+$ , so  $|ab| = ab$ . The LHS =  $ab$ , and the RHS =  $ab$ .

**Case 3:**  $a \in \mathbb{R}^+$ ,  $-b \in \mathbb{R}^+$ . Then,  $|a||b| = (a)(-b) = -ab$ . Then, since  $a(-b) \in \mathbb{R}^+$ ,  $-ab \in \mathbb{R}^+$ , so  $|ab| = -ab$ . Therefore, the LHS and RHS are equal.

Proof of (vii) Having established that  $|a + b| \leq |a| + |b|$ , we will show that  $||a| - |b|| \leq |a - b|$ .

$$\begin{aligned} |a| &= |a - b + b| \\ &\leq |a - b| + |b| \\ |a| - |b| &\leq |a - b| \end{aligned}$$

Similarly, by exchanging  $a$  for  $b$

$$\begin{aligned} |b| - |a| &\leq |b - a| \\ |b| - |a| &\leq |a - b| \end{aligned}$$

Let  $t = |a| - |b|$ . We have shown that

$$\begin{aligned} \pm t &\leq |a - b| \\ -|a - b| &\leq t \leq |a - b| \\ |t| &\leq |a - b| \end{aligned}$$

## Absolute Values, cont'd

Recall:

$$|x| = \begin{cases} x, & x \in \mathbb{R}^+ \\ -x, & x \notin \mathbb{R}^+ \end{cases}$$

If we want to find all  $x \in \mathbb{R}$  such that  $|x - 1| \leq |x|$ , we would split up into cases:

$x \leq 0$   $x - 1 \leq -1$ , so  $|x| = -x$  and  $|x - 1| = 1 - x$ , so  $1 - x \leq -x$ , so  $0 \geq 1$ .  $\perp$

$0 < x \leq 1$   $|x| = x$  and  $|x - 1| = 1 - x$ , so  $1 - x \leq x$ , so  $x \geq \frac{1}{2}$ , so  $\frac{1}{2} \leq x \leq 1$ .

$1 < x$   $|x| = x$  and  $|x - 1| = x - 1$ , so  $x - 1 \leq x$ , so  $-1 \leq 0$ , which is true  $\forall \mathbb{R}$  in the interval, so  $x > 1$ .

Therefore, we have  $x \in (\frac{1}{2}, \infty)$  as that which satisfies this inequality.

## Bounded Sets

A subset  $A \subseteq \mathbb{R}$  is **bounded**  $\Leftrightarrow \exists c \geq 0$  such that  $\forall x \in A$ ,  $|x| \leq c$ .

( $\Rightarrow$ ) Suppose  $A \subseteq \mathbb{R}$  is bounded. Then,  $\exists \ell, u \in \mathbb{R}$  such that  $\ell \leq x \leq u \forall x \in A$ . Let  $c := \max\{|\ell|, |u|\}$ .

Since  $|u| \leq c$ , we have that  $x \leq c$ .

Since  $|\ell| \leq c$ , and  $-\ell \leq x$ , we get that  $-x \leq |\ell| \leq c$ .

Since  $x \leq c$  and  $-x \leq c$ ,  $|x| \leq c$ .

( $\Leftarrow$ ) If such a  $c$  exists, then  $|x| \leq c$  if and only if  $-c \leq x \leq c$ . Thus,  $-c$  is the lower bound and  $c$  is the upper bound.

### Bounded Functions

Let  $D$  be any set. A function  $f : D \rightarrow \mathbb{R}$  is bounded if  $\text{ran}(D) \subseteq \mathbb{R}$  is bounded.

### Example

Let  $f : [3, 7] \rightarrow \mathbb{R}$ ,  $f(x) = \frac{x^2+2x+1}{x-1}$ . Show that  $f$  is bounded.

$$3 \leq x \leq 7 \Rightarrow 2 \leq x - 1 \leq 6 \Rightarrow \frac{1}{6} \leq \frac{1}{x-1} \leq \frac{1}{2}.$$

$$\text{Also, } 4 \leq x + 1 \leq 8 \Rightarrow 16 \leq x^2 + 2x + 1 \leq 64 \Rightarrow |x^2 + 2x + 1| \leq 64.$$

$$\text{So, } |f(x)| \leq 32.$$

### Distance Metrics

For  $s, t \in \mathbb{R}$ , we will define  $d(s, t) = |s - t|$  to be the **distance** between  $s$  and  $t$ .

#### Properties:

(i)

$$\begin{aligned} d : \mathbb{R} \times \mathbb{R} &\rightarrow [0, \infty) \\ (s, t) &\mapsto d(s, t) \geq 0 \end{aligned}$$

(ii)  $d(s, t) = d(t, s)$

(iii)  $d(s, r) \leq d(s, t) + d(t, r)$

(iv)  $d(s, s) = 0$

(v) If  $d(s, t) = 0$ , then  $s = t$ .

Let  $v = \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix}, w = \begin{pmatrix} y_1 \\ \vdots \\ y_n \end{pmatrix} \in \mathbb{R}^n$ .

• 1-norm:

$$\|v\|_1 = \sum_{j=1}^n |x_j|$$

•  $\infty$ -norm:

$$\|v\|_\infty = \max_{j=1}^n |x_j|$$

• 2-norm:

$$\|v\|_2 = \left( \sum_{j=1}^n x_j^2 \right)^{1/2}$$



### Properties of the Norms

**Properties:** With  $v, w$  above, let  $p = 1, 2, \infty$ . The following are true:

- (1)  $\|v\|_p \geq 0$
- (2)  $\|v + w\|_p \leq \|v\|_p + \|w\|_p$
- (3)  $\|\vec{0}\|_p = 0$
- (4)  $\|v\|_p = 0 \Rightarrow v = \vec{0}$
- (5)  $\forall t \in \mathbb{R}, \|tv\|_p = |t|\|v\|_p$

### Proofs

Let  $p = \infty$ . We will prove (2).

Say  $\|v\|_{\infty} = |x_i|$  and  $\|w\|_{\infty} = |y_k|$ . We want to show that  $\|v + w\|_{\infty} = \max_{j=1}^n |x_j + y_j| \leq |x_i| + |y_k|$ .

Note that  $\forall j$

$$\begin{aligned} |x_j + y_j| &\leq |x_j| + |y_j| && \text{Triangle Inequality} \\ &\leq |x_i| + |y_k| \\ &= \|v\|_{\infty} + \|w\|_{\infty} \end{aligned}$$

Therefore,  $\|v + w\|_{\infty} \leq \|v\|_{\infty} + \|w\|_{\infty}$ .

### Distances and Norms

A **norm** on  $\mathbb{R}^n$  is a function  $\|\cdot\| : \mathbb{R}^n \rightarrow \mathbb{R}^+, v \mapsto \|v\|$ , satisfying the following properties for  $v \in \mathbb{R}^n$ :

- (1)  $\|v\| \geq 0$
- (2)  $\|v + w\| \leq \|v\| + \|w\|$
- (3)  $\|\vec{0}\| = 0$
- (4)  $\|v\| = 0 \Rightarrow v = \vec{0}$
- (5)  $\forall t \in \mathbb{R}, \|tv\| = |t|\|v\|$

If  $\|\cdot\| : \mathbb{R}^n \rightarrow \mathbb{R}^+$  is a norm, we define  $d_{\|\cdot\|} : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}^+$ , defined as follows:

$$d_{\|\cdot\|}(v, w) = \|v - w\|$$

for  $v, w \in \mathbb{R}^n$ .

The properties of distance in  $\mathbb{R}$  still hold for distance in  $\mathbb{R}^n$ :

- (1)  $d(v, w) = d(w, v)$
- (2)  $d(u, w) \leq d(u, v) + d(v, w)$
- (3)  $d(v, v) = 0$
- (4)  $d(v, w) = 0 \Rightarrow v = w$

## Metric Spaces

A **metric space** is a nonempty set  $X$  equipped with a function  $d : X \times X \rightarrow \mathbb{R}^+$ ,  $(x, y) \mapsto d(x, y) \geq 0$ . The metric has the following properties:

- (1)  $d(x, y) = d(y, x) \quad \forall x, y \in X$
- (2)  $d(x, z) \leq d(x, y) + d(y, z) \quad \forall x, y, z \in X$
- (3)  $d(x, x) = 0$
- (4)  $d(x, y) = 0 \Leftrightarrow x = y$

The map  $d$  is called a *metric* on  $X$ .