This is going to be a loosely held-together collection of notes on different types of constructions in the field of operator algebras whose primary connection is that they're referred to by "product."

Tensor Products of C^*-Algebras

One of the main complications with C^*-algebras is that the algebraic tensor product $A \odot B$ may admit more than one C^*-norm.

Definition: A C^*-norm on $A \odot B$ is a norm $\|\cdot\|_\alpha$ satisfying $\|xy\|_\alpha \leq \|x\|_\alpha \|y\|_\alpha$ and $\|x^*x\|_\alpha = \|x\|_\alpha^2$ for all $x, y \in A \odot B$. The completion of $A \odot B$ with respect to this norm is denoted $A \otimes_\alpha B$.

However, the existence of there being multiple norms is not necessarily universal.

For instance, since there is a $*$-isomorphism between $\mathbb{M}_n(\mathbb{C}) \odot A$ and $\mathbb{M}_n(A)$, we have that $\mathbb{M}_n(\mathbb{C}) \odot A$ admits a unique C^*-norm, which is the norm that it obtains from $\mathbb{M}_n(A)$.

Definition: Let A and B are C^*-algebras. The *maximal norm* on $A \odot B$ is given by

$$\|x\|_{\max} = \sup\{\|\pi(x)\| \mid \pi\colon A \odot B \to B(H) \text{ a cyclic } *\text{-representation}\}$$

for any $x \in A \odot B$. The completion of $A \odot B$ with respect to $\|\cdot\|_{\max}$ is denoted $A \otimes_{\max} B$.

This is akin to the universal C^*-norm on the group C^*-algebra $C^*(\Gamma) \subseteq B(\ell_2(\Gamma))$.

Definition: If $\pi\colon A \to B(H)$ and $\sigma\colon B \to B(K)$ are faithful representations, then the *spatial* or *minimal* C^*-norm on $A \odot B$ is given by

$$\left\|\sum_{i=1}^{n} a_i \otimes b_i\right\|_{\min} = \left\|\sum_{i=1}^{n} \pi(a_i) \otimes \sigma(b_i)\right\|_{B(H \otimes K)}.$$

The completion of $A \odot B$ with respect to $\|\cdot\|_{\min}$ is denoted $A \otimes_{\min} B$.

Remark: If we consider von Neumann algebras $M \subseteq B(H)$ and $N \subseteq B(K)$, then there are numerous C^*-norms that $M \odot N$ can be endowed with, but their norm completions will not be von Neumann algebras. The von Neumann algebra tensor product is still well-defined, and denoted $M \bar{\otimes} N$, which is the von Neumann algebra generated by $M \otimes N \subseteq B(H \otimes K)$.

Free Products and Ultraproducts

Crossed Products

If we have a group Γ acting on a C^*-algebra A, then a question becomes how we may construct a C^*-algebra that encodes this action.

If a group is acting on another group, there is a construction known as the (outer) semidirect product that enables this. We will construct a C^*-algebra emerging from an action $\alpha\colon \Gamma \to \operatorname{aut}(A)$, which will be dubbed $A \rtimes_\alpha \Gamma$.