

These are notes I'm taking for Christian Gromoll's Probability Theory class. The primary text is *Probability with Martingales* by Williams.

Dictionary between Measure Theory and Probability

We will assume a lot of measure theory background, so we will instead discuss the dictionary between terms from probability and terms from measure theory. It's important to note that the measure spaces in probability theory are probability measure spaces (i.e., they have measure 1).

Measure Theory	Probability
Set X	Sample space Ω
σ -Algebra Σ	σ -Field $\mathcal{F}$
Measurable subsets $A \in \Sigma$	Events $F \in \mathcal{F}$
Measure $\mu(A)$	Probability $P(F)$
Almost Everywhere (a.e.)	Almost Surely (a.s.)
$\limsup_{n \rightarrow \infty} E_n$	$\{E_n \text{ infinitely often}\}, \{E_n \text{ i.o.}\}$
$\liminf_{n \rightarrow \infty} E_n$	$\{E_n \text{ eventually}\}, \{E_n \text{ ev}\}$
Borel Measurable Function $f: X \rightarrow \mathbb{R}$	Random Variable $X: \Omega \rightarrow \mathbb{R}$
Integral $\int f d\mu$	Expectation $E[X]$
Pushforward measure $f_*\mu$	Law Λ_X
$\mu(S)$	$P(\omega \in S)$
$f_*\mu(S')$	$P(X \in S')$

Remark: In probability, rather than the non-calligraphic L^p spaces (whose elements are equivalence classes of almost-everywhere equal functions), one uses the calligraphic $\mathcal{L}^p$, whose elements are the functions themselves. This does not fundamentally change anything in particular (as probabilists are only concerned with almost-everywhere equivalence regardless), but it is useful to note.

Important Theorems

σ -Algebras and Measures

Definition: Let S be a set, and let $\mathcal{D}$ be a collection of subsets. We call $\mathcal{D}$ a π -system on S if it is closed under finite intersection.

We call $\mathcal{D}$ a λ -system if

- $S \in \mathcal{D}$;
- for any $A, B \in \mathcal{D}$ with $A \subseteq B$, one has $B \setminus A \in \mathcal{D}$;
- if $(A_n)_n \subseteq \mathcal{D}$ is such that $\bigcup_n A_n = A$, then $A \in \mathcal{D}$.

For an arbitrary collection of subsets $\mathcal{I}$, we write $\pi(\mathcal{I})$ for the π -system generated by $\mathcal{I}$ and $\lambda(\mathcal{I})$ for the λ -system generated by $\mathcal{I}$.

Proposition: A collection Σ of subsets of S is a σ -algebra if and only if Σ is both a π -system and a λ -system.

Proof. Let Σ be a π -system and a λ -system. If $E, F \in \Sigma, (E_n)_n \subseteq \Sigma$, then

$$E^c = S \setminus E$$

$$E \cup F = S \setminus (E^c \cap F^c).$$

Therefore, $G_n = E_1 \cup \dots \cup E_n \in \Sigma$, with $\bigcup_n G_n = \bigcup_k E_k$, so that $\bigcup_k E_k \in \Sigma$. Finally, we have $\bigcap_k E_k = \left(\bigcup_k E_k^c\right)^c$, meaning that Σ is a σ -algebra. $\square$

Theorem (Dynkin π - λ Theorem): If $\mathcal{I}$ is a π -system, then $\lambda(\mathcal{I}) = \sigma(\mathcal{I})$.

Proof. To prove the theorem, we will use a classic technique by defining a “truth set” on which a desired statement holds, then showing that truth set is the desired σ -algebra.

Define

$$\mathcal{D}_1 = \{B \in \lambda(\mathcal{I}) : B \cap C \in \lambda(\mathcal{I}) \text{ for all } C \in \mathcal{I}\}.$$

We observe that $\mathcal{I}$ is a π system, so that $\mathcal{I} \subseteq \mathcal{D}_1$. Furthermore,

- $S \in \mathcal{D}_1$;
- for any $B_1, B_2 \in \mathcal{D}_1$ with $B_1 \subseteq B_2$, we have that for any given $C \in \mathcal{I}$, $(B_2 \setminus B_1) \cap C = (B_2 \cap C) \setminus (B_1 \cap C)$, so since $\lambda(\mathcal{I})$ is a λ -system, it follows that $B_2 \setminus B_1 \in \mathcal{D}_1$;
- if $(B_n)_n \subseteq \mathcal{D}_1$ increases to B , then for each $C \in \mathcal{I}$, one has $(B_n \cap C)_n \nearrow B \cap C$, so that $B \cap C \in \lambda(\mathcal{I})$, meaning $B \in \mathcal{D}_1$.

Thus, $\mathcal{D}_1$ is a λ -system containing $\mathcal{I}$, so since $\mathcal{D}_1 \subseteq \lambda(\mathcal{I})$ by definition, one has $\mathcal{D}_1 = \lambda(\mathcal{I})$.

Next, define

$$\mathcal{D}_2 = \{A \in \lambda(\mathcal{I}) : B \cap A \in \lambda(\mathcal{I}) \text{ for all } B \in \lambda(\mathcal{I})\}.$$

From above, we have that $\mathcal{D}_2$ contains $\mathcal{I}$. Similarly to above, we can prove that $\mathcal{D}_2$ inherits the λ -system structure from $\lambda(\mathcal{I})$, so that $\mathcal{D}_2 = \lambda(\mathcal{I})$. Yet, this means that $\mathcal{D}_2 = \lambda(\mathcal{I})$, so $\lambda(\mathcal{I})$ is a π -system. $\square$

Corollary: If two probability measures μ_1 and μ_2 agree on a π -system $\mathcal{I}$, then they agree on the σ -algebra $\sigma(\mathcal{I})$.

Theorem (Carathéodory Extension Theorem): Let S be a set, Σ_0 an algebra of subsets on S , and $\Sigma = \sigma(\Sigma_0)$. If $\mu_0: \Sigma_0 \rightarrow [0, \infty]$ is a countably additive map, then there exists a measure μ on (S, Σ) such that $\mu = \mu_0$ on Σ_0 .

Furthermore, if $\mu_0(S) < \infty$, then the extension is unique.

Convergence Theorems

We recall some of the useful convergence theorems from measure theory, specializing to the case with measure 1 and rewriting in the language of probability.

Theorem (Monotone Convergence Theorem): If $(X_n)_n$ is a sequence of real-valued random variables increasing pointwise almost surely to a real-valued random variable X , then

$$E[X] = \lim_{n \rightarrow \infty} E[X_n].$$

Theorem (Fatou’s Lemma): If $(X_n)_n$ is a sequence of positive-valued random variables, then

$$\begin{aligned} E\left[\liminf_{n \rightarrow \infty} X_n\right] &\leq \liminf_{n \rightarrow \infty} E[X_n] \\ \limsup_{n \rightarrow \infty} E[X_n] &\leq E\left[\limsup_{n \rightarrow \infty} X_n\right]. \end{aligned}$$

Theorem (Dominated Convergence Theorem): Let $(X_n)_n$ be a sequence of complex-valued random variables converging pointwise almost surely to a complex-valued random variable X . If there exists some positive-valued integrable random variable Y such that $|X_n| \leq Y$ for all n , then $(X_n)_n \xrightarrow{\mathcal{L}^1} X$. That is,

$$E[|X_n - X|] \rightarrow 0,$$

so that $E[X_n] \rightarrow E[X]$.

Theorem (Scheffé Convergence Theorem): If $(X_n)_n$ is a sequence of (real valued) integrable random variables converging to an integrable random variable X almost surely, and $E(|X_n|) \rightarrow E(|X|)$, then $(X_n)_n \xrightarrow{\mathcal{L}^1} X$.

Remark: The Scheffé convergence theorem is proven by splitting into positive and negative components.

Independence

Definition: We say sub- σ -algebras $\mathcal{G}_1, \mathcal{G}_2, \dots$ of $\mathcal{F}$ are *independent* if for any $G_i \in \mathcal{G}_i$ with distinct i , one has

$$P(G_{i_1} \cap \dots \cap G_{i_n}) = \prod_{k=1}^n P(G_{i_k}).$$

We say random variables $X_1, X_2, \dots$ are independent if their produced σ -algebras $\sigma(X_1), \sigma(X_2), \dots$ are independent.

Events $E_1, E_2, \dots$ are independent if their σ -algebras $\mathcal{E}_n$, where

$$\mathcal{E}_n = \{\emptyset, E_n, E_n^c, \Omega\},$$

are independent.

Theorem: If X and Y are independent random variables, then

$$E[XY] = E[X]E[Y].$$

Proof. Bootstrap from simple functions. □

Asymptotic Results

Theorem (First Borel–Cantelli Lemma): Let $(E_n)_n$ be a family of events such that $\sum_n P(E_n) < \infty$. Then,

$$\begin{aligned} P\left(\limsup_{n \rightarrow \infty} E_n\right) &= P(E_n \text{ i.o.}) \\ &= 0. \end{aligned}$$

Proof. Let

$$G_m = \bigcup_{n=m}^{\infty} E_n.$$

Then, $G_m \searrow \limsup E_n$, and one has

$$\begin{aligned} P(\limsup E_n) &\leq P(G_m) \\ &\leq \sum_{n=m}^{\infty} P(E_n). \end{aligned}$$

Taking $m \rightarrow \infty$ yields our desired result. □

Theorem (Second Borel–Cantelli Lemma): Suppose $(E_n)_n$ is a sequence of independent events. Then, if $\sum_n P(E_n) = \infty$, one has

$$P(\limsup E_n) = 1.$$

Proof. Recall that $(\limsup E_n)^c = \liminf E_n^c$. If $p_n = P(E_n)$, then one has

$$P\left(\bigcap_{n \geq m} E_n^c\right) = \prod_{n \geq m} (1 - p_n).$$

Notice that if $x \geq 0$, then $1 - x \leq e^{-x}$. Therefore, we have

$$\prod_{n \geq m} (1 - p_n) \leq \exp\left(-\sum_{n \geq m} p_n\right) = 0,$$

where the latter equality emerges from the fact that $\sum_n p_n = \infty$.

Therefore $P(\liminf E_n^c) = 0$, meaning $P(\limsup E_n) = 1$. □

Inequalities