

Contents

Locally Convex Topological Vector Spaces	1
Definitions and Examples	1
Metrizability and Normability	4
Hyperplane Separation and Continuous Linear Functionals	5
Weak Topologies and Extremal Structure	7
Duality	7
Duality in Subspaces and Quotient Spaces	8
Banach–Alaoglu Theorem and Applications	9
Separability and Metrizability	11
Extremal Structure	13
C^*-Algebras	14
Spectral Theory	14
Applications of the Spectral Theorem	22

Locally Convex Topological Vector Spaces

Definitions and Examples

Definition: A topological vector space (TVS) is a vector space X with a topology $\mathcal{T}$ such that, with respect to this topology, the maps

- (i) $X \times X \rightarrow X, (x, y) \mapsto x + y$
- (ii) $\mathbb{F} \times X \rightarrow X, (\alpha, x) \mapsto \alpha x$

are continuous with respect to $\mathcal{T}$.

Example: If $\mathcal{P}$ is a family of seminorms on X , and $\mathcal{T}$ is the topology on X with subbasic sets $U(x_0, p, \varepsilon) = \{x \in X : p(x - x_0) < \varepsilon\}$, then $(X, \mathcal{T})$ is a topological vector space.

Definition: A locally convex topological vector space (LCS) is a TVS equipped with a family of seminorms $\mathcal{P}$ that satisfy

$$\bigcap_{p \in \mathcal{P}} \{x : p(x) = 0\} = \{0\}.$$

That is, the family $\mathcal{P}$ separates the points of X .

Proposition: Let X be a TVS, and let p be a seminorm on X . The following are equivalent:

- (a) p is continuous;
- (b) $\{x \in X : p(x) < 1\}$ is open;
- (c) $0 \in \{x \in X : p(x) < 1\}^\circ$;
- (d) $0 \in \{x \in X : p(x) \leq 1\}^\circ$;
- (e) p is continuous at 0;
- (f) there is a continuous seminorm q on X such that $p \leq q$.

Proof. We see immediately that (a) implies (b) since the set is equal to the inverse image of the open subset $[0, 1) \subseteq [0, \infty)$. Similarly, if $p^{-1}([0, 1))$ is open, then it is equal to its own interior, and since $0 \in p^{-1}([0, 1))$, it follows that there is a subbasic open neighborhood of 0 contained in $p^{-1}([0, 1))$, giving (c). Similarly, we obtain (d) by the fact that $0 \in p^{-1}([0, 1))^\circ \subseteq p^{-1}([0, 1])^\circ$.

Next, we see that (d) implies that for every $\varepsilon > 0$, $0 \in p^{-1}([0, \varepsilon])^\circ$, so if $(x_i)_i$ is a net in X converging to 0 and $\varepsilon > 0$, there is some i_0 such that for all $i \geq i_0$, $x_i \in \{x : p(x) \leq \varepsilon\}$. That is, $p(x_i) \leq \varepsilon$ for all $i \geq i_0$, whence p is continuous at 0. This gives (e).

If p is continuous at 0, then if $x_i \rightarrow x$ is a convergent net in X , the reverse triangle inequality gives

$$|p(x) - p(x_i)| \leq p(x_i - x),$$

whence $p(x_i) \rightarrow p(x)$, so p is continuous. This gives that (e) implies (a).

Taking $q = p$ yields (a) implies (f). We will show that (f) implies (e). If $x_i \rightarrow 0$ in X , then $q(x_i) \rightarrow 0$, but since $0 \leq p(x_i) \leq q(x_i)$, it follows that $p(x_i) \rightarrow 0$. $\square$

Example:

- (a) Let X be a completely regular topological space, and let $C(X)$ be the set of all continuous functions from X to $\mathbb{F}$. If K is compact in X , we may define a seminorm $p_K(f) = \sup_{x \in K} |f(x)|$. Then,

$$\{p_K : K \subseteq X \text{ compact}\}$$

forms a separating family of seminorms for $C(X)$.

- (b) Let G be an open subset of $\mathbb{C}$. If $H(G)$ is the set of holomorphic functions on G , then the same seminorms $\{p_K : K \subseteq G \text{ compact}\}$ forms a locally compact space whose topology is uniform convergence on compact subsets.

- (c) Let X be a normed vector space. For each $\varphi \in X^*$, let $p_\varphi(x) = |\varphi(x)|$. Then, $\mathcal{P} = \{p_\varphi : \varphi \in X^*\}$ forms a separating family of seminorms for X . The topology these seminorms induce is called the *weak topology* on X . It is often denoted $\sigma(X, X^*)$.

- (d) If X is a normed space, then for each $x \in X$, we may define a seminorm $p_x : X^* \rightarrow [0, \infty)$ by taking $p_x(\varphi) = |\varphi(x)|$. This defines a separating family of seminorms $\mathcal{P} = \{p_x : x \in X\}$, and the topology on X^* induced by these seminorms is called the *weak**, or *w**, topology. It is often denoted $\sigma(X^*, X)$.

Next, we focus on an examination of the “convexity” part of the notion of locally convex topological vector spaces.

Definition: A subset $A \subseteq X$ is said to be convex if for all $v, w \in A$, the segment

$$[v, w] := \{tv + (1 - t)w : 0 \leq t \leq 1\}$$

is contained in A .

Note that the intersection of any family of convex subsets is convex.

Definition: If $A \subseteq X$ is any subset, then the *convex hull* of A , denoted $\text{conv}(A)$, is the intersection of all the convex subsets that contain A . If X is a TVS, the *closed convex hull* of A , denoted $\overline{\text{conv}}(A)$, is the intersection of all the closed convex sets that contain A .

Given any linear map $T : X \rightarrow Y$ of real vector spaces, and $C \subseteq Y$ a convex subset of Y , then $T^{-1}(C)$ is convex in X . If X^* is the dual of any complex vector space, then since X is also a real vector space, it then follows that the sets

$$A_0 = \{x : |f(x)| \leq 1\}$$

$$A_1 = \{x : \text{Re}(f(x)) \leq 1\}$$

$$A_3 = \{x : \text{Re}(f(x)) > 1\}$$

are all convex.

Proposition: Let X be a TVS, and let A be a convex subset of X . Then,

- (a) $\overline{A}$ is convex;

(b) if $a \in A^\circ$ and $b \in \overline{A}$, then

$$[a, b] := \{tb + (1-t)a : 0 \leq t < 1\}$$

is contained in A° .

Proof. Let $a \in A$, $b \in \overline{A}$, and let $0 \leq t \leq 1$. Let $(x_i)_i$ be a net in A such that $x_i \rightarrow b$. Then, $tx_i + (1-t)a \rightarrow tb + (1-t)a$. In particular, this means that if $b \in \overline{A}$ and $a \in A$, then $[a, b] \subseteq \overline{A}$. So, given any $a, b \in \overline{A}$, we have that the net $(x_i)_i \rightarrow b$ is such that $[x_i, a] \subseteq \overline{A}$, so it follows that $[b, a] \subseteq \overline{A}$, whence $\overline{A}$ is convex.

For (b), fix $0 < t < 1$, and set $c = tb + (1-t)a$ for some $a \in A^\circ$ and $b \in \overline{A}$. Since $a \in A^\circ$, there is an open neighborhood V of 0 such that $a + V \subseteq A$. Therefore, for any $d \in A$, we have

$$\begin{aligned} A &\supseteq td + (1-t)(a + V) \\ &= t(d - b) + tb + (1-t)(a + V) \\ &= (t(d - b) + (1-t)V) + c. \end{aligned}$$

We hope to find some $d \in A$ such that $0 \in t(d - b) + (1-t)V$, which we call U . The preceding inclusion then shows that $c \in A^\circ$, since U is an open neighborhood of 0 with $c + U \subseteq A$.

Finding such a d is equivalent to finding d such that $0 \in \frac{1}{t}(1-t)V + (d - b)$, or $d \in b - \frac{1}{t}(1-t)V$. Yet, we know that $0 \in -\frac{1}{t}(1-t)V$ by the assumption that V is an open neighborhood of 0. Since $b \in \overline{A}$, it follows that $b - \frac{1}{t}(1-t)V \cap A \neq \emptyset$ by the definition of closure, whence such a $d \in A$ exists. $\square$

Corollary: If $A \subseteq X$, then $\overline{\text{conv}(A)} = \overline{\text{conv}(\overline{A})}$.

Definition: A set $A \subseteq X$ is called *balanced* if for any $|\alpha| \leq 1$ and any $x \in A$, we have $\alpha x \in A$.

We say a set A is *absorbing* if for any $x \in X$, there is some $\varepsilon > 0$ such that $tx \in A$ for all $0 \leq t < \varepsilon$. We say A is *absorbing at a* if for any $x \in X$, there is $\varepsilon > 0$ such that $a + tx \in A$ for all $0 \leq t < \varepsilon$.

Given a seminorm p on V , the set $V = \{x : p(x) < 1\}$ is a convex balanced set that is absorbing at each of its points. We will show that the converse actually holds — that is, a vector space is a LCS if and only if the set of all open, convex, balanced subsets of X forms a neighborhood basis at 0.

Proposition: Suppose X is a vector space over F , and V is a nonempty convex balanced set that is absorbing at each of its points. Then, there is a unique seminorm p on X such that

$$V = \{x \in X : p(x) < 1\}.$$

Proof. Let

$$p(x) = \inf\{t \geq 0 : x \in tV\}.$$

Since V is absorbing, it follows that $X = \bigcup_{n=1}^{\infty} nV$, so this is a well-defined quantity. Furthermore, $p(0) = 0$.

To show that $p(\alpha x) = |\alpha|p(x)$, we suppose $\alpha \neq 0$, and since V is balanced,

$$\begin{aligned} p(\alpha x) &= \inf\{t \geq 0 : \alpha x \in tV\} \\ &= \inf\left\{t \geq 0 : x \in \frac{t}{\alpha}V\right\} \\ &= \inf\left\{t \geq 0 : x \in \frac{t}{|\alpha|}V\right\} \\ &= |\alpha| \inf\left\{\frac{t}{|\alpha|} : x \in \frac{t}{|\alpha|}V\right\} \\ &= |\alpha|p(x). \end{aligned}$$

We now observe that if $\alpha, \beta \geq 0$, then for any $a, b \in V$, we have

$$\begin{aligned}\alpha a + \beta b &= (\alpha + \beta) \left(\frac{\alpha}{\alpha + \beta} a + \frac{\beta}{\alpha + \beta} b \right) \\ &\in (\alpha + \beta)V,\end{aligned}$$

since V is convex. For any $x, y \in X$, take $p(x) = \alpha$ and $p(y) = \beta$, and let $\delta > 0$. Then, $x \in (\alpha + \delta)V$ and $y \in (\beta + \delta)V$ by the definition of the infimum, so that $x + y \in (\alpha + \beta + 2\delta)V$. Therefore, we have $p(x + y) \leq \alpha + \beta + 2\delta$, which holds for all $\delta > 0$, so that $p(x + y) \leq p(x) + p(y)$.

Finally, we will have to show that $V = \{x : p(x) < 1\}$. If $p(x) = \alpha < 1$, then for any $\alpha < \beta < 1$, we have $x \in \beta V \subseteq V$, which emerges from the fact that V is balanced. Thus, $V \supseteq \{x : p(x) < 1\}$. If $x \in V$, then $p(x) \leq 1$, and since V is absorbing at x , there is some $\varepsilon > 0$ such that for $0 < t < \varepsilon$, we have $y = x + tx \in V$. Yet, this means $x = \frac{1}{1+t}y$ with $y \in V$, so that $p(x) = \frac{1}{1+t}p(y) \leq \frac{1}{1+t} < 1$. $\square$

The seminorm p is called the Minkowski function for V , or the *gauge* of V .

Proposition: Let X be a TVS, and let $\mathcal{U}$ be the collection of open, convex, balanced subsets of X . Then, X is locally convex if and only if $\mathcal{U}$ is a basis for the neighborhood system at 0.

Proof. Suppose X is locally convex with topology defined by the family of seminorms $\mathcal{P}$. Then, we observe that each of the subbasic open neighborhoods of 0,

$$\{U(0, p, \varepsilon) : \varepsilon > 0, p \in \mathcal{P}\},$$

are open, convex, and balanced; since the set of all finite intersections of all these subbasic open neighborhoods yields a neighborhood basis for 0, and finite intersections of open, balanced, convex subsets are open, balanced, and convex, so it follows that $\mathcal{U}$ contains the set of all finite intersections of all these subbasic open neighborhoods, whence $\mathcal{U}$ contains a basis for the neighborhood system at 0. In particular, this means $\mathcal{U}$ also must generate the neighborhood system at 0, so $\mathcal{U}$ forms a basis for the neighborhood system at 0.

Now, suppose $\mathcal{U}$ forms a basis for the neighborhood system at 0. Observe that an open, convex, balanced subset of V contains 0, and is itself absorbing. We claim that the family of Minkowski functionals

$$\mathcal{P} = \{p_V(\cdot - z) : z \in X, V \in \mathcal{U}\}$$

generates the topology for X . $\square$

Thus, it follows that in a locally convex space, a net $(x_i)_i$ converges in the topology to x if and only if for all seminorms $p \in \mathcal{P}$, we have $p(x_i - x) \rightarrow 0$.

Example:

- (i) Suppose X is a locally compact Hausdorff space (or any Tychonoff space, but we will work with these for uniformity), and let $C(X)$ be the space of all continuous functions on X . Then, the family of seminorms $p_K(f) = \sup_{x \in K} |f(x)|$ yields a locally convex space where convergence is defined as uniform convergence on compact subsets.
- (ii) If X is a normed space, then the family of seminorms $p_\varphi(x) = |\varphi(x)|$ for each $\varphi \in X^*$ defines the weak topology on X .
- (iii) If X is a normed space, then the family of seminorms $p_x(\varphi) = |\varphi(x)|$ for each $x \in X$ defines the weak*, or w^* , topology on X^* .

The weak and weak* topologies are very important, and we will discuss them more in depth.

Metrizability and Normability

When we look at a locally convex space, we concern ourselves with whether the topology of the space is given by a metric or not. It turns out this is (effectively) equivalent to there being countably many seminorms defining the topology.

Proposition: Let $\{p_n : n \geq 1\}$ be a sequence of separating seminorms on X . For any $x, y \in X$, we may take

$$d(x, y) = \sum_{n=1}^{\infty} 2^{-n} \frac{p_n(x-y)}{1 + p_n(x-y)}.$$

Then, the topology on X is defined by this metric.

Definition: A Fréchet space is a locally convex space whose topology is defined by a metric that is complete.

Definition: If X is a topological vector space, and $B \subseteq X$, then we say B is bounded if, for every open set U containing 0 , there is some $\varepsilon > 0$ such that $\varepsilon B \subseteq U$.

Proposition: If X is a locally convex space, then X is normable if and only if X has a nonempty bounded open set.

Proof. The open unit ball U_X is bounded. Therefore, we only need to show one direction. Suppose X is a locally convex space with a bounded open set U . Translating if necessary, we may assume $0 \in U$.

Since X is locally convex, there is an open, convex, balanced subset of U which we call V ; using the gauge of V , we may find a continuous seminorm p such that $\{x : p(x) < 1\} = V$. We will show that p is a norm and that p defines the topology on X .

First, let $0 \neq x \in X$. Let W_0 and W_x be disjoint open subsets with $0 \in W_0$ and $x \in W_x$. Since W_0 is absorbing, there is some $\varepsilon > 0$ such that $W_0 \supseteq \varepsilon U \supseteq \varepsilon V$. We note that $\varepsilon V = \{y : p(y) < \varepsilon\}$. Since $x \notin W_0$, we have $p(x) \geq \varepsilon$, so that p is a norm.

To show that the topology on X is generated by p , it suffices to show that if q is any continuous seminorm, then there is some α such that $q \leq \alpha p$. Since q is continuous, it follows that there is some $\varepsilon > 0$ such that $\{x : q(x) < 1\} \supseteq \varepsilon U \supseteq \varepsilon V$. Yet, this means that if $p(x) < \varepsilon$, then $q(x) < 1$, whence $q \leq \varepsilon^{-1} p$. $\square$

Hyperplane Separation and Continuous Linear Functionals

We recall the classification of continuous linear functionals in a locally convex topological vector space.

Theorem: If X is a locally convex topological vector space with defining seminorms $\mathcal{P}$, and $f : X \rightarrow \mathbb{C}$ is a linear functional, then the following statements are equivalent:

- (i) f is continuous;
- (ii) f is continuous at 0 ;
- (iii) f is continuous at some point;
- (iv) $\ker(f)$ is closed;
- (v) there are $p_1, \dots, p_n \in \mathcal{P}$ and some constant C such that $|f(x)| \leq C \max\{p_1(x), \dots, p_n(x)\}$.

To understand hyperplane separation, we start with the most basic case — namely, that of separating a convex subset from the origin via a continuous linear functional.

Theorem: If X is a topological vector space, and G is an open convex nonempty subset of X that does not contain the origin, then there is a closed hyperplane M such that $M \cap G = \emptyset$.

Proof. We start by assuming that X is an $\mathbb{R}$ -linear space. Choose $x_0 \in G$, and let $H = x_0 - G$. Then, H is an open, convex subset of X that contains 0 . Taking the gauge of H yields a continuous sublinear functional $q : X \rightarrow \mathbb{R}$ with $H = \{x : q(x) < 1\}$. Furthermore, since $x_0 \notin H$, we have that $q(x_0) \geq 1$.

Consider the subspace $Y = \{\alpha x_0 : \alpha \in \mathbb{R}\}$. Let $f_0 : Y \rightarrow \mathbb{R}$ be defined by taking $f_0(\alpha x_0) = \alpha q(x_0)$. Observe that if $\alpha \geq 0$, then $f_0(\alpha x_0) = q(\alpha x_0)$, and if $\alpha < 0$, then $f_0(\alpha x_0) = \alpha q(x_0) \leq \alpha < 0 \leq q(\alpha x_0)$, whence $f_0 \leq q$ on Y . Using the Hahn–Banach–Minkowski extension yields some $f : X \rightarrow \mathbb{R}$ a continuous linear functional such that $f|_Y = f_0$ and $f \leq q$ on X . Then, $M = \ker(f)$ is the desired hyperplane.

If $x \in G$, then $x_0 - x \in H$, so $f(x_0) - f(x) = f(x_0 - x) \leq q(x_0 - x) < 1$, while $f(x) > f(x_0) - 1 = q(x_0) - 1 \geq 0$ for all $x \in G$, whence $M \cap G = \emptyset$.

If X is a $\mathbb{C}$ -linear space, then since X is also an $\mathbb{R}$ -linear space, we obtain a continuous $\mathbb{R}$ -linear functional $f: X \rightarrow \mathbb{R}$ with $G \cap \ker(f) = \emptyset$. If $F(x) = f(x) - if(ix)$, then F is a $\mathbb{C}$ -linear functional with $f = \operatorname{Re}(F)$ with $F(x) = 0$ if and only if $f(x) = f(ix) = 0$, so that $M = \ker(F)$ is the desired closed hyperplane. $\square$

Affine hyperplanes in X are given by $M' = M + x_0$ for some $x_0 \in X$ and hyperplane M . It turns out that we can extend any separating affine (closed) subspace of a topological vector space to an affine hyperplane.

Corollary: Let X be a topological vector space, and let G be an open, convex, nonempty subset of X . If Y is an affine subspace of X with $Y \cap G = \emptyset$, then there is a closed affine hyperplane M in X such that $Y \subseteq M$ and $M \cap G = \emptyset$.

Proof. Choosing some $x_0 \in Y$, we may consider $G - x_0$ and $Y - x_0$, so that Y may be assumed to be a subspace of X . Let $Q: X \rightarrow X/Y$ be the quotient map. Notice that $Q^{-1}(Q(G)) = \{y + G : y \in Y\}$, so that $Q(G)$ is open in X/Y . Furthermore, $Q(G)$ is also convex since Q is a linear map. Furthermore, since $Y \cap G = \emptyset$, $0 \notin Q(G)$. In particular, there is then a closed hyperplane N in X/Y such that $N \cap Q(G) = \emptyset$. Setting $M = Q^{-1}(N)$, we have that M is our desired hyperplane. $\square$

In the special case of a real topological vector space, it's useful to note that $X \setminus \ker(f)$ for a continuous linear functional $f: X \rightarrow \mathbb{R}$ has two connected components. This yields the following notion.

Definition: Let X be a real topological vector space. A subset $S \subseteq X$ is called an open half space if there is some continuous linear functional $f: X \rightarrow \mathbb{R}$ such that $S = f^{-1}((\alpha, \infty))$. We call S a closed half-space if there is a continuous linear functional $f: X \rightarrow \mathbb{R}$ such that $S = f^{-1}([\alpha, \infty))$.

We say A and B are strictly separated if they are contained in disjoint open half-spaces, while they are separated if they are contained in two closed half-spaces whose intersection is of the form $M = f^{-1}(\{\alpha\})$. That is, their intersection is a closed affine hyperplane. Notice also that continuous linear functionals are open maps, so that the interior of a closed half-space is an open half-space, and the closure of an open half-space is a closed half-space.

Theorem (General Hyperplane Separation): Let X be a real topological vector space, and A and B disjoint convex subsets with A open. Then, there is a continuous linear functional $f: X \rightarrow \mathbb{R}$ and a real scalar α such that $A \subseteq f^{-1}((-\infty, \alpha))$ and $B \subseteq f^{-1}([\alpha, \infty))$.

Proof. Let $G = A - B$. Then, G is convex, and $G = \bigcup_{b \in B} (A - b)$, meaning G is open. Since $A \cap B = \emptyset$, it follows that $0 \notin G$. Thus, there is a closed hyperplane M in X such that $M \cap G = \emptyset$.

Let $f: X \rightarrow \mathbb{R}$ be a linear functional such that $M = \ker(f)$. Notice that $f(G)$ is a convex subset of $\mathbb{R}$ with $0 \notin f(G)$. Either, $f(x) > 0$ for all $x \in G$ or $f(x) < 0$ for all $x \in G$. Without loss of generality, we may assume that $f(x) > 0$ for all $x \in G$, so that $f(b) < f(a)$ for any $a \in A$ and $b \in B$, meaning there is some real number α such that $\sup_{b \in B} f(b) \leq \alpha \leq \inf_{a \in A} f(a)$. Since $f(A)$ and $f(B)$ are open intervals if B is open, it follows that $f < \alpha$ on B and $f > \alpha$ on A . $\square$

Lemma: If X is a topological vector space, K is a compact subset of X , and V is an open subset satisfying $K \subseteq V$, then there is an open neighborhood U of 0 such that $K + U \subseteq V$.

Proof. Let $\mathcal{O}_0$ denote the system of open neighborhoods of 0 . Suppose toward contradiction that for each $U \in \mathcal{O}_0$ that $K + U$ is not contained in V . We may choose vectors $x_U \in K$ and $y_U \in U$ such that $x_U + y_U \in V^c$. This yields a net $(y_U)_{U \in \mathcal{O}_0}$ and $(x_U)_{U \in \mathcal{O}_0}$ with $y_U \rightarrow 0$ and a subnet $x_{U'} \rightarrow x \in K$. Yet, this yields $(x_{U'} + y_{U'}) \rightarrow x$, implying that $x \in \overline{V^c} = V^c$, which yields a contradiction. $\square$

Theorem (Hyperplane Separation for Locally Convex Spaces): Let X be a real locally convex topological vector space, and let A and B be two disjoint closed convex subsets of X with B compact. Then, A and B are contained in disjoint open half-spaces.

Proof. Notice that B is a compact subset of A^c . Therefore, there is an open neighborhood U_1 of 0 such that $B + U_1 \subseteq A^c$. Since X is locally convex, we may find a continuous seminorm p on X such that $\{x : p(x) < 1\} \subseteq U_1$.

Define $U = \{x : p(x) < 1/2\}$. Then, $(B + U) \cap (A + U) = \emptyset$, and $A + U$ and $B + U$ are open convex subsets of X that contain A and B . Therefore, the result follows from the general hyperplane separation. $\square$

Rephrasing this theorem for locally convex spaces by using the fact that any linear functional φ over $\mathbb{C}$ is given by $\varphi(x) = \operatorname{Re}(\varphi(x)) - i \operatorname{Re}(\varphi(ix))$ yields the following.

Theorem (Hyperplane Separation for Locally Convex Spaces, Encore): Let X be a locally convex space, and let A and B be disjoint closed convex subsets of X with B compact. Then, there is a continuous linear functional $f: X \rightarrow \mathbb{C}$, some $\alpha \in \mathbb{R}$, and some $\varepsilon > 0$ such that for all $a \in A$ and $b \in B$, we have

$$\operatorname{Re}(f(a)) \leq \alpha < \alpha + \varepsilon \leq \operatorname{Re}(f(b)).$$

We will now discuss the dual space of one particular locally convex space.

Proposition: Let X be a locally compact Hausdorff space, and let $C(X)$ be topologized by uniform convergence on compact subsets. If $L: C(X) \rightarrow \mathbb{F}$ is a continuous linear functional, then there is a compact set K and a regular (complex) Borel measure μ on K such that $L(f) = \int_K f d\mu$ for any $f \in C(X)$. Conversely, all these measures define elements of $C(X)^*$.

Proof. Any compactly support measure defines an element of $C(X)^*$, and satisfies $|L(f)| \leq \|\mu\| \sup_{x \in K} |f(x)|$ for any such measure μ .

Conversely, let $L \in C(X)^*$. By the classification of continuous linear functionals, there are compact sets $K_1, \dots, K_n$ and a constant C such that $|L(f)| \leq C \max\{p_{K_j}(f) : j = 1, \dots, n\}$. Let $K = \bigcup_{j=1}^n K_j$. Then, $|L(f)| \leq \alpha p_K(f)$. In particular, for any $f \in C(X)$ with $f|_K = 0$, then $L(f) = 0$.

Define a linear functional $F: C(K) \rightarrow \mathbb{F}$ as follows. For any $g \in C(K)$, let $\tilde{g}$ be a continuous extension to X , and set $F(g) = L(\tilde{g})$. Suppose $\tilde{g}_1$ and $\tilde{g}_2$ are any such extensions; then, $\tilde{g}_1 - \tilde{g}_2 = 0$ on K , so that $L(\tilde{g}_1) = L(\tilde{g}_2)$, so that F is well-defined. Furthermore, F is linear. Since, for any $g \in C(K)$ and extension $\tilde{g} \in C(X)$, we have $|F(g)| \leq C p_K(g) = C \|g\|$. In particular, this means F is continuous, so there is some regular complex Borel measure μ in $M(K)$ such that $F(g) = \int g d\mu$. Notice that if $f \in C(X)$ is such that $f|_K = g \in C(K)$, it follows that $L(f) = F(g) = \int f d\mu$. $\square$

Weak Topologies and Extremal Structure

Duality

Let X be a locally convex space. Then, X^* denotes the space of continuous linear functionals on X , and admits a vector space structure. We will write $\langle x, \varphi \rangle = \varphi(x)$ whenever $x \in X$ and $\varphi \in X^*$. We will write $\hat{x} = \langle x, \cdot \rangle$ to denote the analogous linear functional on φ such that $\hat{x}(\varphi) = \langle x, \varphi \rangle$.

Definition: If X is a locally convex space, then the weak topology on X is the topology defined by the family of seminorms

$$p_\varphi(x) = |\langle x, \varphi \rangle|.$$

The weak* topology on X is the topology defined by the seminorms

$$p_x(\varphi) = |\langle x, \varphi \rangle|.$$

We have that $(x_i)_i \xrightarrow{\text{wk}} x$ if, for all $\varphi \in X^*$, $\varphi(x_i) \rightarrow \varphi(x)$. Similarly, we have that $(\varphi_i)_i \xrightarrow{\text{wk}^*} \varphi$ if, for all $x \in X$, $\varphi_i(x) \rightarrow \varphi(x)$.

Theorem: If X is a locally convex space, then $(X^*, \text{wk}^*)^* = X$.

Proof. First, if $x \in X$, then $\varphi \mapsto \langle x, \varphi \rangle$ is a weak*-continuous linear functional on X^* by definition.

Conversely, if $f \in (X^*, \text{wk}^*)^*$, then there are vectors $x_1, \dots, x_n \in X$ such that

$$|f(\varphi)| \leq \sum_{k=1}^n |\langle x_k, \varphi \rangle|$$

for all $\varphi \in X^*$. This implies that

$$\bigcap_{k=1}^n \ker(\hat{x}_k) \subseteq \ker(f),$$

so there are $\alpha_1, \dots, \alpha_n$ such that $f = \sum_{k=1}^n \alpha_k \hat{x}_k$, meaning that $f \in X$. $\square$

Local convexity is a very useful property to have, as the following result exhibits.

Theorem: Let X be a locally convex space, and let A be a convex subset of X . Then, $\overline{A} = \overline{A}^{\text{wk}}$.

Proof. Notice that the weak topology is weaker than the topology on X , so $\overline{A} \subseteq \overline{A}^{\text{wk}}$.

Conversely, suppose there were $x \in \overline{A}^{\text{wk}} \setminus \overline{A}$. Then, by geometric Hahn–Banach, there is $\varphi \in X^*$, $\alpha \in \mathbb{R}$, and $\varepsilon > 0$ such that

$$\operatorname{Re}(\langle a, \varphi \rangle) \leq \alpha < \alpha + \varepsilon \leq \operatorname{Re}(\langle x, \varphi \rangle)$$

for all $a \in \overline{A}$. Yet, this means that $\overline{A} \subseteq \{y \in X : \operatorname{Re}(\langle y, \varphi \rangle) \leq \alpha\}$, even though the latter is weakly closed. Since $x \notin \overline{A}$, it follows that $x \notin \overline{A}$, which yields a contradiction. $\square$

Corollary: A convex subset of X is closed if and only if it is weakly closed.

Duality in Subspaces and Quotient Spaces

Proposition: Let p be a seminorm on X , M a subspace of X , and $\overline{p}: X/M \rightarrow [0, \infty)$ is defined by

$$\overline{p}(x + M) = \inf\{p(x + y) : y \in M\},$$

then $\overline{p}$ is a seminorm on X/M . If $\mathcal{P}$ is a family of continuous seminorms on X defining a locally convex topology, then $\overline{\mathcal{P}}$ defines the quotient topology on X/M .

In particular, if X is a locally convex space and $M \leq X$ is a subspace, then X/M is a locally convex space. Suppose $f \in (X/M)^*$. Then, $f \circ \pi \in X^*$, where $\pi: X \rightarrow X/M$ is the projection. Furthermore, observe that $f \circ \pi$ is an element of

$$M^\perp = \{\varphi \in X^* : \varphi(m) = 0 \text{ for all } m \in M\}.$$

In particular, we obtain the map $f \mapsto f \circ \pi$, yielding a linear map $(X/M)^* \rightarrow M^\perp \subseteq X^*$.

Theorem: Let X be a locally convex space, $M \leq X$ a subspace, and $\pi: X \rightarrow X/M$ the natural map. Then, $f \mapsto f \circ \pi$ defines a bijection between $(X/M)^*$ and $M^\perp$.

Equipping $(X/M)^*$ with the weak* topology, and $M^\perp$ with the relative weak* topology, yields that this bijection is a homeomorphism. If X is a normed space, then the bijection is an isometry and a weak*–weak* homeomorphism.

Proof. Define $\rho: (X/M)^* \rightarrow M^\perp$ by $\rho(f) = f \circ \pi$. Then, we have that ρ is well-defined and takes $(X/M)^*$ to $M^\perp$. Notice that if $0\rho(f)$, then $f \circ \pi = 0$, but since π is surjective, f must be the zero map, whence ρ is injective.

Let $\varphi \in M^\perp$, and let $f: X/M \rightarrow \mathbb{C}$ be given by $f(x + M) = \langle x, \varphi \rangle$. Since $M \subseteq \ker(\varphi)$, f is well-defined and linear. Since

$$\pi^{-1}(\{x + M : |f(x + M)| < 1\}) = \{x \in X : |\langle x, \varphi \rangle| < 1\}$$

is open in X , it follows that $\{x + M : |f(x + M)| < 1\}$ is open in X/M , so f is continuous.

If X is a normed space, then results from Banach space theory yields that ρ is an isometry. If $(f_i)_i \subseteq (X/M)^*$ converges to 0 in the weak* topology on $(X/M)^*$, then for each $x \in X$, we have that

$$\langle x, \rho(f_i) \rangle = f_i(\pi(x))$$

$$\rightarrow 0,$$

so that $\rho(f_i) \rightarrow 0$ weak* in X^* . Conversely, if $\rho(f_i) \rightarrow 0$ weak* in X^* , it follows that for each $x \in X$, we have $f_i(x + M) = \langle x, \rho(f_i) \rangle \rightarrow 0$ weak* in $(X/M)^*$. $\square$

Banach–Alaoglu Theorem and Applications

Theorem: If X is a normed space, then the unit ball B_{X^*} of X^* is weak*-compact.

Proof. Let $D_x = \{\alpha \in \mathbb{C} : |\alpha| \leq 1\}$. Set $D = \prod_{x \in B_X} D_x$. By Tychonoff's theorem, D is compact. Let $\tau: B_{X^*} \rightarrow D$ be defined by

$$\tau(\varphi)(x) = \langle x, \varphi \rangle.$$

Then, $\tau(\varphi) \in D$ with x coordinate given by $\langle x, \varphi \rangle$. We will show that τ is a homeomorphism from B_{X^*} with the weak* topology onto $\tau(B_{X^*}) \subseteq D$ with the subspace topology, and that $\tau(B_{X^*}) \subseteq D$ is closed.

Suppose $\tau(\varphi_1) = \tau(\varphi_2)$. Then, for each $x \in B_X$, $\langle x, \varphi_1 \rangle = \langle x, \varphi_2 \rangle$. Since B_X separates the points of X^* , it follows that $\varphi_1 = \varphi_2$.

Let $(\varphi_i)_i$ be a net in B_{X^*} with $\varphi_i \xrightarrow{\text{wk}^*} \varphi$. For each $x \in B_X$, it follows that $\tau(\varphi_i)(x) \rightarrow \tau(\varphi)(x)$, or that each coordinate of $(\tau(\varphi_i))_i$ converges to $\tau(\varphi)$, so that $\tau(\varphi_i) \rightarrow \tau(\varphi)$, meaning τ is continuous.

Let $(\varphi_i)_i$ be a net in B_{X^*} , $f \in D$, and $\tau(\varphi_i) \rightarrow f$ in D . Then, $f(x) = \lim \langle x, \varphi_i \rangle$ exists for each $x \in B_X$. If $x \in X$, we choose $\alpha > 0$ such that $\|\alpha x\| \leq 1$. Take $f(x) = \alpha^{-1} f(\alpha x)$.

If $\beta > 0$ is also such that $\|\beta x\| \leq 1$, then

$$\begin{aligned} \alpha^{-1} f(\alpha x) &= \alpha^{-1} \lim \langle \alpha x, \varphi_i \rangle \\ &= \beta^{-1} \lim \langle \beta x, \varphi_i \rangle \\ &= \beta^{-1} f(\beta x), \end{aligned}$$

so that $f(x)$ is well-defined. Notice that by the way f is defined, it is necessarily a linear functional. If $\|x\| \leq 1$, then $f(x) \in D_x$, so $|f(x)| \leq 1$, meaning that there is $\varphi \in B_{X^*}$ such that $\tau(\varphi) = f$. Thus, $\tau(B_{X^*})$ is closed in D , so $\tau(B_{X^*})$ is compact. $\square$

Recall that a Banach space X is called reflexive if the embedding of X into its double dual X^{**} is surjective, where we recall that $\hat{x} \in X^{**}$ is defined by

$$\langle \varphi, \hat{x} \rangle = \langle x, \varphi \rangle.$$

We write $\iota: X \mapsto X^{**}$, taking $x \mapsto \hat{x}$, for this canonical embedding.

We will usually use σ -notation for weak topologies. That is, if X is a locally convex space, and X^* is the continuous dual space, then

- $\sigma(X, X^*)$ denotes the weak topology on X induced by X^* ;
- $\sigma(X^*, X)$ denotes the weak* topology on X^* induced by X .

Recall that convergence of nets in these topologies is given by

- $(x_i)_i \xrightarrow{\sigma(X, X^*)} x$ if and only if for all $\varphi \in X^*$, $\langle x_i, \varphi \rangle \rightarrow \langle x, \varphi \rangle$;
- $(\varphi_i)_i \xrightarrow{\sigma(X^*, X)} \varphi$ if and only if for all $x \in X$, $\langle x, \varphi_i \rangle \rightarrow \langle x, \varphi \rangle$ if and only if for all $x \in X$, $\langle \varphi_i, \hat{x} \rangle \rightarrow \langle \varphi, \hat{x} \rangle$.

Theorem (Goldstine's Theorem): Let X be a normed space. Then, $\iota(B_X)$ is $\sigma(X^{**}, X^*)$ -dense in X^{**} .

Proof. We will denote by w^{**} the weak* topology on X^{**} induced by X^* . Let $B = \overline{\iota(B_X)}^{w^{**}}$, and notice that $B \subseteq B_{X^{**}}$. Suppose there is some $\theta \in B_{X^{**}} \setminus B$. The Hahn–Banach theorem implies there is some $\varphi \in X^*$, $\alpha \in \mathbb{R}$,

and $\varepsilon > 0$ such that

$$\operatorname{Re}(\langle x, \varphi \rangle) < \alpha < \alpha + \varepsilon < \operatorname{Re}(\langle \varphi, \theta \rangle)$$

for all $x \in B_X$. Since $0 \in B_X$, it follows that $0 < \alpha$. Dividing, we may assume that there is $\varphi \in X^*$ and $\varepsilon > 0$ such that

$$\operatorname{Re}(\langle x, \varphi \rangle) < 1 < 1 + \varepsilon < \operatorname{Re}(\langle \varphi, \theta \rangle).$$

Notice that $|\langle x, \varphi \rangle| \leq 1$ (multiplying by a phase) for any $\|x\| \leq 1$, so that $\varphi \in B_{X^*}$. Yet, we also have

$$\begin{aligned} 1 + \varepsilon &< \operatorname{Re}(\langle \varphi, \theta \rangle) \\ &\leq |\langle \varphi, \theta \rangle| \\ &\leq \|\theta\| \\ &\leq 1, \end{aligned}$$

which is a contradiction. $\square$

Theorem: Let X be a Banach space. The following are equivalent:

- (a) X is reflexive;
- (b) X^* is reflexive;
- (c) we have $\sigma(X^*, X) = \sigma(X^*, X^{**})$;
- (d) B_X is weakly compact.

Proof. If X is reflexive, then we have that $(\varphi_i)_i \rightarrow \varphi$ weak* if and only if $\langle x, \varphi_i \rangle \rightarrow \langle x, \varphi \rangle$ for all $x \in X$ if and only if $\langle \varphi_i, \hat{x} \rangle \rightarrow \langle \varphi, \hat{x} \rangle$ for all $x \in X$ if and only if $\langle \varphi_i, \theta \rangle \rightarrow \langle \varphi, \theta \rangle$ for all $\theta \in X^{**}$. This gives (a) implies (c).

If the unit ball B_X of X is weakly compact, then it is the case that $\iota(B_X) \subseteq B_{X^{**}}$ is $\sigma(X^{**}, X^*)$ -closed. Yet, Goldstine's theorem gives that $\iota(B_X)$ is dense in $B_{X^{**}}$, so that $\iota(B_X) = B_{X^{**}}$, whence $\iota(X) = X^{**}$. This gives (d) implies (a).

By the Banach–Alaoglu theorem, B_{X^*} is $\sigma(X^*, X)$ -compact. By the assumption in (c), we have $\sigma(X^*, X^{**}) = \sigma(X^*, X)$, so B_{X^*} is $\sigma(X^*, X^{**})$ -compact. Therefore, since (d) implies (a), it follows that X^* is reflexive, giving (b).

We assume (b). The unit ball B_X is norm-closed in X^{**} . Therefore, since B_X is convex, it follows that B_X is $\sigma(X^{**}, X^{***})$ -closed. Since $X^* \cong X^{***}$, it follows that $\iota(B_X)$ is $\sigma(X^{**}, X^*)$ -closed. By Goldstine's theorem, we know that B_X is $\sigma(X^{**}, X^*)$ -dense in $B_{X^{**}}$, so that $B_X = B_{X^{**}}$, whence X is reflexive. This yields (a).

Finally, assuming (a), we have that $B_{X^{**}}$ is $\sigma(X^{**}, X^*)$ -compact in $B_{X^{**}}$. Since $X \cong X^{**}$, it follows that B_X is $\sigma(X, X^*)$ -compact. $\square$

Corollary: If X is a reflexive Banach space, and $M \leq X$ is a closed subspace, then M is a reflexive Banach space.

Proof. Notice that the unit ball of M is the intersection of M with the unit ball of X , so the unit ball of M is $\sigma(X, X^*)$ -compact. Since $\sigma(X, X^*)|_M = \sigma(M, M^*)$, it follows that the unit ball of M is weakly compact with respect to $\sigma(M, M^*)$. $\square$

Corollary: If X is reflexive, then X is weakly sequentially complete.

Proof. Let $(x_n)_n$ be weakly Cauchy. Then, $(\langle x_n, \varphi \rangle)_n$ is Cauchy for each $\varphi \in X^*$. So, by the principle of uniform boundedness, $\|x_n\| \leq M$ for some constant M .

Since X is reflexive, $\{x \in X : \|x\| \leq M\}$ is weakly compact. So, there is some $x \in X$ such that $(x_{n_k})_{n_k} \rightarrow x$ weakly for some subsequence of $(x_n)_n$. Since $(\langle x_n, \varphi \rangle)_n$ converges for all $\varphi \in X^*$, it follows that $\langle x_n, \varphi \rangle \rightarrow \langle x, \varphi \rangle$ for all $\varphi \in X^*$, whence $(x_n)_n \rightarrow x$ weakly. $\square$

Corollary: If X is a reflexive Banach space, $M \leq X$ is a closed subspace, and $x_0 \in X \setminus M$, then there is some $y_0 \in M$ such that $\|x_0 - y_0\| = \text{dist}_M(x_0)$.

Proof. The map $x \mapsto \|x - x_0\|$ is weakly lower semicontinuous. If $d = \text{dist}_M(x_0)$, then the set $M \cap \{x : \|x - x_0\| \leq 2d\}$ is weakly compact. Since lower semicontinuous functions attain their minima on compact sets, the result follows. $\square$

Closed subspaces satisfying the property that for any $x_0 \in X \setminus M$, there exists $y_0 \in M$ such that $\|x_0 - y_0\| = \text{dist}_M(x_0)$ are known as proximal subspaces. Not every subspace of a normed vector space is proximal.

Proposition: Let X be a Banach space, $\varphi \in X^*$. Then, $\ker(\varphi)$ is proximal if and only if there is some $x \in X$ with $\|x\| = 1$ such that $\langle x, \varphi \rangle = \|\varphi\|$.

Proof. Suppose $M = \ker(\varphi)$ is proximal. If $f: X/M \rightarrow \mathbb{C}$ is given by $f(x + M) = \langle x, \varphi \rangle$, then f is a linear functional with $\|f\| = \|\varphi\|$. Since X/M is one-dimensional, there is $x \in X$ such that $\|x + M\| = 1$ and $f(x + M) = \|f\|$. Since M is proximal, there is $y \in M$ such that $1 = \|x + M\| = \|x + y\|$. Therefore,

$$\begin{aligned} \langle x + y, \varphi \rangle &= \langle x, \varphi \rangle \\ &= f(x + M) \\ &= \|f\| \\ &= \|\varphi\|. \end{aligned}$$

Now, suppose there is $x_0 \in X$ with $\|x_0\| = 1$ and $\langle x_0, \varphi \rangle = \|\varphi\|$. If $x \in X$ with $\|x + M\| = \alpha > 0$, then $\|\alpha^{-1}x + M\| = 1$; since $\|x_0 + M\| = 1$ and X/M is one-dimensional, it follows that there is $\beta \in \mathbb{C}$ with $|\beta| = 1$ such that $\alpha^{-1}x + M = \beta x_0 + M$, whence $\alpha^{-1}x - \beta x_0 \in M$. Equivalently, we have $x - \alpha\beta x_0 \in M$. Since

$$\begin{aligned} \|x - (x - \alpha\beta x_0)\| &= \|\alpha\beta x_0\| \\ &= \alpha \\ &= \text{dist}_M(x), \end{aligned}$$

it follows that the distance from x to M is attained at the element $x - \alpha\beta x_0 \in M$. $\square$

Separability and Metrizability

Theorem: If X is a Banach space, then B_{X^*} with the weak* topology is metrizable if and only if X is separable.

Proof. Let X be separable, $(x_n)_n$ a countable dense subset of B_X . Let $D = \prod_{n=1}^{\infty} B_{\mathbb{C}}(0, 1)$. Since D is a compact metric space, it follows that if B_{X^*} with the weak* topology is homeomorphic to a subset of D , then B_{X^*} is weak* metrizable.

Let $\tau: B_{X^*} \rightarrow D$ be defined by $\tau(\varphi) = (\langle x_n, \varphi \rangle)_n \in D$. If $(\varphi_i)_i$ is a net in B_{X^*} converging weak* to $\varphi \in X^*$, it follows that for each $n \geq 1$, we have that $\tau(\varphi_i) \rightarrow \tau(\varphi)$, so τ is weak* continuous. Furthermore, $\tau(\varphi) = 0$ if and only if $\varphi(x_n) = 0$ for all n ; since φ is then zero on a dense subset, it follows that φ is the zero functional, so τ is injective. Since B_{X^*} is compact, τ is a homeomorphism onto its image, so B_{X^*} is weak* metrizable.

Suppose B_{X^*} is weak* metrizable. Let $\{U_n : n \geq 1\}$ be a family of weak*-open neighborhoods of 0 whose intersection is 0. Then, for each n , there is some finite $F_n \subseteq X$ such that

$$\{\varphi \in B_{X^*} : |\langle x, \varphi \rangle| < 1 \text{ for all } x \in F_n\} \subseteq U_n.$$

Let $F = \bigcup_{n=1}^{\infty} F_n$. Then, F is countable. The pre-annihilator of $F^\perp$ is the closed linear span of F , which is separable. Yet, if $\varphi \in F^\perp$, then for each $n \geq 1$ and each $x \in F_n$, we have

$$\left| \left\langle x, \frac{\varphi}{\|\varphi\|} \right\rangle \right| = 0$$

$$< 1,$$

so $\frac{\varphi}{\|\varphi\|} \in U_n$ for all $n \geq 1$, whence $\varphi = 0$. Therefore, $F^\perp = \{0\}$, meaning its pre-annihilator is X . Thus, X is separable. $\square$

We may be curious about an analogous result for the weak topology. That is, if X^* is separable, is the weak topology on B_X metrizable? This follows immediately from the above theorem when we consider the embedding of X into X^{**} . However, this isn't very useful unless we're considering a separable reflexive space.

However, in general, if X is separable, it is not necessarily the case that B_X equipped with the weak topology is metrizable.

Theorem: If a sequence in ℓ^1 converges weakly, then the sequence converges in norm.

Proof. Since ℓ^1 is separable, we have that B_{ℓ^∞} is weak* metrizable. Since B_{ℓ^∞} is weak* compact (i.e., it is a compact metrizable space), we may use the Baire Category Theorem.

Let $(f_n)_n \rightarrow 0$ weakly in ℓ^1 , and let $\varepsilon > 0$. For each $m \geq 0$, let

$$F_m = \{\phi \in B_{\ell^\infty} : |\langle f_n, \phi \rangle| \leq \varepsilon/3 \text{ for all } n \geq m\}.$$

Then, F_m is weak*-closed in B_{ℓ^∞} . Since $(f_n)_n \rightarrow 0$ weakly, it follows that the union of all such F_m is equal to B_{ℓ^∞} .

By the Baire Category Theorem, there is some F_m with nonempty interior. A metric on B_{ℓ^∞} inducing the weak* topology is given by

$$d(\phi, \psi) = \sum_{j=1}^{\infty} 2^{-j} |\phi(j) - \psi(j)|.$$

Since F_m has nonempty weak* interior, there is $\phi \in F_m$ and $\delta > 0$ such that $U(\phi, \delta) \subseteq F_m$, where we recall that

$$U(\phi, \delta) = \{\psi \in B_{\ell^\infty} : d(\phi, \psi) < \delta\}.$$

Let J be such that $2^{-(J-1)} < \delta$. Let $n \geq m$, and let $\psi \in \ell^\infty$ be defined by $\psi(j) = \phi(j)$ for each $1 \leq j \leq J$ and $\psi(j) = \text{sgn}(f_n(j))$ for each $j > J$. Then, $\psi(j)f_n(j) = |f_n(j)|$ for each $j > J$. We have that $\psi \in B_{\ell^\infty}$.

Furthermore, we have

$$\begin{aligned} d(\phi, \psi) &= \sum_{j=J+1}^{\infty} 2^{-j} |\phi(j) - \psi(j)| \\ &\leq 2^{-(J-1)} \\ &< \delta. \end{aligned}$$

Thus, $\psi \in F_m$, so $|\langle f_n, \psi \rangle| \leq \varepsilon/3$ for all $n \geq m$. In particular, we have

$$\left| \sum_{j=1}^J \phi(j)f_n(j) + \sum_{j=J+1}^{\infty} |f_n(j)| \right| \leq \frac{\varepsilon}{3}$$

for all $n \geq m$.

Yet, since $(f_n)_n \rightarrow 0$ in ℓ^1 , we have some $m_1 \geq m$ such that for all $n \geq m_1$, we have

$$\sum_{j=1}^J |f_n(j)| < \varepsilon/3.$$

Therefore, we have

$$\begin{aligned}
 \|f_n\| &= \sum_{j=1}^{\infty} |f_n(j)| \\
 &< \frac{\varepsilon}{3} + \left| \sum_{j=j+1}^{\infty} |f_n(j)| + \sum_{j=1}^J \phi(j)f_n(j) \right| + \left| \sum_{j=1}^J \phi(j)f_n(j) \right| \\
 &< \frac{2\varepsilon}{3} + \sum_{j=1}^J |f_n(j)| \\
 &< \varepsilon
 \end{aligned}$$

whenever $n \geq m_1$. This gives norm convergence. $\square$

In particular, if it were the case that B_{ℓ^1} with the weak topology was metrizable, then we would have that the weak and norm topologies on ℓ^1 agreed. Yet, this is not the case.

Extremal Structure

Definition: Let X be a vector space, and let $K \subseteq X$ be a convex set. We call a subset $F \subseteq K$ a *face* if F is a convex subset satisfying the following criterion: if $0 < t < 1$, and $x, y \in K$ with $(1-t)x + ty \in F$ then $x, y \in F$.

A singleton set $\{z\} \in K$ that is a face is called an extreme point. We write $\text{ext}(K)$ for the set of extreme points of K .

In particular, $z \in K$ is an extreme point if and only if, for any $0 < t < 1$, if there are $x, y \in K$ with $(1-t)x + ty = z$, then $x = y = z$. In fact, we can take $t = 1/2$ — that is, we only need to concern ourselves with the midpoint of any such line segment. To see this, we assume without loss of generality that $t < 1/2$. Then, the vector $w = (1-2t)x + 2ty$ lies in C since C is convex, and satisfies $\frac{1}{2}(x+w) = z$. In particular, this means that $x = z = w$, so that $y = z$.

There are convex subsets not admitting extreme points.

Example: Let $X = c_0$. We will show that the unit ball B_{c_0} has no extreme points.

Let $z \in \text{ext}(B_{c_0})$. Notice that for any $0 < \|z\| < 1$, we have

$$z = (1 - \|z\|)0 + \|z\| \frac{z}{\|z\|},$$

meaning in particular that z cannot be an extreme point if $\|z\| < 1$. In particular, this means that $\|z\| = 1$.

Since $z \in c_0$, there is $n_0 \in \mathbb{N}$ with $|z(n_0)| < 1$. Write $z(n_0) = (1-t)\lambda + t\mu$ for some $0 < t < 1$ and $|\lambda| = |\mu| = 1$. Consider the sequences defined by

$$\begin{aligned}
 x(n) &= \begin{cases} z(n) & n \neq n_0 \\ \lambda & n = n_0 \end{cases} \\
 y(n) &= \begin{cases} z(n) & n \neq n_0 \\ \mu & n = n_0. \end{cases}
 \end{aligned}$$

Then, $\|x\|, \|y\| \leq 1$, but $(1-t)x + ty = z$ with $x \neq z$ and $y \neq z$, meaning that z cannot be extreme.

Proposition: Let Ω be a locally compact Hausdorff space.

- (i) If Ω is not compact, then $\text{ext}(B_{C_0(\Omega)}) = \emptyset$.

(ii) If Ω is compact, then

$$\text{ext}(B_{C(\Omega)}) = \{u \in C(\Omega) : |u(x)| = 1 \text{ for all } x \in \Omega\}.$$

Proof. Let $f \in \text{ext}(B_{C_0(\Omega)})$. Then, $\|f\| = 1$. Suppose there is some x_0 with $|f(x_0)| < 1$. Choose δ such that $|f(x_0)| < \delta < 1$. By continuity, there is some open neighborhood U of x_0 with $|f| < \delta$ on U . By Urysohn's Lemma, there is a continuous function $g: \Omega \rightarrow [0, 1]$ with $\text{supp}(g) \subseteq U$ and $g(x_0) = 1$.

Then, we may write

$$f = \frac{1}{2}((f + (1 - \delta)g) + (f - (1 - \delta)g)).$$

First, we observe that on U^c , we have that $|f \pm (1 - \delta)g| = |f| \leq 1$, while on U we have

$$|f \pm (1 - \delta)g| \leq |f| + (1 - \delta) \leq 1.$$

In particular, since $f \pm (1 - \delta)g \neq f$, this contradicts the fact that f is extreme, so in particular, we must have $|f(x)| = 1$ for all $x \in \Omega$, yielding that Ω is compact. This gives (i).

Having seen that (i) holds, we have that $\text{ext}(B_{C(\Omega)})$ is contained in the set of all $u \in C(\Omega)$ with $|u| = 1$ on Ω . Our next task is to show that any such u is extreme.

Write $u = \frac{1}{2}(f + g)$ with $f, g \in B_{C(\Omega)}$. Then, $u(x) = \frac{1}{2}(f(x) + g(x))$ with $|f(x)|, |g(x)| \leq 1$ and $|u(x)| = 1$. Since the extreme points of the closed unit ball of $\mathbb{C}$ is S^1 , it follows that $f(x) = g(x) = u(x)$ for all $x \in \Omega$. $\square$

C^* -Algebras

Spectral Theory

Spectral theory is primarily concerned with representations of abelian C^* -algebras. Consider a normal operator N on H . Then, $C^*(N)$ is an abelian C^* -algebra, and the continuous functional calculus $f \mapsto f(N)$ yields a $*$ -isomorphism of $C(\sigma(N))$ onto $C^*(N)$. In particular, $f \mapsto f(N)$ is a representation of the abelian C^* -algebra $C^*(N)$.

We will find a way to extend this further, as well as find some useful applications to the structure of abelian von Neumann algebras and understanding unitary invariants for normal operators.

Definition: Let X be a set, Σ a σ -algebra on X , and H a Hilbert space. A *spectral measure* for (X, Σ, H) is a map $E: \Sigma \rightarrow B(H)$ satisfying

- (a) for each $\Delta \in \Sigma$, $E(\Delta)$ is a projection;
- (b) $E(\emptyset) = 0$ and $E(X) = 1$;
- (c) $E(\Delta_1 \cap \Delta_2) = E(\Delta_1)E(\Delta_2)$ for any $\Delta_1, \Delta_2 \in \Sigma$;
- (d) given pairwise disjoint $\{\Delta_n\}_{n=1}^\infty$, we have

$$E\left(\bigcup_{n=1}^\infty \Delta_n\right) = \sum_{n=1}^\infty E(\Delta_n),$$

where the latter is an SOT-convergent sum of pairwise orthogonal projections converging to the projection onto the supremum $\bigwedge_{n=1}^\infty E(\Delta_n)H$.

Recall that the SOT and WOT are defined by the seminorms

- $p_h(A) = \|Ah\|$ for the SOT, and

- $p_{h,k}(H) = |\langle Ah, k \rangle|$ for the WOT.

Example: If (X, Ω, μ) is a σ -finite measure space, and $\phi \in L^\infty(X, \mu)$, then $M_\phi \in B(L^2(X, \mu))$ can be defined by $M_\phi(f) = \phi f$.

A net $(\phi_i)_{i \in I}$ in $L^\infty(X, \mu)$ is w^* -convergent to ϕ if and only if $M_{\phi_i} \rightarrow M_\phi$ in WOT. This follows from the fact that if $f, g \in L^2(X, \mu)$, then $f\bar{g} \in L^1(X, \mu)$, where we have $L^1(X, \mu)^* = L^\infty(X, \mu)$, so that if $(\phi_i)_i \rightarrow \phi$ weak*, then

$$\begin{aligned} \langle M_{\phi_i} f, g \rangle &= \int \phi_i(f\bar{g}) d\mu \\ &\rightarrow \int \phi f\bar{g} d\mu \\ &= \langle M_\phi f, g \rangle, \end{aligned}$$

while the converse emerges from the fact that any element $f \in L^1(X, \mu)$ can be written as $f = g_1 \bar{g}_2$.

Example: If E is a spectral measure for (X, Ω, H) , then $E^{(n)}(\Delta) = E(\Delta)^{(n)}$ (under the diagonal embedding) is a spectral measure for $(X, \Omega, H^{(n)})$.

We will now use spectral measures to define a representation of a C^* -algebra.

Lemma: Let E be a spectral measure on (X, Ω, H) , and let $g, h \in H$. Then,

$$E_{g,h}(\Delta) = \langle E(\Delta)g, h \rangle$$

defines a countably additive measure on Ω with total variation bounded above by $\|g\| \|h\|$.

Proof. For countable additivity, we note the classic result that a sequence of projections $(P_n)_n \rightarrow P$ converges in WOT if and only if it converges in SOT. In particular, this means that if $\{\Delta_n\}_{n \in \mathbb{N}}$ is a sequence of disjoint subsets of X , then we have a sequence of projections

$$\left(\sum_{k=0}^n E(\Delta_k) \right)_n \rightarrow \sum_{k=0}^{\infty} E(\Delta_k)$$

converging in SOT, hence converging in WOT. Yet, this means that

$$\begin{aligned} \left\langle \left(\sum_{k=0}^n E(\Delta_k) \right) g, h \right\rangle &= \sum_{k=0}^n \langle E(\Delta_k)g, h \rangle \\ &\rightarrow \left\langle \left(\sum_{k=0}^{\infty} E(\Delta_k) \right) g, h \right\rangle, \end{aligned}$$

so that

$$\sum_{k=0}^{\infty} \langle E(\Delta_k)g, h \rangle = \left\langle E \left(\bigcup_{k=0}^{\infty} \Delta_k \right) g, h \right\rangle,$$

yielding our desired result.

If we have a finite set $\Delta_1, \dots, \Delta_n$ of pairwise disjoint sets, we choose $|\alpha_j| = 1$ such that $|\langle E(\Delta_j)g, h \rangle| = \alpha_j \langle E(\Delta_j)g, h \rangle$, whence

$$\sum_{j=1}^n |\mu(\Delta_j)| = \left\langle \sum_{j=1}^n \alpha_j E(\Delta_j)g, h \right\rangle$$

$$\leq \left\| \sum_{j=1}^n E(\Delta_j) \alpha_j g \right\| \|h\|.$$

Notice that the family of $E(\Delta_j)$ are pairwise orthogonal projections, meaning that by Pythagoras's Theorem and the additivity of the spectral measure, we have

$$\begin{aligned} \left\| \sum_{j=1}^n E(\Delta_j) \alpha_j g \right\|^2 &= \sum_{j=1}^n \|E(\Delta_j) g\|^2 \\ &= \left\| E\left(\bigcup_{j=1}^n \Delta_j\right) g \right\|^2 \\ &\leq \|g\|^2, \end{aligned}$$

whence $\sum_{j=1}^n |\mu(\Delta_j)| \leq \|g\| \|h\|$. Since this holds for any finite collection of subsets, we have $\|\mu\|_{\text{TV}} \leq \|g\| \|h\|$. $\square$

We next concern ourselves with establishing “integration” with respect to the spectral measure.

Proposition: Let E be a spectral measure for (X, Ω, H) , and let $\phi: X \rightarrow \mathbb{C}$ be a bounded Ω -measurable function. Then, there is a unique operator $A \in B(H)$ such that for any $\varepsilon > 0$ and partition $\{\Delta_1, \dots, \Delta_n\}$ with $\sup_{x, x' \in \Delta_k} |\phi(x) - \phi(x')| < \varepsilon$ for each k , we have

$$\left\| A - \sum_{k=1}^n \phi(x_k) E(\Delta_k) \right\| < \varepsilon$$

for any $x_k \in \Delta_k$.

Proof. For each $g, h \in H$, define a (bounded) sesquilinear form

$$B(g, h) = \int \phi dE_{g, h},$$

with $|B(g, h)| \leq \|\phi\|_{L^\infty} \|g\| \|h\|$. Then, by the structure of bounded sesquilinear forms, there is a unique operator A such that $B(g, h) = \langle Ag, h \rangle$ for all $g, h \in H$.

Consider a partition $\{\Delta_1, \dots, \Delta_n\}$ satisfying the conditions in the proposition. For any arbitrary vectors $g, h \in H$ and $x_k \in \Delta_k$, we have

$$\begin{aligned} \left| \langle Ag, h \rangle - \sum_{k=1}^n \phi(x_k) \langle E(\Delta_k) g, h \rangle \right| &= \left| \sum_{k=1}^n \int_{\Delta_k} (\phi(x) - \phi(x_k)) dE_{g, h} \right| \\ &\leq \sum_{k=1}^n \int_{\Delta_k} |\phi(x) - \phi(x_k)| d|E_{g, h}| \\ &\leq \varepsilon \int d|E_{g, h}| \\ &\leq \varepsilon \|g\| \|h\|. \end{aligned}$$

This yields the desired result. $\square$

For the operator A in the above statement, we write

$$A = \int \phi dE.$$

That is, for any $g, h \in H$ and bounded Ω -measurable function ϕ on X , we have

$$\left\langle \left(\int \phi dE \right) g, h \right\rangle = \int \phi dE_{g,h}.$$

If $B_\infty(X, \Omega)$ denotes the set of bounded Ω -measurable functions $\phi: X \rightarrow \mathbb{C}$ with the norm $\|\phi\| = \sup_{x \in X} |\phi(x)|$, then $B_\infty(X, \Omega)$ is a unital Banach algebra — in fact, it is an abelian C^* -algebra.

Proposition: Suppose E is a spectral measure for (X, Ω, H) , and $\rho: B_\infty(X, \Omega) \rightarrow B(H)$ is defined by taking $\rho(\phi) = \int \phi dE$.

Then, ρ is a representation of $B_\infty(X, \Omega)$ and $\rho(\phi)$ is a normal operator for every $\phi \in B_\infty(X, \Omega)$.

Proof. We will only focus on showing that ρ is multiplicative. Suppose $\phi, \psi \in B_\infty(X, \Omega)$. Let $\varepsilon > 0$, and let $\{\Delta_1, \dots, \Delta_n\}$ be a Borel partition of X satisfying

$$\sup \{ |\omega(x) - \omega(x')| : x, x' \in \Delta_k, \omega = \phi, \psi, \phi\psi \} < \varepsilon.$$

In particular, we have that for any $x_k \in \Delta_k$ and $\omega = \phi, \psi, \phi\psi$, we have

$$\left\| \int \omega dE - \sum_{k=1}^n \omega(x_k) E(\Delta_k) \right\| < \varepsilon.$$

The triangle inequality then gives

$$\begin{aligned} \left\| \int \phi\psi dE - \left(\int \phi dE \right) \left(\int \psi dE \right) \right\| &\leq \varepsilon + \left\| \sum_{k=1}^n \phi(x_k) \psi(x_k) E(\Delta_k) - \left(\sum_{i=1}^n \phi(x_i) E(\Delta_i) \right) \left(\sum_{j=1}^n \psi(x_j) E(\Delta_j) \right) \right\| \\ &\quad + \left\| \left(\sum_{i=1}^n \phi(x_i) E(\Delta_i) \right) \left(\sum_{j=1}^n \psi(x_j) E(\Delta_j) \right) - \left(\int \phi dE \right) \left(\int \psi dE \right) \right\|. \end{aligned}$$

Since $E(\Delta_i)E(\Delta_j) = E(\Delta_i \cap \Delta_j)$, and $\{\Delta_1, \dots, \Delta_n\}$ is a partition, we have that the second term on the right-hand side is zero. Thus, we have

$$\begin{aligned} \left\| \int \phi\psi dE - \left(\int \phi dE \right) \left(\int \psi dE \right) \right\| &\leq \varepsilon + \left\| \left(\sum_{i=1}^n \phi(x_i) E(\Delta_i) \right) \left(\sum_{j=1}^n \psi(x_j) E(\Delta_j) - \int \psi dE \right) \right\| \\ &\quad + \left\| \left(\sum_{i=1}^n \phi(x_i) E(\Delta_i) - \int \phi dE \right) \left(\int \psi dE \right) \right\| \\ &\leq \varepsilon (1 + \|\phi\| + \|\psi\|). \end{aligned}$$

Thus, we get multiplicativity. $\square$

Corollary: If X is a compact Hausdorff space, and E is a spectral measure on H defined on the Borel sets of X , then $\rho: C(X) \rightarrow B(H)$ given by $\rho(u) = \int u dE$ is a representation of $C(X)$.

Thus far, all we have obtained is a variety of the properties of a spectral measure. However, we do not know if such a measure exists. The following is the proof that it does.

Theorem: Suppose $\rho: C(X) \rightarrow B(H)$ is a representation. Then, there is a unique spectral measure E defined on the Borel subsets of X such that for all $g, h \in H$, $E_{g,h}$ is a regular measure, and we have

$$\rho(u) = \int u dE$$

for all $u \in C(X)$.

Theorem: Let $g, h \in H$. The map $u \mapsto \langle \rho(u)g, h \rangle$ is a linear functional on $C(X)$ with norm at most $\|g\|\|h\|$. So, there is a unique measure $\mu_{g,h}$ in $M(X)$ such that

$$\langle \rho(u)g, h \rangle = \int u d\mu_{g,h}.$$

The pairing $(g, h) \mapsto \mu_{g,h}$ is sesquilinear, and for fixed $\phi \in B_\infty(X)$, the form

$$[g, h] = \int \phi d\mu_{g,h}$$

is a sesquilinear form with $\|[g, h]\| \leq \|\phi\|\|g\|\|h\|$. Therefore, there is a unique bounded operator A such that $[g, h] = \langle Ag, h \rangle$ with $\|A\| \leq \|\phi\|$. We will denote this operator A by $\tilde{\rho}(\phi)$. We obtain a well-defined map $\tilde{\rho}: B_\infty(X) \rightarrow B(H)$ satisfying $\|\tilde{\rho}(\phi)\| \leq \|\phi\|$ such that

$$\langle \tilde{\rho}(\phi)g, h \rangle = \int \phi d\mu_{g,h}.$$

We start by claiming that $\tilde{\rho}: B_\infty(X) \rightarrow B(H)$ is a representation with $\tilde{\rho}|_{C(X)} = \rho$. That $\tilde{\rho}$ restricted to $C(X)$ is equal to ρ follows directly from the definition of $\tilde{\rho}$.

Now, let $\phi \in B_\infty(X)$. Consider ϕ as an element of $M(X)^* = C(X)^{**}$ by viewing it as corresponding to the linear functional

$$\mu \mapsto \int \phi d\mu.$$

By the Banach–Alaoglu theorem, we have that $\{u \in C(X) : \|u\| \leq \|\phi\|\}$ is weak*-dense in $\{L \in M(X)^* : \|L\| \leq \|\phi\|\}$. Then, there is a net $(u_i)_i \subseteq C(X)$ with $\|u_i\| \leq \|\phi\|$ for all i and

$$\int \phi d\mu = \lim_i \int u_i d\mu$$

for all $\mu \in M(X)$. For any $\psi \in B_\infty(X)$, we have $\psi\mu \in M(X)$ for any $\mu \in M(X)$. In particular, we have

$$\int \phi\psi d\mu = \lim_i \int u_i\psi d\mu$$

for all $\psi \in B_\infty(X)$ and $\mu \in M(X)$, whence $\tilde{\rho}(u_i\psi) \rightarrow \tilde{\rho}(\phi\psi)$ in WOT. For any $\psi \in C(X)$, we have that

$$\begin{aligned} \tilde{\rho}(\phi\psi) &= \lim_i \tilde{\rho}(u_i\psi) \\ &= \lim_i \rho(u_i)\rho(\psi) \\ &= \tilde{\rho}(\phi)\rho(\psi). \end{aligned}$$

In particular, we have $\tilde{\rho}(u_i\psi) = \rho(u_i)\tilde{\rho}(\psi)$ for any $\psi \in B_\infty(X)$ for all $u_i \in C(X)$. Therefore, we have $\tilde{\rho}(u_i\psi) \rightarrow \tilde{\rho}(\phi\psi)$ in WOT, whence

$$\tilde{\rho}(\phi\psi) = \tilde{\rho}(\phi)\tilde{\rho}(\psi)$$

for any $\phi, \psi \in B_\infty(X)$.

The linearity of measures gives that $\tilde{\rho}$ itself is linear. A similar bootstrapping argument gives $\tilde{\rho}(\phi)^* = \tilde{\rho}(\overline{\phi})$.

Now, let $\Delta \subseteq X$ be Borel. Let $E(\Delta) = \tilde{\rho}(\chi_\Delta)$. We will show that E is a spectral measure.

First, since χ_Δ is a hermitian idempotent in $B_\infty(X)$, it follows that $E(\Delta)$ is a projection by the classification of projections on a Hilbert space. Furthermore, $\chi_\emptyset = 0$ and $\chi_X = 1$, so that $E(\emptyset) = 0$ and $E(X) = 1$. Since

$$E(\Delta_1 \cap \Delta_2) = \tilde{\rho}(\chi_{\Delta_1 \cap \Delta_2})$$

$$\begin{aligned} &= \widetilde{\rho}(\chi_{\Delta_1} \chi_{\Delta_2}) \\ &= E(\Delta_1)E(\Delta_2). \end{aligned}$$

Finally, let $\{\Delta_n : n \geq 0\}$ be a pairwise disjoint sequence of Borel sets. Then, for fixed $h \in H$ and $\Lambda_n = \bigcup_{k=n+1}^{\infty} \Delta_k$, we have

$$\begin{aligned} \left\| E\left(\bigcup_{k=1}^{\infty} \Delta_k\right) - \sum_{k=1}^n E(\Delta_k) \right\|^2 &= \langle E(\Lambda_n)h, E(\Lambda_n)h \rangle \\ &= \langle E(\Lambda_n)h, h \rangle \\ &= \int \chi_{\Lambda_n} d\mu_{h,h} \\ &= \sum_{k=n+1}^{\infty} \mu_{h,h}(\Delta_k) \\ &\xrightarrow{n \rightarrow \infty} 0. \end{aligned}$$

Thus, E is a spectral measure.

Finally, we must show that $\rho(u) = \int u dE$ for all $u \in C(X)$. In fact, we will show that $\widetilde{\rho}(\phi) = \int \phi dE$ for all $\phi \in B_{\infty}(X)$. For a fixed $\phi \in B_{\infty}(X)$, we choose a Borel partition $\{\Delta_1, \dots, \Delta_n\}$ such that $\sup\{|\phi(x) - \phi(x')| : x, x' \in \Delta_k\} < \varepsilon$ for each k . Then,

$$\left\| \phi - \sum_{k=1}^n \phi(x_k) \chi_{\Delta_k} \right\| < \varepsilon$$

for any choices of $x_k \in \Delta_k$. Since $\|\widetilde{\rho}\| = 1$, it follows that

$$\left\| \widetilde{\rho}(\phi) - \sum_{k=1}^n \phi(x_k) E(\Delta_k) \right\| < \varepsilon,$$

so that $\widetilde{\rho}(\phi) = \int \phi dE$ for any $\phi \in B_{\infty}(X)$.

The uniqueness of E follows from the fact that the measures $E_{g,h}$ are unique for each $g, h \in H$ as part of the Riesz Representation Theorem.

The primary use of the existence of spectral measures is to enable the Borel functional calculus for normal operators on Hilbert spaces.

Theorem: Let N be a normal operator on H . There is a unique spectral measure E on the Borel subsets of $\sigma(N)$ such that:

- (a) $N = \int z dE$;
- (b) if G is a nonempty open subset of $\sigma(N)$, then $E(G) \neq 0$;
- (c) if $T \in B(H)$, then $TN = NT$ and $TN^* = N^*T$ if and only if $TE(\Delta) = E(\Delta)T$ for all Borel $\Delta \subseteq \sigma(N)$.

Proof. Let $A = C^*(N)$. Then, A is the closure of the polynomials in N and N^* , so there is an isometric $*$ -isomorphism $\rho: C(\sigma(N)) \rightarrow A$ given by the functional calculus, $\rho(u) = u(N)$. There is a unique spectral measure E defined on the Borel subsets of $\sigma(N)$ such that $\rho(u) = \int u dE$ for all $u \in C(\sigma(N))$. This gives (a).

If G is a nonempty open subset of $\sigma(N)$, Urysohn's Lemma yields a continuous function u on $\sigma(N)$ such that $0 \leq u(z) \leq \chi_G(z)$ for all $z \in \sigma(N)$. Since $E(G) = \widetilde{\rho}(\chi_G) \geq \rho(u) \neq 0$, it follows that $E(G) \neq 0$.

Let $T \in B(H)$ be such that $TN = NT$ and $TN^* = N^*T$. Then, by the Stone–Weierstrass theorem, it follows that $Tu(N) = u(N)T$ for all $u \in C(\sigma(N))$.

Consider the set Ω defined as follows:

$$\Omega = \{\Delta \subseteq \sigma(N) : \Delta \text{ is a Borel subset and } AE(\Delta) = E(\Delta)A\}.$$

Notice that by the definition of the spectral measure, Ω is closed under finite intersections, countable unions (using SOT convergence), and contains $\sigma(N)$, so it is a σ -algebra by the Dynkin π - λ theorem. If $G \subseteq \sigma(N)$ is open, there is a sequence $(u_n)_n$ of positive continuous functions on $\sigma(N)$ such that $u_n(z) \nearrow \chi_G(z)$ for all $z \in \sigma(N)$. Dominated convergence then gives

$$\begin{aligned} \langle AE(G)g, h \rangle &= \langle E(G)g, A^*h \rangle \\ &= E_{g, A^*h}(G) \\ &= \int \chi_G(z) dE_{g, A^*h} \\ &= \int \lim_{n \rightarrow \infty} u_n(z) dE_{g, A^*h} \\ &= \lim_{n \rightarrow \infty} \int u_n(z) dE_{g, A^*h} \\ &= \lim_{n \rightarrow \infty} \langle Au_n(N)g, h \rangle \\ &= \lim_{n \rightarrow \infty} \langle u_n(N)Ag, h \rangle \\ &= \langle E(G)Ag, h \rangle. \end{aligned}$$

Thus, Ω contains all the open subsets of $\sigma(N)$, so it must be the collection of Borel sets. The converse follows from bootstrapping by approximating with simple functions. $\square$

The spectral measure will provide us a way to reframe normal operators as multiplication operators on $L^\infty(X, \mu)$ for some measure space (X, μ) .

Definition: We call $e_0 \in H$ *star-cyclic* for an operator A if the smallest reducing subspace for A that contains e_0 is H .

The operator A is *star-cyclic* if it contains a star-cyclic vector.

Similarly, $e_0 \in H$ is *cyclic* for A if H is the smallest invariant subspace for A that contains e_0 , and A is *cyclic* if it contains a cyclic vector.

Equivalently, e_0 is star-cyclic if and only if

$$H = \overline{\{Te_0 : T \in C^*(A)\}}$$

and e_0 is cyclic if and only if

$$H = \overline{\{p(A)e_0 : p \in \mathbb{C}[x]\}}.$$

Proposition: If A has either a cyclic or star-cyclic vector, then H is separable.

Proof. Notice that $C^*(A)$ and $\{p(A) : p \in \mathbb{C}[x]\}$ are separable subalgebras of $B(H)$. $\square$

If μ is a compactly supported measure on $K \subseteq \mathbb{C}$, and N_μ is the operator defined by $N_\mu \equiv M_z$, where $(M_z f)(z) = zf(z)$, then we observe that $C^*(N_\mu)$ is given by M_u for all $u \in C(K)$. Since continuous functions are dense in $L^2(K, \mu)$, it follows that 1 is a star-cyclic vector for N_μ .

Furthermore, the converse holds.

Theorem: A normal operator N is star-cyclic if and only if N is unitarily equivalent to N_μ for some compactly supported measure μ on $\mathbb{C}$. If e_0 is a star-cyclic vector for N , then μ can be chosen such that there is an isomorphism $V : H \rightarrow L^2(K, \mu)$ with $Ve_0 = 1$ and $VNV^{-1} = N_\mu$. Furthermore, V is unique.

Proof. Suppose $N \cong N_\mu$. Then, we have already seen that N is star-cyclic.

Suppose N has a star-cyclic vector e_0 . Let E be the spectral measure for N . Set $\mu(\Delta) = \langle E(\Delta)e_0, e_0 \rangle = \|E(\Delta)e_0\|^2$ for each Borel subset Δ of $\mathbb{C}$.

Let $K = \text{supp}(\mu)$. For any $\phi \in B_\infty(K)$, we have

$$\begin{aligned} \|\phi(N)e_0\|^2 &= \langle |\phi|^2(N)e_0, e_0 \rangle \\ &= \int |\phi(z)|^2 d\langle E(z)e_0, e_0 \rangle \\ &= \int |\phi|^2 d\mu. \end{aligned}$$

Viewing $B_\infty(K) \subseteq L^2(K, \mu)$, we have that $U\phi = \phi(N)e_0$ defines an isometry from $B_\infty(K)$ onto $\{\phi(N)e_0 : \phi \in B_\infty(K)\}$.

Since e_0 is star-cyclic, it follows that the range of U is dense in H , so U extends to an isomorphism $U : L^2(K, \mu) \rightarrow H$.

For any $\phi \in B(K)$, then we have

$$\begin{aligned} UN_\mu U^{-1}(\phi(N)e_0) &= UN_\mu(\phi) \\ &= U(z\phi) \\ &= N\phi(N)e_0, \end{aligned}$$

so that $UN_\mu U^{-1} = N$ on the dense subset $\{\phi(N)e_0 : \phi \in B_\infty(K)\}$ of $L^2(K, \mu)$. Therefore $UN_\mu U^{-1} = N$, and $V = U^{-1}$ is our desired selection. $\square$

We say two measures μ_1 and μ_2 are mutually absolutely continuous (or of the same measure class) if they have the same null sets. We will write $[\mu_1] = [\mu_2]$. If the respective Radon–Nikodym derivatives $\frac{d\mu_1}{d\mu_2}$ and $\frac{d\mu_2}{d\mu_1}$ are essentially bounded, then we say μ_1 and μ_2 are boundedly mutually absolutely continuous.

Theorem: We have that $N_{\mu_1} \cong N_{\mu_2}$ if and only if $[\mu_1] = [\mu_2]$.

Proof. Suppose $[\mu_1] = [\mu_2]$, and let $\phi = \frac{d\mu_1}{d\mu_2}$. If $g \in L^1(\mu_1)$, then $g\phi \in L^1(\mu_2)$, with

$$\int g\phi d\mu_2 = \int g d\mu_1.$$

For any $f \in L^2(\mu_1)$, we have $f\sqrt{\phi} \in L^2(\mu_2)$ with

$$\|f\sqrt{\phi}\|_{L^2(\mu_2)} = \|f\|_{L^2(\mu_1)}.$$

Define $U : L^2(\mu_1) \rightarrow L^2(\mu_2)$ by $Uf = f\sqrt{\phi}$. This is an isometry.

For any $g \in L^2(\mu_2)$, we have that $f = \phi^{-1/2}g \in L^2(\mu_1)$ with $Uf = g$, so U is surjective with $U^{-1}g = \phi^{-1/2}g$ for any $g \in L^2(\mu_2)$. For any $g \in L^2(\mu_2)$, $UN_{\mu_1}U^{-1}g = zg$, so $UN_{\mu_1}U^{-1} = N_{\mu_2}$.

Now, suppose $V : L^2(\mu_1) \rightarrow L^2(\mu_2)$ is an isomorphism satisfying $VN_{\mu_1}V^{-1} = N_{\mu_2}$. Let $\psi = V(1)$, with $\psi \in L^2(\mu_2)$. We will write $N_j = N_{\mu_j}$. Notice that for any polynomial $p \in \mathbb{C}[z, \bar{z}]$, we have $Vp(N_1, N_1^*)V^{-1} = p(N_2, N_2^*)$. Since $N_1 \cong N_2$, $\sigma(N_1) = \sigma(N_2)$, so μ_1 and μ_2 have the same support, which we will dub K .

Taking uniform limits of polynomials yields for all $u \in C(K)$, we have $Vu(N_1)V^{-1} = u(N_2)$, so we have $V(u) = u\psi$. Since V is an isometry, we have

$$\int |u|^2 d\mu_1 = \int |u|^2 |\psi|^2 d\mu_2$$

for every $u \in C(K)$. Therefore, we have

$$\int v d\mu_1 = \int v|\psi|^2 d\mu_2$$

for all $v \in C(K)$ with $v \geq 0$. By the Riesz representation theorem, we have that $\mu_1 = |\psi|^2 \mu_2$, so $\mu_1 \ll \mu_2$. Reversing the roles of N_1 and N_2 , and substituting V^{-1} for V where necessary, we obtain that $\mu_2 \ll \mu_1$, whence $[\mu_1] = [\mu_2]$. $\square$

Applications of the Spectral Theorem

Proposition: If N is a normal operator with spectral decomposition $N = \int z dE(z)$, we have that N is compact if and only if for every $\varepsilon > 0$, $E(\{z : |z| > \varepsilon\})$ is finite rank.

Proof. Let $\varepsilon > 0$, and let $\Delta_\varepsilon = \{z : |z| > \varepsilon\}$, with $E_\varepsilon = E(\Delta_\varepsilon)$. Then,

$$\begin{aligned} N - NE_\varepsilon &= \int z\chi_{\Delta_\varepsilon^c} dE(z) \\ &= \phi(N), \end{aligned}$$

where $\phi(z) = z\chi_{\Delta_\varepsilon^c}(z)$. In particular, $\|N - NE_\varepsilon\| \leq \varepsilon$. If E_ε has finite rank for every $\varepsilon > 0$, then NE_ε has finite rank, so $N \in \overline{F(H)}$.

If N is compact, then $\phi(z) = z^{-1}\chi_{\Delta_\varepsilon}(z)$ has $\phi(z) \in B(\mathbb{C})$. Since N is compact, we have that $N\phi(N)$ is compact, but then

$$\begin{aligned} N\phi(N) &= \int zz^{-1}\chi_{\Delta_\varepsilon}(z) dE(z) \\ &= E_\varepsilon, \end{aligned}$$

so E_ε is a compact projection, whence it is finite rank. $\square$

Theorem: If H is separable, and I is an ideal of $B(H)$ that contains a noncompact operator, then $I = B(H)$.

Proof. Let $A \in I \setminus K(H)$. Take the spectral decomposition

$$A^*A = \int t dE(t).$$

By the preceding proposition, there is some $\varepsilon > 0$ such that $P = E(\{t \geq 0 : \varepsilon < t < \infty\})$ has infinite rank. This yields that

$$\begin{aligned} P &= \left(\int t^{-1}\chi_{(\varepsilon, \infty)}(t) dE(t) \right) A^*A \\ &\in I. \end{aligned}$$

Since H is separable, $\text{hdim}(PH) = \aleph_0 = \text{hdim}(H)$. There is a surjective isometry $U : H \rightarrow PH$ satisfying $1 = U^*PU$. Since $P \in I$, $1 \in I$, so $I = B(H)$. $\square$

Corollary: The only nontrivial closed ideal of $B(H)$ is $K(H)$.

Proposition: If N is a normal operator on H , then there are reducing subspaces H_i such that $H = \bigoplus_{i \in I} H_i$ is a direct sum of star-cyclic subspaces.

Proof. Use Zorn's lemma to find a maximal set of vectors $\mathcal{E}$ such that, for any $e, f \in \mathcal{E}$ with $e \neq f$, we have that

$$\begin{aligned} H_e &= \overline{\{p(T, T^*)e : p \in \mathbb{C}[z, \bar{z}]\}} \\ H_f &= \overline{\{p(T, T^*)f : p \in \mathbb{C}[z, \bar{z}]\}} \end{aligned}$$

are orthogonal. Then, we have that $H = \bigoplus_{e \in \mathcal{E}} H_e$. $\square$

Corollary: Every normal operator is unitarily equivalent to the direct sum of star-cyclic normal operators.

Theorem: If N is a normal operator on H , then there is a measure space (X, Ω, μ) and a function $\phi \in L^\infty(X, \mu)$ such that $N \cong M_\phi$ on $L^2(X, \mu)$.

Proof. If $\{N_i\}_{i \in I}$ is a collection of star-cyclic normal operators such that $N \cong \bigoplus_{i \in I} N_i$, then $\sigma(N_i) \subseteq \sigma(N)$ for each N_i .

Let (X_i, Ω_i, μ_i) be measure spaces with μ_i a Borel measure, $X_i = \text{supp}(\mu_i)$, and $N_i \cong N_{\mu_i}$ acting on the measure space $L^2(X_i, \mu_i)$. Then, defining $X = \bigsqcup_{i \in I} X_i$, the corresponding disjoint union σ -algebra is defined to be the collection of all $\Delta \subseteq X$ such that $\Delta \cap X_i \in \Omega_i$ for all $i \in I$. The corresponding measure is then given by

$$\mu(\Delta) = \sum_{i \in I} \mu_i(\Delta \cap X_i),$$

and (X, Ω, μ) is then a measure space. For any $f \in L^2(X, \mu)$, we have that $f_i = f|_{X_i}$ is an element of $L^2(X_i, \mu_i)$, and the map $U: L^2(X, \mu) \rightarrow \bigoplus_{i \in I} L^2(X_i, \mu_i)$ given by taking $Uf = \bigoplus_{i \in I} f_i$ is an isomorphism.

Define $\phi: X \rightarrow \mathbb{C}$ by taking $\phi(z) = z$ if $z \in X_i$, where we observe that $X_i \subseteq \sigma(N)$ for each i . Therefore, ϕ is a bounded function. For any open subset of $\mathbb{C}$, we have that $\phi^{-1}(G) \cap X_i = G \cap X_i \in \Omega_i$, so ϕ is Ω -measurable, whence $\phi \in L^\infty(X, \mu)$.

It then follows that $UM_\phi U^{-1} = \bigoplus_{i \in I} N_{\mu_i} \cong N$. $\square$

Proposition: If A is a von Neumann subalgebra of $B(H)$, then A is the norm-closed linear span of its projections.

Proof. For any $a \in A$, we have that $a + a^*$ and $a - a^*$ are in A , so A is the linear span of $\text{Re}(A)$.

Let $a \in \text{Re}(A)$, and set the spectral decomposition

$$a = \int t dE(t).$$

Approximating by piecewise linear functions, we have for any $[a, b] \subseteq \mathbb{R}$, there is a sequence $(u_n)_n$ of continuous functions over $\mathbb{R}$ such that $0 \leq u_n \leq 1$, $u_n(t) = 1$ for any $a \leq t \leq b - \frac{1}{n}$ and $u_n(t) = 0$ for all $t \leq a - \frac{1}{n}$ and $t \geq b$. Therefore, $u_n(t) \rightarrow \chi_{[a, b]}(t)$ as $n \rightarrow \infty$.

For any $h \in H$, we have

$$\|(u_n(a) - E([a, b]))h\| = \int |u_n(t) - \chi_{[a, b]}(t)|^2 dE_{h, h}(t),$$

which goes to zero by dominated convergence. Therefore, $u_n(a) \rightarrow E([a, b])$ in SOT. Thus, all the spectral projections of the form $E([a, b])$ are in A .

For any open interval (α, β) containing $\sigma(A)$, we have for any $\varepsilon > 0$ a partition $\mathcal{P} = \{\alpha = t_0 < \dots < t_n = \beta\}$ such that

$$\left| t - \sum_{k=1}^n t_k \chi_{[t_{k-1}, t_k)} \right| < \varepsilon$$

for any $t \in \sigma(A)$. Therefore,

$$\left\| a - \sum_{k=1}^n E([t_{k-1}, t_k)) \right\| < \varepsilon,$$

meaning that $\text{Re}(A) \subseteq \overline{\text{span}}(P(A))$, yielding the desired result. $\square$

References

- [Ped89] Gert K. Pedersen. *Analysis Now*. Vol. 118. Graduate Texts in Mathematics. Springer-Verlag, New York, 1989, pp. xiv+277. ISBN: 0-387-96788-5. DOI: [10.1007/978-1-4612-1007-8](https://doi.org/10.1007/978-1-4612-1007-8). URL: <https://doi.org/10.1007/978-1-4612-1007-8>.
- [Con90] John B. Conway. *A Course in Functional Analysis*. Second. Vol. 96. Graduate Texts in Mathematics. Springer-Verlag, New York, 1990, pp. xvi+399. ISBN: 0-387-97245-5.