

frame=single, breaklines=true, columns=fullflexible

Matrix 4.8.11

ai25btech11015 – M Sai Rithik

Question

Find the vector equation of the plane determined by the points

$$A(3, -1, 2), \quad B(5, 2, 4), \quad C(-1, -1, 6),$$

and hence find the distance of this plane from the origin.

Solution

Step 1: Position vectors and edge vectors

$$\mathbf{A} = \begin{pmatrix} 3 \\ -1 \\ 2 \end{pmatrix}, \quad \mathbf{B} = \begin{pmatrix} 5 \\ 2 \\ 4 \end{pmatrix}, \quad \mathbf{C} = \begin{pmatrix} -1 \\ -1 \\ 6 \end{pmatrix}. \quad (1)$$

$$\mathbf{B} - \mathbf{A} = \begin{pmatrix} 5 \\ 2 \\ 4 \end{pmatrix} - \begin{pmatrix} 3 \\ -1 \\ 2 \end{pmatrix} = \begin{pmatrix} 2 \\ 3 \\ 2 \end{pmatrix}, \quad (2)$$

$$\mathbf{C} - \mathbf{A} = \begin{pmatrix} -1 \\ -1 \\ 6 \end{pmatrix} - \begin{pmatrix} 3 \\ -1 \\ 2 \end{pmatrix} = \begin{pmatrix} -4 \\ 0 \\ 4 \end{pmatrix}. \quad (3)$$

Step 2: Normal vector via cross product

Using the determinant/submatrix definition of the cross product,

$$(\mathbf{B} - \mathbf{A}) \times (\mathbf{C} - \mathbf{A}) = \begin{pmatrix} \begin{vmatrix} 3 \\ 2 \end{vmatrix} & \begin{vmatrix} 0 \\ 4 \end{vmatrix} \\ \begin{vmatrix} 2 \\ 2 \end{vmatrix} & \begin{vmatrix} -4 \\ 4 \end{vmatrix} \\ \begin{vmatrix} 2 \\ 3 \end{vmatrix} & \begin{vmatrix} -4 \\ 0 \end{vmatrix} \end{pmatrix}. \quad (4)$$

Now evaluate the 2×2 determinants:

$$(\mathbf{B} - \mathbf{A}) \times (\mathbf{C} - \mathbf{A}) = \begin{pmatrix} (3)(4) - (2)(0) \\ (2)(4) - (2)(-4) \\ (2)(0) - (3)(-4) \end{pmatrix}. \quad (5)$$

$$(\mathbf{B} - \mathbf{A}) \times (\mathbf{C} - \mathbf{A}) = \begin{pmatrix} 12 \\ 16 \\ 12 \end{pmatrix}. \quad (6)$$

We may simplify (factor out 4):

$$\mathbf{N}_0 = \begin{pmatrix} 3 \\ -4 \\ 3 \end{pmatrix} \times (-1) \quad \text{or} \quad \mathbf{N} = \begin{pmatrix} 3 \\ -4 \\ 3 \end{pmatrix}, \quad (7)$$

(we choose $\mathbf{N} = (3, -4, 3)^\top$, sign is arbitrary: any nonzero scalar multiple is a valid normal).

Step 3: Finding plane Equation

Compute the RHS constant d for the plane $\mathbf{N}^\top x = d$ using point A :

$$d = \mathbf{N}^\top \mathbf{A} = \begin{pmatrix} 3 & -4 & 3 \end{pmatrix} \begin{pmatrix} 3 \\ -1 \\ 2 \end{pmatrix} = 3 \cdot 3 + (-4) \cdot (-1) + 3 \cdot 2. \quad (8)$$

$$d = 9 + 4 + 6 = 19. \quad (9)$$

Thus the plane equation in standard form is

$$\boxed{\mathbf{N}^\top x = 19 \quad \text{with} \quad \mathbf{N} = \begin{pmatrix} 3 \\ -4 \\ 3 \end{pmatrix}}. \quad (10)$$

Scale the normal so that the right-hand side becomes 1. Define

$$\tilde{\mathbf{N}} = \frac{\mathbf{N}}{19} = \begin{pmatrix} \frac{3}{19} \\ -\frac{4}{19} \\ \frac{3}{19} \end{pmatrix}. \quad (11)$$

Then the plane can be written as

$$\boxed{\tilde{\mathbf{N}}^\top x = 1}, \quad (12)$$

since $\tilde{\mathbf{N}}^\top x = (\mathbf{N}^\top x)/19$ and $\mathbf{N}^\top \mathbf{A} = 19$.

Step 5: Distance of the plane from the origin

For a plane written as $\tilde{\mathbf{N}}^\top x = 1$, the perpendicular distance D from the origin to the plane equals

$$D = \frac{|1|}{\|\tilde{\mathbf{N}}\|}. \quad (13)$$

Compute $\|\mathbf{N}\|$ first:

$$\|\mathbf{N}\| = \sqrt{3^2 + (-4)^2 + 3^2} = \sqrt{9 + 16 + 9} = \sqrt{34}. \quad (14)$$

Since $\tilde{\mathbf{N}} = \mathbf{N}/19$, we have $\|\tilde{\mathbf{N}}\| = \|\mathbf{N}\|/19$. Hence

$$D = \frac{1}{\|\tilde{\mathbf{N}}\|} = \frac{19}{\|\mathbf{N}\|}. \quad (15)$$

$$D = \frac{19}{\sqrt{34}} = \frac{19\sqrt{34}}{34} \approx 3.260. \quad (16)$$

Final Answer

$$\tilde{\mathbf{N}}^\top x = 1, \quad \tilde{\mathbf{N}} = \begin{pmatrix} \frac{3}{19} \\ \frac{4}{19} \\ -\frac{3}{19} \end{pmatrix}, \quad (17)$$

$$\text{Distance from origin to plane} = \frac{19}{\sqrt{34}} \approx 3.260. \quad (18)$$

Plane through A, B, C and its distance from origin




Figure 1: