

## 4.8.30

AI25BTECH11034 - Sujal Chauhan

October 2, 2025

# Question

Find the equation of a line passing through the point  $(2,3,2)$  and parallel to the line  $\mathbf{r} = (-2\hat{i} + 3\hat{j}) + \lambda(2\hat{i} - 3\hat{j} + 6\hat{k})$ . Also, find the distance between these two lines.

# Theory:

Consider two parallel lines in 3D:

$$\mathbf{r}_1 = \mathbf{a}_1 + \lambda \mathbf{b}, \quad \lambda \in \mathbb{R}, \quad (1)$$

$$\mathbf{r}_2 = \mathbf{a}_2 + \mu \mathbf{b}, \quad \mu \in \mathbb{R}, \quad (2)$$

where  $\mathbf{a}_1, \mathbf{a}_2$  are points on the respective lines and  $\mathbf{b}$  is the common direction vector.

Shortest distance between them can be given by :

$$d^2 = (\mathbf{a}_2 - \mathbf{a}_1)^\top (\mathbf{a}_2 - \mathbf{a}_1) - \left( \frac{(\mathbf{a}_2 - \mathbf{a}_1)^\top \mathbf{b}}{\|\mathbf{b}\|} \right)^2 \quad (3)$$

Now we can calculate d.

## Solution:

The direction vector of the given parallel lines is

$$\mathbf{b} = \begin{pmatrix} 2 \\ -3 \\ 6 \end{pmatrix}. \quad (4)$$

The first line is given by

$$\mathbf{r}_1 = \begin{pmatrix} 2 \\ 3 \\ 2 \end{pmatrix} + \mu \begin{pmatrix} 2 \\ -3 \\ 6 \end{pmatrix}, \quad \mu \in \mathbb{R}. \quad (5)$$

The second line is

$$\mathbf{r}_2 = \begin{pmatrix} -2 \\ 3 \\ 0 \end{pmatrix} + \lambda \begin{pmatrix} 2 \\ -3 \\ 6 \end{pmatrix}, \quad \lambda \in \mathbb{R}. \quad (6)$$

Now, we compute the difference between the position vectors  $\mathbf{a}_1$  and  $\mathbf{a}_2$ :

$$\mathbf{a}_2 - \mathbf{a}_1 = \begin{pmatrix} -2 \\ 3 \\ 0 \end{pmatrix} - \begin{pmatrix} 2 \\ 3 \\ 2 \end{pmatrix} = \begin{pmatrix} -4 \\ 0 \\ -2 \end{pmatrix}. \quad (7)$$

Next, we calculate the required components for the distance formula. The squared magnitude of  $(\mathbf{a}_2 - \mathbf{a}_1)$  is:

$$(\mathbf{a}_2 - \mathbf{a}_1)^\top (\mathbf{a}_2 - \mathbf{a}_1) = \|\mathbf{a}_2 - \mathbf{a}_1\|^2 \quad (8)$$

$$= (-4)^2 + (0)^2 + (-2)^2 \quad (9)$$

$$= 16 + 0 + 4 = 20. \quad (10)$$

$$(\mathbf{a}_2 - \mathbf{a}_1)^\top \mathbf{b} = \begin{pmatrix} -4 & 0 & -2 \end{pmatrix} \begin{pmatrix} 2 \\ -3 \\ 6 \end{pmatrix} \quad (11)$$

$$= (-4)(2) + (0)(-3) + (-2)(6) \quad (12)$$

$$= -8 + 0 - 12 = -20. \quad (13)$$

The magnitude of the direction vector  $\mathbf{b}$  is:

$$\|\mathbf{b}\| = \sqrt{2^2 + (-3)^2 + 6^2} \quad (14)$$

$$= \sqrt{4 + 9 + 36} = \sqrt{49} = 7. \quad (15)$$

Substituting these values into the formula for the squared distance:

$$d^2 = (\mathbf{a}_2 - \mathbf{a}_1)^\top (\mathbf{a}_2 - \mathbf{a}_1) - \left( \frac{(\mathbf{a}_2 - \mathbf{a}_1)^\top \mathbf{b}}{\|\mathbf{b}\|} \right)^2 \quad (16)$$

$$= 20 - \left( \frac{-20}{7} \right)^2 \quad (17)$$

$$= 20 - \frac{400}{49} \quad (18)$$

$$= \frac{20 \times 49 - 400}{49} \quad (19)$$

$$= \frac{980 - 400}{49} \quad (20)$$

$$= \frac{580}{49}. \quad (21)$$

Finally, the shortest distance  $d$  is the square root of this value:

$$d = \sqrt{\frac{580}{49}} = \frac{\sqrt{580}}{7} = \frac{\sqrt{4 \times 145}}{7} = \frac{2\sqrt{145}}{7}. \quad (22)$$

Thus, the distance between the two parallel lines is

$$d = \frac{2\sqrt{145}}{7}.$$



# Graph

