

Bacterial Purine Base Oxidation to Urate: A Convergent, Modular XdhAB-Centered System

A commissioned review-style synthesis for a molecular biology audience

1. Executive Summary

Bacterial oxidation of purine bases to urate is best understood not as a single linear pathway but as a **compact, reusable metabolic module** organized around one catalytic core: a **molybdo-flavo-iron-sulfur enzyme of the xanthine oxidase (XO) family**. The defining chemistry of the system is oxidative hydroxylation at a molybdenum center, in which a metal-bound, water-derived hydroxide attacks a purine carbon with concomitant hydride transfer to a Mo=S group. This same chemistry is used twice in the fully realized pathway — first to convert **hypoxanthine** → **xanthine**, and then to perform the obligatory terminal step **xanthine** → **urate**.

The system's most instructive architectural feature is **substrate-gated convergence**. Xanthine, the committed intermediate, can be supplied by two mechanistically independent branches: (i) **NAD⁺-dependent oxidation of hypoxanthine** by the XO-family enzyme itself, and (ii) **zinc-dependent deamination of guanine** by an amidohydrolase-superfamily guanine deaminase that is evolutionarily and mechanistically unrelated to the molybdenum enzyme. Both branches deliver xanthine to the same terminal oxidation. In the **two-subunit "XdhAB" realization** — best characterized in the phototroph *Rhodobacter capsulatus* — a single $\alpha_2\beta_2$ heterotetramer carries out *both* NAD⁺-dependent oxidations (hypoxanthine→xanthine and xanthine→urate), so one enzyme spans the hypoxanthine-supply and terminal steps, whereas the guanine branch is enzymatically distinct but convergent.

Three cross-cutting conclusions define the boundaries and reliability of this model. First, the catalytic **core chemistry and cofactor set are ancient and conserved**, but the **packaging is variable**: the same three cofactor modules (Mo-molybdopterin, FAD, two [2Fe-2S] clusters, plus NAD⁺ binding) appear as a single eukaryotic polypeptide, as the two-subunit bacterial XdhAB,

and as three-subunit heterotrimers such as *Escherichia coli* PaoABC. Second, each architectural variant requires a **dedicated molybdenum-cofactor (Moco) insertion chaperone** — XdhC for XdhAB, PaoD for PaoABC — that is not itself part of the mature catalytic enzyme. Third, the "one enzyme does both oxidations" statement is **realization-specific, not universal**: at least one lineage (*Klebsiella pneumoniae*) possesses a **molybdenum-cofactor-independent route** for hypoxanthine utilization, and the downstream conversion of urate to allantoin belongs to a **separate regulon** that should be treated as outside this system's boundary.

2. Definition and Biological Boundaries

2.1 What is included

The biological system reviewed here comprises the **enzymatic oxidation of purine bases to urate in bacteria**, bounded by two entry substrates and one exit product:

- **Entry point A** — **guanine**, converted to xanthine by **guanine deaminase** (a Zn-dependent amidohydrolase-superfamily enzyme).
- **Entry point B** — **hypoxanthine**, converted to xanthine by **XdhAB hypoxanthine dehydrogenase activity** (NAD⁺-dependent).
- **Committed intermediate** — **xanthine**, on which both branches converge.
- **Terminal step** — **xanthine → urate**, catalyzed by **XdhAB xanthine dehydrogenase activity** (NAD⁺-dependent), the obligatory reaction shared by all routes.

The system is therefore a two-input, one-output module: two substrate-gated supply branches feeding a single obligatory terminal oxidation.

2.2 What should be treated separately

Several neighboring processes are frequently conflated with this module but lie outside its boundaries:

- **Downstream urate degradation (urate → allantoin → glyoxylate/carbamoyl phosphate)**. In bacteria such as *E. coli* this is governed by a distinct regulatory system (the ALLR/ALLS/RutR network) that switches purine-derived nitrogen between assimilation and energy production ([PMID: 18957590](#)). Urate production is the *endpoint* of the system reviewed here; its consumption is a separate regulon.

- **Moco-insertion chaperones (XdhC, PaoD).** These are essential for producing active enzyme but are **not catalytic subunits** and do not participate in the reaction chemistry (see §4.2).
- **Other molybdenum-enzyme families.** The XO family is one of three structurally defined mononuclear Mo-enzyme families (xanthine oxidase, sulfite oxidase, DMSO reductase) (PMID: 19452052). Nitrate reductases and formate dehydrogenases (bis-MGD, DMSO reductase family) share Mo chemistry but are mechanistically distinct and unrelated to purine oxidation (PMID: 15311335).
- **Mammalian/eukaryotic purine oxidation.** Eukaryotic XDH performs the same net chemistry but as a single-chain enzyme, and eukaryotic guanine deaminase is a different protein family (see §2.3, §5). Data should not be transferred uncritically between kingdoms.


2.3 Competing definitions

The chief definitional tension is whether "the enzyme" is **one protein or two (or three)**. In eukaryotes, the Mo-molybdopterin, FAD, and [2Fe-2S] modules reside in a single polypeptide; in bacteria they are split. Notably, the bacterial *R. capsulatus* enzyme is **more similar in sequence to eukaryotic XDH** than to other prokaryotic molybdenum enzymes such as the *Desulfovibrio gigas* aldehyde oxidoreductase, despite the eukaryotic enzyme being single-chain (PMID: 9515710). Thus "XDH" as a functional label spans several distinct quaternary realizations, and the "two-subunit XdhAB" is one point on a continuum of architectures.

3. Mechanistic Overview

3.1 The sequence of events

The best-supported model is a two-input converging pathway:



- **Obligatory step:** xanthine → urate. Every route to urate passes through this terminal oxidation.
- **Conditional/substrate-gated steps:** the two supply branches. Which branch operates depends on which purine base is available; the branches are **non-exclusive** (both can run in the same cell when both substrates are present) and are induced under nitrogen-source control.
- **Accessory (not part of catalysis):** Moco-insertion chaperones (XdhC/PaoD), required for biogenesis of active enzyme but not for the reaction cycle.

3.2 The conserved catalytic chemistry

Both oxidative hydroxylations use the canonical XO-family mechanism, established structurally on the *D. gigas* aldehyde oxidoreductase (MOP) prototype ([PMID: 8799115](#)). At the Mo(VI) center bearing oxo and sulfido ligands:

1. A **molybdenum-bound water/hydroxide** is activated (proton transfer to an active-site glutamate — Glu-869 in the MOP numbering).
2. The hydroxide performs **nucleophilic attack on the substrate carbon** (the C2/C8 carbonyl carbon of the purine).
3. This is **concerted with hydride transfer to the Mo=S sulfido group**, reducing Mo(VI)→Mo(IV).
4. **Water is regenerated** from an internal chain of catalytically relevant waters, resetting the active site.

Crucially, the oxygen atom incorporated into the product comes from **water**, not molecular O₂ — a hallmark distinguishing this chemistry from oxygenases. Electrons abstracted at Mo are then routed intramolecularly through the **[2Fe-2S] clusters to FAD**, where in the dehydrogenase form they reduce NAD⁺. This electron-relay wiring is why the enzyme is a "molybdo-flavo-iron-sulfur" complex: each cofactor module is a station on an electron conduit.

3.3 Why the same enzyme can do both oxidations

Hypoxanthine→xanthine and xanthine→urate are both **oxidative hydroxylations of a purine carbon** — the same reaction type applied to successive positions. Because the XO-family active site catalyzes purine-carbon hydroxylation generically, a single XdhAB complex is chemically competent for both. In the two-subunit realization, this is exactly what happens: the hypoxanthine-supply branch and the terminal step are catalyzed by **one $\alpha_2\beta_2$ enzyme**, while the guanine branch (a hydrolytic deamination, not a hydroxylation) necessarily requires a **different enzyme** (PMID: 9515710; PMID: 11101664).

4. Major Molecular Players and Active Assemblies

4.1 The two-subunit XdhAB complex (Finding F001)

In *R. capsulatus*, xanthine dehydrogenase is encoded by adjacent **xdhA** and **xdhB** genes and assembles as an **$\alpha_2\beta_2$ heterotetramer**. The cofactor division is explicit:

Subunit	Cofactors carried	Functional role
XdhA	two [2Fe-2S] clusters + FAD	electron relay and NAD ⁺ reduction
XdhB	molybdopterin (Moco) center	substrate hydroxylation

The deduced XdhA sequence contains binding sites for two [2Fe-2S] clusters and FAD, whereas XdhB contains the molybdopterin cofactor; in eukaryotic XDH these three modules reside in a single chain (PMID: 9515710). The two-subunit composition of the active enzyme is confirmed independently (PMID: 10217763). Expression is induced when **hypoxanthine or xanthine** is supplied as sole nitrogen source, tying the same *xdhA* locus to both the supply and terminal reactions (PMID: 9515710).

4.2 XdhC — a maturation chaperone, not a subunit (Finding F002)

xdhC is cotranscribed with *xdhAB* and is required for XDH activity, yet **XdhC is not part of the active enzyme**. XDH purified from an *xdhC* mutant contains FAD and iron–sulfur clusters but **lacks the molybdopterin cofactor**, showing that XdhC functions specifically as a **Moco/MPT insertase and chaperone**. The paper states plainly that "XDHC is not a subunit of active XDH, which forms an $\alpha_2\beta_2$ heterotetramer in *R. capsulatus*," and that "in the absence of

XDHC, no molybdopterin cofactor MPT is present in the XDHAB tetramer" (PMID: 10217763). This distinction — biogenesis factor versus catalytic subunit — is a common source of confusion in describing the system.

4.3 Guanine deaminase — an independent, Zn-dependent branch (Finding F003)

The guanine branch is catalyzed by **guanine deaminase (GDEase)**, in *Bacillus subtilis* encoded by *yknA/gde*. Mutants cannot use guanine as a nitrogen source and lack detectable GDEase activity; the enzyme is induced by purines under **GlnR/TnrA nitrogen control**. Critically, "the first step is the deamination of guanine to xanthine catalysed by guanine deaminase (GDEase)," and "the GDEase amino acid sequence shows no homology with the mammalian enzyme" (PMID: 11101664). Structurally, bacterial guanine deaminases belong to the **amidohydrolase superfamily** and use an active-site **zinc** center, with substrate-selectivity determinants distinguishing them within the superfamily; guanine deaminase "is a metabolic enzyme, found in all forms of life, which catalyzes the conversion of guanine to xanthine" (PMID: 31283204). This branch is thus **mechanistically orthogonal** to the molybdenum chemistry: hydrolytic deamination at Zn, not redox hydroxylation at Mo.

4.4 The conserved catalytic mechanism and a Moco-independent exception (Finding F004)

The XO-family mechanism (§3.2) is shared by the terminal oxidation and the hypoxanthine oxidation. The MOP structural work describes the reaction proceeding "by transfer of the molybdenum-bound water molecule as OH⁻ after proton transfer to Glu-869 to the carbonyl carbon of the substrate in concert with hydride transfer to the sulfido group" (PMID: 8799115). However, the assumption that hypoxanthine oxidation *always* requires Moco is refuted in at least one organism: in *Klebsiella pneumoniae*, molybdenum-cofactor (chlorate-resistant) mutants and the wild-type parent "grew equally well with hypoxanthine as the sole nitrogen source, suggesting that *K. pneumoniae* has a molybdenum cofactor-independent pathway for hypoxanthine utilization" (PMID: 1400180). This is a genuine boundary/controversy point: the "one XdhAB does both oxidations" model is realization-specific.

4.5 Architectural variation across the XO family (Finding F005)

The same cofactor set is packaged in **one, two, or three chains**:

Architecture	Example	Cofactors	Dedicated chaperone
Single-chain	eukaryotic XDH	Mo-MPT + FAD + 2×[2Fe-2S] in one polypeptide	(eukaryotic system)
Two-subunit ($\alpha_2\beta_2$)	<i>R. capsulatus</i> XdhAB	XdhA: FAD + 2×[2Fe-2S]; XdhB: Mo-MPT	XdhC
Three-subunit ($\alpha\beta\gamma$)	<i>E. coli</i> PaoABC	2×[2Fe-2S], FAD, MCD (molybdopterin-cytosine-dinucleotide), + extra [4Fe-4S]	PaoD

PaoABC is "the only heterotrimer of the XO family so far structurally characterized," is the first *E. coli* protein shown to contain an MCD cofactor, and unexpectedly harbors an extra [4Fe-4S] cluster; it does **not** dimerize through its Mo-domain (PMID: 27622978; PMID: 24492481). Each architecture uses its own maturation chaperone: "PaoD is the chaperone of the periplasmic aldehyde oxidoreductase PaoABC ... its presence is crucial for obtaining mature enzyme" (PMID: 24498065), exactly paralleling XdhC's role for XdhAB.

4.6 The modular, convergent architecture (Finding F006)

Synthesizing the above: guanine (via Zn-dependent deaminase) and hypoxanthine (via NAD⁺-dependent XdhAB hydroxylation) both generate xanthine, which is oxidized to urate by the same XO-family Mo mechanism. In the two-subunit realization, one XdhAB complex carries both NAD⁺-dependent oxidations, so the hypoxanthine-supply and terminal steps share an enzyme, while the guanine branch is enzymatically independent but convergent. The terminal xanthine→urate step is **obligatory**; the two supply branches are **conditional and non-exclusive** (PMID: 11101664; PMID: 9515710; PMID: 8799115).

5. Evolutionary and Cell-Biological Variation

5.1 Conservation of the core, variation in packaging

The **catalytic core is ancient and conserved**: the XO-family Mo-hydroxylation chemistry, the pyranopterin (molybdopterin) cofactor, the [2Fe-2S]→FAD electron relay, and the water-derived oxygen atom are shared from sulfate-reducing bacteria (*Desulfovibrio*) through phototrophs (*Rhodobacter*) to eukaryotes. Molybdenum/tungsten pyranopterin enzymes are "ubiquitous in

Nature" and catalyze two-electron redox reactions central to C, N, and S metabolism (PMID: 19452052). The **variable, later-elaborated features** are quaternary architecture (1/2/3 subunits), the specific pyranopterin dinucleotide form (bare MPT vs. MCD vs. MGD), auxiliary Fe-S content (the extra [4Fe-4S] in PaoABC), and lineage-specific chaperone identity.

5.2 Best representatives of the ancestral role

For understanding the ancestral catalytic role of the expanded XO family, the **best-characterized structural prototypes** are the *Desulfovibrio* aldehyde oxidoreductases (MOP from *D. gigas*, MOD from *D. desulfuricans*), refined to near-atomic resolution and defining the family's active-site geometry, water chain, and Mo coordination (PMID: 11713686; PMID: 27520791; PMID: 10704312). These enzymes anchor the mechanistic model even though their physiological substrates are aldehydes rather than purines — the point is that the **chemistry and scaffold** are ancestral, and substrate specialization (purine hydroxylation) is a lineage-specific tuning of a conserved active site.

The breadth of the family within a single organism is illustrated by *Starkeya novella*, whose molybdoproteome encodes **18 gene loci** across the XO, sulfite oxidase, and DMSO reductase families, ~70% of which have no characterized close relatives (PMID: 23310928). This underscores that "xanthine dehydrogenase" is one specialized outgrowth of a large, diversified enzyme family, and that functional assignment from sequence alone is hazardous.

5.3 Physiological and regulatory context

Across bacteria, the purine-oxidation module is typically deployed under **nitrogen limitation**, allowing purines to serve as N (and sometimes C) sources. Induction is governed by nitrogen-control regulators — GlnR/TnrA in *B. subtilis* for the guanine branch (PMID: 11101664) — and substrate availability gates which branch operates. The downstream fate of urate (to allantoin and beyond) is controlled independently, e.g., by AllR/AllS/RutR in *E. coli*, which partitions purine nitrogen between assimilation (anaerobic) and energy production (aerobic) (PMID: 18957590). Compartmentalization can also vary: PaoABC is **periplasmic**, whereas classic XdhAB is cytoplasmic — a cell-biological difference relevant to substrate access and electron-acceptor choice.

5.4 Plants, eukaryotes, and cross-kingdom caveats

XDH is found in eukaryotes, bacteria, and archaea, catalyzing xanthine/hypoxanthine → uric acid with roles extending into nitrogen metabolism, ROS metabolism, hormone metabolism, and stress responses (PMID: 39467736). The conserved net chemistry across kingdoms is real,

but the **quaternary structure and the identity of the guanine deaminase differ** (bacterial GDEase has no homology to the mammalian enzyme; [PMID: 11101664](#)). Overgeneralizing from mammalian urate physiology (much of the surveyed literature concerns hyperuricemia and gut-microbial urate degradation, e.g., [PMID: 42417957](#), [PMID: 41703989](#)) to bacterial pathway architecture is not warranted.

6. Constraints, Dependencies, and Failure Modes

Ordering constraints (obligatory sequence). - Guanine must be **deaminated to xanthine before** it can be oxidized to urate — the guanine branch converges on, and cannot bypass, the terminal step. - Hypoxanthine must be **oxidized to xanthine before** proceeding to urate — the hypoxanthine branch runs *through* xanthine. - The terminal **xanthine → urate** step is obligatory for all routes.

Cofactor and biogenesis dependencies (failure modes). - Active XdhAB requires **Moco insertion by XdhC**; without XdhC, the tetramer assembles with FAD and Fe-S but is Moco-deficient and inactive ([PMID: 10217763](#)). Loss of the chaperone is a specific, cofactor-level failure mode distinct from loss of the catalytic subunits. - Analogously, PaoABC requires **PaoD** for maturation ([PMID: 24498065](#)). - The catalytic cycle depends on the intact **electron relay** ($\text{Mo} \rightarrow [2\text{Fe-2S}] \rightarrow \text{FAD} \rightarrow \text{NAD}^+$); disruption of any station uncouples substrate oxidation from cofactor re-oxidation.

Substrate/mechanism exclusivity. - The **guanine branch is hydrolytic (Zn)**; the **hypoxanthine and terminal branches are redox (Mo)**. These are mechanistically mutually exclusive chemistries carried by different proteins — a guanine deaminase cannot perform hydroxylation and vice versa. - Compartment specificity (periplasmic PaoABC vs. cytoplasmic XdhAB) constrains which substrates and electron acceptors are physiologically accessible.

Evidence that rules out otherwise-plausible paths. - The *Klebsiella* Moco-mutant experiment rules out the assumption that hypoxanthine utilization is universally Moco-dependent: a **Moco-independent route** exists in that lineage ([PMID: 1400180](#)). This forbids treating "XdhAB does both oxidations" as a universal law. - The *xdhC* purification data rule out XdhC being a catalytic subunit ([PMID: 10217763](#)).


7. Controversies and Open Questions

1. **Is hypoxanthine oxidation always Moco-dependent?** No, not universally. The *K. pneumoniae* result implies an alternative (possibly non-XO-family) hypoxanthine-utilization route ([PMID: 1400180](#)). The molecular identity of this Moco-independent activity remains unresolved and is arguably the single most important open mechanistic question for the hypoxanthine branch.
 2. **How universal is the "one XdhAB, both oxidations" model?** It is well supported for *R. capsulatus* ([PMID: 9515710](#)) but should not be generalized. Many bacteria encode multiple XO-family paralogs (18 Mo-enzyme loci in *S. novella*; [PMID: 23310928](#)), and it is unclear how often a single enzyme, versus dedicated hypoxanthine- and xanthine-specific enzymes, serves the two steps.
 3. **Sequence-based functional assignment is unreliable.** With ~70% of *S. novella* Mo-enzymes lacking characterized relatives ([PMID: 23310928](#)), and with the bacterial XdhAB being paradoxically more eukaryote-like in sequence than other bacterial Mo-enzymes ([PMID: 9515710](#)), assigning purine-oxidation function from homology alone risks error. Direct biochemistry remains necessary.
 4. **Cofactor form and its functional consequence.** Why some family members use bare MPT (XdhB) and others MCD (PaoABC, MOP) or MGD, and how the extra [4Fe-4S] cluster of PaoABC tunes electron flow, are only partly understood ([PMID: 27622978](#)).
 5. **Boundary with downstream metabolism.** The regulatory logic coupling urate *production* (this module) to urate *consumption* (AllR/RutR networks; [PMID: 18957590](#)) is organism-specific and incompletely mapped.
 6. **Indirect and cross-organism evidence.** Much of the broader literature is mammalian/clinical (hyperuricemia models, probiotic urate degradation), which informs urate physiology but not bacterial pathway mechanism; mixing these sources risks overgeneralization.
-

8. Mechanistic Model — Consolidated

NITROGEN-LIMITED BACTERIUM (purines as N source)

SUPPLY BRANCHES (conditional, substrate-gated, non-exclusive)



BIOGENESIS (accessory, non-catalytic): XdhC inserts Moco into XdhAB;
PaoD matures PaoABC.

ARCHITECTURE varies: 1 chain (eukaryotic XDH) | 2 chains (XdhAB α₂β₂) | 3 chains (PaoABC αβγ)
CAVEAT: Klebsiella has a Moco-INDEPENDENT hypoxanthine route → model is realization-specific.

9. Evidence Base

PMID	Paper (abbrev.)	How it supports / challenges the model
9515710	<i>R. capsulatus</i> XDH more similar to eukaryotic counterparts	Establishes two-subunit XdhA (FAD+2×[2Fe-2S]) / XdhB (Moco) division; hypoxanthine/xanthine induction; eukaryote-like sequence (F001, F006)
10217763	Role of XdhC in Moco insertion	Confirms $\alpha_2\beta_2$ active enzyme; XdhC is a Moco-insertase chaperone, not a subunit (F002)
11101664	<i>B. subtilis</i> guanine deaminase (yknA)	Guanine→xanthine as branch entry; enzyme non-homologous to mammalian GDA; purine/nitrogen induction (F003, F006)
31283204	Guanine deaminase substrate selectivity	Places GDEase in Zn-dependent amidohydrolase superfamily; universal guanine→xanthine reaction (F003)
8799115	Structure-based XO-family mechanism	Defines Mo-hydroxylation chemistry (H ₂ O-derived OH ⁻ , Glu-869, hydride to Mo=S) common to both oxidations (F004, F006)
1400180	<i>Klebsiella</i> Moco mutants use hypoxanthine	Challenges universality: Moco-independent hypoxanthine route exists (F004)
27622978	<i>E. coli</i> PaoABC	Three-subunit XO-family enzyme with MCD + extra [4Fe-4S]; architectural variation (F005)
24498065	PaoD chaperone	Dedicated Moco-maturation chaperone per architecture, paralleling XdhC (F005)
24492481	PaoABC SAXS/crystallography	Confirms $\alpha\beta\gamma$ heterotrimer that does not dimerize via Mo-domain
11713686 / 27520791	<i>D. gigas</i> MOP at 1.28 Å	Atomic-resolution XO-family prototype: MCD, 2×[2Fe-2S], Mo water ligand, active-site water chain
10704312	<i>D. desulfuricans</i> MOD	Conserved active site/water chain across the family (ancestral scaffold)
15030483	DgAOR direct electrochemistry	Redox behavior of Fe-S and Mo cofactors (electron relay)
19452052		

PMID	Paper (abbrev.)	How it supports / challenges the model
	Mo/W crystallographic overview	Three Mo-enzyme families; situates XO family and boundaries
15311335	Bis-MGD nitrate reductases/FDHs	Defines separate DMSO-reductase-family enzymes to exclude
23310928	<i>S. novella</i> molybdoproteome	Family expansion; caution on sequence-based function assignment
39467736	Plant XDH review	Cross-kingdom conservation of net chemistry; broader physiological roles
18957590	AllR/AllS/RutR regulation	Downstream urate degradation is a separate regulon (boundary)

10. Limitations and Knowledge Gaps

- **Single-organism anchoring.** The two-subunit "one enzyme does both oxidations" model rests principally on *R. capsulatus*; genome-wide, most bacteria encode multiple XO-family paralogs whose individual substrate assignments are unverified.
- **Unidentified alternative route.** The molecular basis of the *Klebsiella* Moco-independent hypoxanthine utilization is unknown — a concrete gap that directly qualifies the model.
- **Guanine branch under-characterized in the terminal-oxidation context.** Most guanine-deaminase structural work is not from organisms in which the full guanine→xanthine→urate flux has been dissected; branch-to-terminal coupling is inferred, not directly measured in one system.
- **Regulatory coupling.** How induction of the supply branches is coordinated with terminal-step expression and with downstream urate catabolism varies by organism and is not comprehensively mapped.
- **Literature imbalance.** A substantial fraction of retrievable "urate" literature is mammalian/clinical and does not bear on bacterial mechanism; conclusions about bacterial architecture rest on a relatively small set of primary studies.

11. Proposed Follow-up Experiments / Actions

1. **Identify the *Klebsiella* Moco-independent hypoxanthine enzyme.** Combine transposon mutagenesis or Tn-seq under hypoxanthine-as-sole-N selection with biochemistry to pin down the responsible activity and test whether it produces xanthine or bypasses it.
 2. **Test the "one XdhAB, both oxidations" claim across lineages.** Purify XdhAB from several taxa and directly assay both hypoxanthine→xanthine and xanthine→urate kinetics on the same enzyme; determine how often dedicated paralogs instead split the two reactions.
 3. **Reconstitute the convergent module in vitro.** Combine recombinant Zn-guanine deaminase and XdhAB (matured with XdhC) to demonstrate flux from both guanine and hypoxanthine to urate in a defined system, quantifying branch throughput.
 4. **Structure–function of cofactor variants.** Compare MPT (XdhB) vs. MCD (MOP/PaoABC) enzymes and dissect the functional role of PaoABC's extra [4Fe-4S] cluster in electron routing.
 5. **Map regulatory coupling.** Use reporter fusions and ChIP/transcriptomics to define how nitrogen regulators coordinate the two supply branches, the terminal step, and downstream urate catabolism within a single organism.
 6. **Phylogenetic reconstruction.** Build a family-wide phylogeny anchored on the *Desulfovibrio* prototypes to identify the best ancestral representatives and trace when purine-specific substrate specialization and multi-subunit splitting arose.
-

*Prepared as a review-style synthesis. Claims are attributed to the primary literature cited; uncertainty is flagged explicitly, and single-organism results (notably from *R. capsulatus* and *K. pneumoniae*) should not be generalized to all bacteria without direct verification.*
