

Bacterial Pst Phosphate Uptake: A Reusable ABC-Import Module for High-Affinity Phosphate Scavenging

A review-style synthesis for a molecular-biology audience

1. Executive Summary

Inorganic phosphate (Pi) is an irreplaceable nutrient — the backbone of nucleic acids, the currency of energy metabolism, and a near-universal covalent regulator — yet it is frequently scarce in the environments bacteria inhabit. The **Pst system (phosphate-specific transport)** is the dedicated high-affinity machine that bacteria deploy to scavenge Pi when it is limiting. It is a canonical **Type I ATP-binding-cassette (ABC) importer** built from a small, conserved parts list: a periplasmic phosphate-binding protein (**PstS**) that captures Pi with exquisite selectivity; a pair of integral-membrane subunits (**PstC** and **PstA**) that form the translocation pathway; and a homodimeric cytoplasmic ATPase (**PstB**) that hydrolyzes ATP to power the transport cycle

([P 8918249](#)).

In *Escherichia coli* these are encoded together with a fifth gene, *phoU*, in the *pstSCAB-phoU* operon, whose transcription is switched on during phosphate starvation by the PhoR/PhoB two-component system

([P 12471449](#)),

([P 8432742](#)).

The system's boundaries are the crux of understanding it. The **transport module proper is PstSCAB. PhoU and the PhoR/PhoB two-component pair are an adjacent regulatory/homeostatic layer, not translocation steps**

([P 25220976](#)),

(
P 8432742

The low-affinity, proton-motive-force-driven **Pit transporter** is a **separate route** for Pi that is often conflated with Pst but is mechanistically and physiologically distinct

(
P 11489853

Complicating matters, PstS and the intact Pst complex have **moonlighting and sensory roles** — PstS acts as an adhesin/biofilm factor in pathogens

(
P 25223609

),
P 25359607

and loss of a functional Pst complex constitutively derepresses the whole Pho regulon, so many "*pst* phenotypes" reflect signaling rather than transport

(
P 11489853

),
P 10629178

Mechanistically, the best-supported model is borrowed from the well-characterized Type I prototypes (maltose, molybdate, methionine importers): liganded PstS docks onto the periplasmic face of PstC/PstA and, together with ATP binding at the PstB dimer, drives an alternating-access conformational cycle; ATP hydrolysis resets the transporter and releases Pi to the cytoplasm

(
P 19395376

),
P 21953468

The **NBD/ATPase motor (PstB)** is **ancient and deeply conserved** across the entire ABC superfamily, whereas the **transmembrane fold (PstA/PstC)** is the **more lineage-variable element**, and the **periplasmic capture protein is the most plastic part** — subject to paralog expansion, replacement (e.g., by PhoX), or outright loss in some operons

(
P 32978974

),
P 8628229

This review lays out the system's parts, the sequence of events, its variation and origin, the physical constraints on the pathway, and the points where the literature remains genuinely uncertain.

2. Definition and Biological Boundaries

2.1 What the system *is*

The Pst system is a **binding-protein-dependent (Type I) ABC importer specific for inorganic phosphate**. Its irreducible functional unit is four proteins (

P 8918249

Component	Location	Activity	Role in the pathway
PstS	Periplasm (or lipoprotein-anchored in mycobacteria/spirochetes)	Pi binding	Capture and presentation of substrate
PstC	Inner membrane (TMD)	Permease	Forms the translocation pathway; docks PstS
PstA	Inner membrane (TMD)	Permease	Forms the translocation pathway; docks PstS
PstB	Cytoplasmic face (NBD), homodimer	ATP hydrolysis	Energy coupling / power stroke

Functionally, the pathway is a strict three-step sequence: **(1) periplasmic capture** by PstS → **(2) phosphate-selective translocation** through the PstC/PstA membrane channel → **(3) ATP-dependent energy coupling** by the PstB dimer, whose nucleotide cycle drives steps 1–2. This maps directly onto the commissioned biological outline.

2.2 What is *adjacent* and often confused

Three neighboring processes are routinely conflated with the transporter and should be kept conceptually separate:

- 1. The Pho regulon signaling apparatus (PhoR/PhoB) and PhoU.** PhoR is the membrane histidine kinase and PhoB the response regulator; under Pi limitation PhoR phosphorylates PhoB, which activates Pho-box promoters, and under Pi excess PhoR, Pst and PhoU together shut the regulon off

P 8432742

PhoU is a negative regulator of Pi signaling that modulates transport through the Pst proteins via PhoR/PhoB (P 25220976).

These are *regulatory* events; they turn the transporter's gene *expression* on and off and communicate Pi status to the cell, but they do not themselves move phosphate across the membrane. The Pst complex participates in **sensing** (see §6), which is why *phoU* is co-transcribed in the operon, but sensing and translocation are separable activities.

2. **The Pit low-affinity transporter.** *E. coli* has a second major Pi-uptake route, **PitA**, that is **constitutively expressed and driven by the proton-motive force** — a secondary transporter, not an ABC system — and a related PitB (P 11489853).

Pit handles bulk Pi uptake when phosphate is plentiful; Pst is the inducible, high-affinity scavenger for starvation. Reports that do not distinguish the two can misattribute uptake defects.

3. **Downstream phosphate storage and homeostasis (polyphosphate).** Once imported, Pi feeds central metabolism and can be polymerized into inorganic polyphosphate (polyP) by polyphosphate kinases for storage and stress buffering (P 41864548).

PolyP synthesis is especially important during recovery from starvation (the "overplus"/nutrient-upshift response) (P 42234530).

This is a separate metabolic fate of imported phosphate, not part of the transport step.

2.3 Competing definitions

The main definitional tension in the literature is whether "the Pst system" means the **four-protein transporter** or the **five-gene operon including phoU** (or even the broader Pi-signaling module PhoR/PhoB/PhoU/Pst). For a mechanistic review the transporter-centric definition (PstSCAB) is the most defensible: it isolates the biochemical activity (Pi translocation) from the regulatory functions layered on top. We adopt that boundary throughout, while noting that the physical Pst complex is genuinely bifunctional (transport + sensing).

3. Mechanistic Overview

3.1 The best current model of the sequence of events

Because there is not yet a high-resolution, multi-conformation structural series for the assembled PstSCAB complex itself, the mechanistic model is built by analogy to the extensively characterized Type I ABC importers — maltose (MalFGK₂-E), molybdate (ModBC-A), and methionine (MetNIQ) — supplemented by direct biochemical data on Pst subunits. The consensus alternating-access cycle is:

1. **Capture (obligatory).** Periplasmic PstS binds a single Pi ion in a cleft between two lobes and closes around it (a "Venus-flytrap" motion). Binding is high-affinity and highly selective (§3.2).
2. **Docking (obligatory).** Liganded, closed PstS docks onto the periplasmic surface of the PstC/PstA transmembrane dimer. In the maltose prototype, the binding protein remains associated with the transporter throughout the cycle rather than dissociating each round, and its liganded state is communicated across the membrane to the ATPase (P 19395376).
3. **ATP binding and gate opening (obligatory, energy-requiring).** ATP binding closes the two PstB nucleotide-binding domains into a sandwich dimer. This closure is mechanically transmitted through coupling helices to the PstC/PstA gates, switching the permease toward an outward-facing state that accepts Pi from PstS. Direct evidence that nucleotide binding drives a large, physiological conformational change in Pst's ATPase comes from *M. tuberculosis* PstB, where ATP (or a non-hydrolyzable analog) triggers a global change involving movement of an α -helical subdomain (Arg137–Trp150) relative to the core (P 15936994).
4. **Translocation and hydrolysis (obligatory).** Pi passes into the transmembrane pathway; ATP hydrolysis and Pi/ADP release open the NBD dimer, resetting the permease to an inward-facing state and delivering phosphate to the cytoplasm. Following hydrolysis the translocation gates return to an apo-like conformation (P 21953468).
5. **Reset.** The transporter returns to its resting state, ready to accept another liganded PstS.

Steps 1–4 are **obligatory and ordered**: capture must precede docking, and productive gating requires both a liganded binding protein and the nucleotide cycle. Regulation of gene expression (PhoR/PhoB/PhoU) is **conditional** — it determines whether the transporter is present at all under a given Pi regime but is not part of the per-molecule transport cycle. Moonlighting activities (adhesion, biofilm) are **accessory** and lineage-specific (§5).

3.2 Why the capture step is so selective

Phosphate selectivity, not just affinity, defines the system. PstS-family binding proteins bury Pi in a dense network of hydrogen bonds — for example, *Clostridium perfringens* PBP-1 forms an unusually high number (14) of hydrogen bonds with the anion

([P 25338617](#))

— and residues implicated in Pi binding are conserved even in distantly related homologs such as *Borrelia burgdorferi* BbPstS (1.3 Å structure)

([P 24318969](#))

The most striking demonstration of selectivity is discrimination against **arsenate**, a near-perfect steric and electronic mimic of phosphate: all tested PBPs reject arsenate by ≥ 500 -fold, and a PBP from the arsenate-rich Mono Lake strain GFAJ-1 reaches $\sim 4,500$ -fold via a distinctive binding geometry resolved at sub-ångström resolution

([P 23034649](#))

This selectivity is achieved by binding-site geometry and hydrogen-bond angle rather than net charge alone, which is what allows the protein to reject the similarly-charged sulfate and arsenate anions.

3.3 Energy coupling

The PstB homodimer is the motor. Like all ABC-transporter NBDs it uses the Walker A/B motifs and the ABC signature motif to sandwich two ATP molecules at the dimer interface; ATP-driven dimer closure and hydrolysis-driven opening are converted into the alternating-access motion of the permease. The Pst-specific evidence that this is a real, nucleotide-driven conformational cycle (rather than an inference from static structures) comes from solution studies of *M. tuberculosis* PstB

([P 15936994](#))

Importantly, coupling is **not universally identical** across the ABC superfamily: exporters, small importers and large importers use measurably different coupling schemes

(
P 21967052
),

so quantitative details (e.g., stoichiometry of ATP per Pi) should be transferred to Pst with caution.

4. Major Molecular Players and Active Assemblies

- **PstS (phosphate-binding protein).** A bilobed periplasmic protein of the cluster-D/ phosphate-binding-protein family. In Gram-negatives it is a soluble periplasmic protein; in Actinobacteria and spirochetes it is a **lipoprotein** tethered to the outer face of the membrane (e.g., *M. tuberculosis* PstS-1, a 38 kDa mannosylated glycolipoprotein; *B. burgdorferi* BB0215)

(
P 25359607
),

P 24318969
).

It is the specificity determinant of the whole system.

- **PstC and PstA (permease pair).** Two homologous but non-identical integral-membrane subunits that together build the translocation pathway and provide the periplasmic docking surface for PstS. Their asymmetry (two different genes rather than a homodimer) is a general feature of Type I importers. In some lineages these subunits carry extra hydrophilic domains not present in *E. coli* (e.g., *P. aeruginosa* PstC/PstA)

(
P 8628229
).

- **PstB (ABC ATPase).** The nucleotide-binding, ATP-hydrolyzing subunit; functions as a homodimer. It is the most conserved component and the best representative of the ancestral ABC "motor"

(
P 8918249
),

P 15936994
).

- **The assembled transporter.** The functional stoichiometry, by analogy to Type I importers, is **PstS : PstC : PstA : PstB₂** (one binding protein, one heterodimeric permease, one ATPase homodimer). Direct high-resolution structural confirmation of the full assembled *E. coli* PstSCAB complex remains a gap in the field (see §7).

- **Accessory/regulatory proteins (outside the transport module).** PhoU (negative regulator; dimeric three-helix-bundle fold)

(
P 25220976);

PhoR (sensor histidine kinase) and PhoB (response regulator, Pho-box binding)

(
P 8432742).

5. Evolutionary and Cell-Biological Variation

5.1 Deep conservation and origin

The Pst system is ancient. Its ATPase belongs to the **ABC superfamily, one of the largest and most ancient protein families**, present in all three domains of life. Across the superfamily the **NBDs are highly conserved** while the **TMDs adopt distinct folds** — evidence that ancient motor domains were repeatedly combined with different transmembrane systems, and the basis for classifying ABC transporters by their TMD fold

(
P 32978974).

Pst falls in the **Type I importer** class alongside the maltose, molybdate and methionine transporters, and is mechanistically distinct from Type II importers such as the vitamin-B₁₂ transporter BtuCD, in which ATP binding drives the gates in the opposite direction

(
P 21953468).

Thus:

- **Most ancient/conserved:** the PstB ATPase (NBD motor) — the best family member for reasoning about the ancestral energy-coupling role.
- **Intermediate:** the PstA/PstC permease fold — conserved within Type I importers but variable across the wider superfamily.
- **Most plastic:** the periplasmic PstS capture protein — the element most subject to lineage-specific gain, loss and replacement.

5.2 Lineage-specific variation

- **Operon architecture varies.** *E. coli*: *pstS-pstC-pstA-pstB-phoU*

(
P 12471449
).

P. aeruginosa: *pstC-pstA-pstB-phoU* — **notably lacking *pstS* in the operon**, though the three membrane/ATPase genes are absolutely required for Pi transport

(
P 8628229
).

Mycobacteria carry a **rearranged operon** (*pstB*, *pstS-1*, *pstC-1*, *pstA-2*) and **multiple PstS paralogs** (*PstS-1/-2/-3*)

(
P 8918249
).

- **Alternative capture proteins.** *Xanthomonas citri* encodes two periplasmic Pi-binding proteins — an ABC-type PstS and a separate PhoX associated with a putative porin — indicating parallel/alternative uptake routes to the same end

(
P 25484207
).

Where *pstS* is absent from the operon, an unlinked binding protein presumably supplies the capture function.

- **Membrane topology of PstS.** Soluble periplasmic protein in classic Gram-negatives vs. surface lipoprotein in Actinobacteria/spirochetes

(
P 25359607
),

P 24318969
).

5.3 Physiological-state and "cell-type" variation

Because bacteria lack tissues, the analog of cell-type variation is **physiological state and developmental program**:

- **Starvation induction.** Pst is OFF when Pi is replete and strongly induced during Pi limitation via the Pho regulon

(
P 12471449
),

P 8432742
).

- **Developmental coupling.** In *Caulobacter crescentus*, Pi status routed through the Pst/Pho system controls **stalk elongation** — Pi-starved cells make stalks up to ~30× longer, and this response requires PhoB
([P 10629178](#)).
- **Virulence programs.** In several pathogens Pi limitation, sensed partly through Pst, triggers virulence: intestinal Pi depletion shifts *P. aeruginosa* to a lethal phenotype with 32-fold-elevated PstS
([P 18656625](#)),
and in uropathogenic *E. coli* a *pst* lesion reprograms c-di-GMP signaling and type 1 fimbriae
([P 28924030](#)).
- **Moonlighting at the cell surface.** PstS has been co-opted for functions independent of transport: in *P. aeruginosa*, phosphate binding and biofilm formation are genetically separable (a binding-dead point mutant still forms biofilm; an N'-loop truncation loses biofilm but retains uptake)
([P 25223609](#));
in *M. tuberculosis*, PstS-1 is an adhesin that engages the macrophage mannose receptor and promotes phagocytosis
([P 25359607](#)).

6. Constraints, Dependencies, and Failure Modes

6.1 Obligatory ordering

- **Capture precedes translocation.** PstS must bind Pi and adopt its closed, liganded conformation before productive docking onto PstC/PstA; empty binding protein does not efficiently stimulate the transporter.
- **Docking + ATP are jointly required.** Neither a liganded binding protein alone nor ATP alone completes a transport stroke; the alternating-access cycle requires both the periplasmic docking signal and the cytoplasmic nucleotide cycle, which are reciprocally

coupled across the membrane

(
P 19395376
).

- **Hydrolysis resets, it does not "pull."** ATP *binding* (NBD closure) drives the outward-facing transition; *hydrolysis* and product release reset the transporter

(
P 21953468
).

This ordering rules out simple models in which hydrolysis directly drags substrate inward.

6.2 Substrate-specificity constraint

The capture step is a **discrimination checkpoint**. PstS actively excludes arsenate (≥ 500 -fold) and sulfate despite their mimicry

(
P 23034649
).

This constraint is why the pathway is "phosphate-specific" — selectivity is set at capture, upstream of translocation.

6.3 Compartmental constraints

The system is intrinsically **two-membrane-aware** in diderms: substrate must first cross the outer membrane (via porins) to reach periplasmic PstS, which then feeds the inner-membrane permease. In monoderms and in lipoprotein-anchored variants, PstS is surface-tethered rather than freely periplasmic, changing the geometry of capture

(
P 24318969
,

P 25359607
).

6.4 Failure modes and the transport-vs-signaling trap

The most important interpretive pitfall: **loss of a functional Pst complex constitutively activates the Pho regulon**, mimicking starvation even when Pi is abundant (the *E. coli* Δ pstC345 phenotype)

(
P 11489853
).

Consequently, many phenotypes of *pst* mutants — Caulobacter hyper-stalks, altered fimbriation, biofilm and virulence changes — are **downstream consequences of Pho derepression, not direct transport defects**

(
P 10629178

P 28924030

pst and *phoB* mutants often show opposite phenotypes, precisely because one derepresses and the other abolishes the regulon

(
P 10629178

Rigorous attribution of a phenotype to "phosphate transport" therefore requires separating the transporter's uptake function from its sensory function — for example with transport-dead but signaling-competent alleles, or with epistasis against *phoB*.

7. Controversies and Open Questions

1. **Mechanism of PhoU.** PhoU is genetically a central negative regulator that couples the Pst transporter to PhoR/PhoB, yet its biochemical mechanism "is still unknown" even after its structure was solved

(
P 25220976

How PhoU physically transmits Pi-replete status from the Pst complex to PhoR (and whether it acts as a molecular "brake," a scaffold, or via a metabolite such as ATP/polyP) remains unresolved.

2. **How is Pi status actually sensed?** The Pst complex is required to keep the Pho regulon off, but the identity of the sensed signal — occupancy of PstS, a conformational state of PstB, an intracellular Pi/ATP ratio, or PhoU-mediated contacts — is not settled. Much of the evidence is genetic and indirect.

3. **Absence of a full Pst complex structure.** The mechanistic model is largely transferred from maltose/molybdate/methionine importers. There is direct structural information on isolated PstS proteins (many species) and biochemical data on PstB, but a high-resolution, multi-conformation structural series of an **assembled PstSCAB** transporter — the direct test of the borrowed alternating-access model — is still lacking. Coupling schemes are known to differ across ABC classes

(
P 21967052

so extrapolation carries genuine uncertainty.

4. **The arsenate / "arsenic life" debate.** The GFAJ-1 claim of arsenate-for-phosphate substitution is contradicted by the strong arsenate discrimination of PstS-family proteins ([P 23034649](#)); computational work shows arsenate *can* substitute kinetically but gives less stable products ([P 23789648](#)). The transporter's selectivity is a key line of evidence, but the broader question of arsenate handling in high-arsenate niches is not fully closed.

5. **Moonlighting vs. transport in pathogenesis.** How much of PstS's contribution to virulence/biofilm is due to surface adhesion functions versus phosphate acquisition versus Pho-regulon signaling is entangled and organism-specific ([P 25223609](#) , [P 25359607](#) , [P 18656625](#)). Data from *E. coli*, *P. aeruginosa* and *M. tuberculosis* should not be pooled uncritically.

6. **Cross-regulation and organism transferability.** In *E. coli* the Pho regulon can be activated Pi-independently (e.g., by CreC or acetyl phosphate) ([P 8432742](#)), and regulatory wiring differs across taxa (e.g., *Streptococcus pneumoniae* PnpR-PnpS behaves differently from *E. coli*) ([P 9973337](#)). Mechanistic claims from one organism may not generalize.

Most important open questions: (i) a definitive assembled-complex structure and transport cycle for Pst specifically; (ii) the molecular mechanism of PhoU and the identity of the sensed phosphate signal; (iii) clean genetic separation of Pst's transport, sensing and moonlighting functions in virulence.

8. Key References

1. Agüena M, Yagil E, Spira B. *Transcriptional analysis of the pst operon of Escherichia coli*. Mol Genet Genomics. 2002.

P 12471449

— Operon structure (*pstSCAB-phoU*), starvation induction, post-transcriptional processing.

2. Wanner BL. *Gene regulation by phosphate in enteric bacteria*. J Cell Biochem. 1993.

P 8432742

— Pho regulon; roles of PhoR, PhoB, Pst, PhoU; cross-regulation.

3. Braibant M, et al. *A Mycobacterium tuberculosis gene cluster encoding proteins of a phosphate transporter homologous to the E. coli Pst system*. 1996.

P 8918249

— Subunit assignments (PstB ATPase; PstA/PstC permease; PstS binding); mycobacterial paralogs/operon.

4. Harris RM, et al. *Characterization of PitA and PitB from Escherichia coli*. J Bacteriol. 2001.

P 11489853

— Pit (low-affinity, PMF-driven) vs Pst (high-affinity, ABC); Δ pstC constitutively activates Pho regulon.

5. Thomas C, et al. *Structural and functional diversity calls for a new classification of ABC transporters*. FEBS Lett. 2020.

P 32978974

— Conserved NBD motor vs divergent TMD folds; evolutionary framing.

6. Joseph B, et al. *Transmembrane gate movements in the type II ABC importer BtuCD-F during nucleotide cycle*. J Biol Chem. 2011.

P 21953468

— Type I vs Type II importer mechanisms; alternating access.

7. Grote M, et al. *Transmembrane signaling in the maltose ABC transporter MalFGK2-E*. J Biol Chem. 2009.

P 19395376

— Type I prototype: binding protein stays docked and signals substrate availability to the ATPase.

8. Al-Shawi MK. *Catalytic and transport cycles of ABC exporters*. Essays Biochem. 2011.

P 21967052

— Coupling mechanisms are not universally conserved across ABC classes.

9. Gupta S, et al. *Nucleotide-induced conformational change in the catalytic subunit (PstB) of the phosphate-specific transporter from M. tuberculosis*. 2005.

P 15936994

— Direct evidence that ATP drives a global conformational change in Pst's ATPase.

10. Elias M, et al. *The molecular basis of phosphate discrimination in arsenate-rich environments*. Nature. 2012.

P 23034649

— PstS-family proteins discriminate Pi over arsenate ≥ 500 -fold (GFAJ-1 PBP $\sim 4,500$ -fold).

11. Gonzalez D, et al. *Crystal structure of the phosphate-binding protein (PBP-1) of an ABC-type phosphate transporter from Clostridium perfringens*. 2014.

P 25338617

— Dense hydrogen-bond network underlying phosphate capture.

12. Brautigam CA, et al. *Structural analyses of the PstS lipoprotein (BB0215) from Borrelia burgdorferi*. 2014.

P 24318969

— Conserved PBP fold/binding residues; lipoprotein-anchored PstS.

13. Nikata T, et al. *Molecular analysis of the pst operon of Pseudomonas aeruginosa*. 1996.

P 8628229

— Operon lacking *pstS*; extra hydrophilic domains in PstC/PstA.

14. Lee SJ, et al. *Crystal structure of PhoU from Pseudomonas aeruginosa, a negative regulator of the Pho regulon*. 2014.

P 25220976

— PhoU as negative regulator; mechanism still unknown.

15. Neznansky A, et al. *The P. aeruginosa PstS plays a phosphate-independent role in biofilm formation*. 2014.

P 25223609

— Genetic separation of PstS transport and biofilm/moonlighting functions.

16. Esparza M, et al. *PstS-1, the 38-kDa M. tuberculosis glycoprotein, is an adhesin*. 2015.

P 25359607

— PstS-1 binds the macrophage mannose receptor; moonlighting.

17. Long J, et al. *Depletion of intestinal phosphate activates the virulence of P. aeruginosa*. 2008.

P 18656625

— Pi limitation → lethal virulence; PstS upregulation.

18. Gonin M, et al. *Regulation of stalk elongation by phosphate in Caulobacter crescentus*. 2000.

P 10629178

— *pst* mutants derepress Pho regulon; developmental coupling; opposite *pst* vs *phoB* phenotypes.

19. Crépin S, et al. *Altered regulation of YaiC reduces type 1 fimbriae in a Pst mutant of uropathogenic E. coli*. 2017.

P 28924030

— *pst* lesion reprograms c-di-GMP signaling/fimbriation.

20. Novak R, et al. *An ABC phosphate transporter and two-component system in Streptococcus pneumoniae*. 1999.

P 9973337

— Divergent regulatory wiring; caution on organism transferability.

21. Jissy AK, Datta A. *Can arsenates replace phosphates in natural biochemical processes? A computational study*. 2013.

P 23789648

— Arsenate can substitute kinetically but yields less stable products.

22. Krukenberg V, et al. *Polyphosphate metabolism: enzymatic pathways and regulation*. 2026.

P 41864548

White G, et al. *Polyphosphate synthesis is essential for phosphate and ATP homeostasis during nutrient upshift*. 2026.

P 42234530

— Downstream polyP storage/homeostasis (adjacent process).

A note on scope and uncertainty

This synthesis deliberately draws mechanistic detail from Type I ABC-importer prototypes (maltose, molybdate, methionine) where direct assembled-Pst structural data are lacking, and it flags where evidence is genetic/indirect (PhoU mechanism, Pi sensing) or organism-specific (moonlighting, regulatory wiring). Claims about the transport cycle are strongest for the general ABC framework and for PstS capture/selectivity; they are weakest for the quantitative energetics of the Pst complex specifically and for the molecular logic that couples the transporter to Pho-regulon signaling.

Generated by OpenScientist — Scientific Hypothesis Agent for Novel Discovery