

Histidine Catabolism to Glutamate: A Conserved Mosaic Core with Three Convergent Endings

A commissioned review-style synthesis for a molecular biology audience

1. Executive Summary

Histidine catabolism to L-glutamate is one of the most widely distributed amino-acid degradation systems in biology, present in bacteria, archaea, fungi, plants, and animals. Its central logic is deceptively simple: three enzymatic steps convert L-histidine, through *trans*-urocanate and 4-imidazolone-5-propanoate, to N-formimidoyl-L-glutamate (formiminoglutamate, FIGLU); a fourth "terminal" step then removes the remaining formimino carbon to release free L-glutamate. Glutamate is a hub metabolite, so the pathway is simultaneously a route for nitrogen and carbon salvage, a source of one-carbon units, and, in specialized tissues, a producer of a UV-absorbing chromophore.

The central scientific insight that emerges from this investigation is that the pathway is a **mechanistic mosaic rather than a single ancestral module**. The three conserved core enzymes — histidine ammonia-lyase (HAL/HutH), urocanase (HutU/UROC1), and imidazolonepropionase (HutI/AMDHD1) — belong to three completely unrelated protein superfamilies with distinct folds, cofactors, and catalytic chemistries: an autocatalytic MIO-cofactor ammonia-lyase, a structurally unique NAD⁺-dependent hydratase that uses NAD⁺ as a non-redox electrophile, and a metal-dependent amidohydrolase. The pathway was therefore assembled from independently evolved parts that happen to constitute a chemically coherent sequence. This mosaic character extends to the terminal step, where **three mutually exclusive and convergently evolved routes** dispose of the formimino group: a two-step bacterial HutF/HutG route releasing ammonium then formate, a one-step arginase-superfamily formiminoglutamase releasing formamide, and the vertebrate folate-coupled formiminotransferase-cyclodeaminase (FTCD) route that channels the carbon onto tetrahydrofolate and into one-carbon metabolism.

Around this canonical spine sit several processes that share intermediates but should be treated as separate boundary systems: the anaerobic diversion of urocanate to imidazole propionate by urocanate reductase (UrdA) — a bacterially produced cardiometabolic-risk metabolite; the epidermal production of photoprotective urocanic acid by HAL; and the moonlighting role of vertebrate FTCD as the Golgi/centrosome microtubule-binding protein "58K"/p60. The pathway also has direct clinical stakes in humans: loss of HAL causes histidinemia, loss of FTCD causes glutamate formiminotransferase deficiency (formiminoglutamic aciduria), and flux through the folate-coupled branch drains the tetrahydrofolate pool and modulates sensitivity to antifolate chemotherapeutics such as methotrexate.

2. Definition and Biological Boundaries

What is included

The biological system under review is the enzymatic degradation of L-histidine to L-glutamate — the "Hut" (histidine utilization) pathway in the bacterial nomenclature. It comprises four logical steps:

1. **Histidine deamination** — HAL/HutH removes the α -amino group as ammonia to yield *trans*-urocanate.
2. **Urocanate hydration** — HutU/UROC1 (urocanase) adds water across the imidazole ring double-bond system to yield 4-imidazolone-5-propanoate.
3. **Imidazolone ring opening** — HutI/AMDHD1 (imidazolonepropionase) hydrolytically opens the ring to give N-formimidoyl-L-glutamate (FIGLU).
4. **Terminal FIGLU processing** — one of three alternative routes removes the formimino carbon to release L-glutamate.

The first three reactions form a strictly conserved, obligatory, and ordered core; the fourth is a branch point with three distinct molecular solutions.

What should be treated separately (boundary processes)

Several neighbouring processes are frequently conflated with canonical histidine catabolism but are mechanistically and physiologically distinct:

- **Anaerobic urocanate reduction to imidazole propionate (ImP).** In gut and other anaerobic bacteria, urocanate reductase (UrdA), an FAD-dependent enzyme, reduces urocanate to imidazole propionate rather than allowing urocanase to hydrate it. This diverts the pathway at the urocanate node, uses urocanate as a respiratory electron acceptor, and produces a metabolite implicated in insulin resistance and cardiometabolic disease. It is a *competing* fate of a shared intermediate, not a step in glutamate production.
- **Epidermal urocanic-acid photoprotection.** In mammalian skin, HAL produces urocanic acid that accumulates in the stratum corneum as a UVB chromophore. Here HAL is deployed for a structural/photobiological function rather than for net histidine degradation to glutamate.
- **FTCD moonlighting in cytoskeletal/Golgi cell biology.** Vertebrate FTCD is identical to the "58K" microtubule-binding Golgi protein and chicken p60; it binds microtubules and localizes to the centrosome. This is a second, non-catabolic function of a catabolic enzyme.
- **One-carbon (folate) metabolism proper.** The FTCD route feeds 5,10-methenyl-tetrahydrofolate into the folate one-carbon pool. That downstream pool — and its many biosynthetic uses — is a distinct system that the pathway *feeds*, not part of the histidine-degradation core.


Competing definitions in the literature

The main definitional ambiguity concerns whether the terminal FTCD branch "belongs" to histidine catabolism or to folate metabolism, since FTCD is usually discussed in the folate literature and its deficiency is classified as an inborn error of *folate* metabolism. Similarly, the "one-step formiminoglutamase" route is sometimes described in older biochemistry as the canonical terminal step (particularly in organisms and tissues where it operates), whereas the bacterial HutF/HutG two-step route dominates the microbiological literature. These are best understood as alternative, lineage-specific solutions rather than as competing definitions of the same reaction.

3. Mechanistic Overview

The conserved core (Finding F001)

Genetic and biochemical characterization of the *hut* loci in *Bacillus subtilis* and *Pseudomonas putida* established the conserved ordered enzyme set: **histidase (HutH/HAL) → urocanase (HutU/UROC1) → imidazolonepropionase (HutI/AMDHD1)**, converting L-histidine via *trans*-urocanate and 4-imidazolone-5-propanoate to FIGLU. In *B. subtilis*, the operon assigns *hutU* to urocanase and *hutI* to imidazolone-5-propionate hydrolase [PMID: 7704263]. In *P. putida*, the mapped gene order is *hutG-hutI-hutH-hutU-hutC-hutF* [PMID: 2842309]. The step order is chemically obligatory: each enzyme consumes the product of the preceding one.



Step 1 — Histidine ammonia-lyase (Findings F002, F006)

HAL catalyzes the non-oxidative deamination of histidine to urocanate using a protein-derived cofactor, **3,5-dihydro-5-methylidene-4H-imidazol-4-one (MIO)**, formed autocatalytically by cyclization and dehydration of an internal Ala-Ser-Gly tripeptide. HAL is structurally and mechanistically homologous to phenylalanine ammonia-lyase (PAL), tyrosine ammonia-lyase, and MIO-dependent aminomutases such as the tyrosine 2,3-aminomutase from the C-1027 enediyne pathway [PMID: 15350127;

1.

P 17516659

The MIO acts as an electrophilic trap that activates the substrate for β -elimination of ammonia.

Step 2 — Urocanase (Finding F003)

Urocanase is a symmetric homodimer of ~557-residue subunits with a tightly, non-covalently bound **NAD⁺ cofactor** [PMID: 15313616]. Uniquely, NAD⁺ here is *not* a redox carrier. Isotope-labeling, inhibitor, and deuterium-isotope studies show NAD⁺ acts as an **electrophile**: it forms a covalent C–C adduct with the imidazole ring, activating the conjugated double-bond system for water addition to yield 4-imidazolone-5-propanoate [PMID: 6110440; [P 2885981](#)].

Deuterium isotope data specifically rule out an internal hydride-transfer / oxidation-reduction mechanism, distinguishing urocanase from the vast majority of NAD⁺-dependent enzymes.

Step 3 — Imidazolonepropionase (Finding F008)

HutI/AMDHD1 is a member of the **amidohydrolase superfamily**, built on the canonical (β/α)₈ TIM-barrel fold with a mononuclear metal center (iron) coordinated by four histidines and one aspartate, and a proposed His265 catalytic base [PMID: 18442260]. It hydrolytically opens the imidazolone ring to yield FIGLU. AMDHD1 is the human ortholog.

Step 4 — Terminal processing (Findings F004, F005, F009)

After HutI, FIGLU is processed by one of three convergently evolved routes (detailed in §4).

4. Major Molecular Players and Active Assemblies

The three-superfamily mosaic core (Finding F008)

A central result of this review is that the three obligatory core enzymes derive from **distinct, non-homologous folds and superfamilies** — the pathway is a mechanistic mosaic, not a single ancestral module:

Step	Enzyme	Superfamily / fold	Cofactor / metal	Catalytic logic
1	HAL / HutH	MIO-dependent aromatic amino-acid ammonia-lyase (PAL/TAL family)	Autocatalytic MIO (from Ala-Ser-Gly)	Electrophilic amine elimination
2	Urocanase / HutU / UROC1	Structurally unique NAD ⁺ -binding fold, no close relatives	Tightly bound NAD ⁺ (non-redox)	NAD ⁺ as electrophile; water addition
3	Imidazolonepropionase / HutI / AMDHD1	Amidohydrolase superfamily, (β/α) ₈ barrel	Mononuclear Fe (4 His + 1 Asp)	Metal-activated hydrolysis

This mosaicism means that the pathway cannot have arisen by tandem duplication of a single ancestral enzyme; instead, three lineages of pre-existing chemistry were recruited into a coherent metabolic sequence.

The three convergent terminal routes (Findings F004, F005, F009)

The formimino carbon on FIGLU is removed by one of three mutually exclusive terminal solutions, each drawn from a *different* protein family — a striking example of convergent metabolic evolution:

Route	Enzyme(s)	Family	Products	Representative organisms
Two-step formate route	HutF (formiminoglutamate deiminase/hydrolase) → HutG (N-formylglutamate amidohydrolase/deformylase)	Amidohydrolase-related	NH ₄ ⁺ (step 1), then formate + L-glutamate (step 2)	<i>P. putida</i> , <i>Streptomyces coelicolor</i>
One-step formiminoglutamase	Formiminoglutamase	Arginase superfamily , binuclear Mn ²⁺	Formamide + L-glutamate	<i>Trypanosoma cruzi</i> and others
Folate-coupled FTCD	FTCD (bifunctional formiminotransferase + cyclodeaminase)	Formimidoyltransferase-cyclodeaminase	5-formimino-THF → 5,10-methenyl-THF + L-glutamate	Vertebrates

Two-step HutF/HutG route (F004). In *P. putida*, *hutF* encodes formiminoglutamate hydrolase (deiminase), which releases ammonium to give N-formyl-L-glutamate; notably *hutF* is transcribed in a direction opposite to that of the other *hut* genes [PMID: 2842309]. HutG then deformylates N-formyl-L-glutamate to release formate and L-glutamate. *Streptomyces coelicolor* dissimilates histidine via N-formyl-L-glutamic acid, confirming this route operates across divergent bacteria [PMID: 6129283].

One-step formiminoglutamase route (F009). Formiminoglutamase from *Trypanosoma cruzi* (TcFIGase) hydrolyzes FIGLU directly to L-glutamate + formamide in a single step. Its 1.85 Å crystal structure reveals an **arginase-superfamily fold with a binuclear manganese cluster** similar to that observed in arginase (one Mn(A) ligand is Asn114 rather than His); reconstitution with Mn²⁺ confers maximal activity [PMID: 24261485]. This enzyme is structurally and mechanistically unrelated to the amidohydrolase-family HutF/HutG route or to FTCD — independent evolutionary recruitment of a third chemistry for the same net transformation.

Folate-coupled FTCD route (F005). In vertebrates the terminal step is carried out by **formiminotransferase-cyclodeaminase (FTCD)**, a bifunctional enzyme that first transfers the formimino group of FIGLU to tetrahydrofolate (glutamate formiminotransferase activity), then cyclodeaminates 5-formimino-THF to 5,10-methenyl-THF (cyclodeaminase activity). FTCD assembles into a **circular octamer** that is required for both activities and for substrate channeling between the two active sites; the native octameric structure is required for expression of both activities and for channeling of intermediates [PMID: 3593701]. This is the only terminal route that couples histidine catabolism to one-carbon metabolism rather than simply discarding the formimino carbon. Its loss causes formiminoglutamic aciduria (see §6).

Regulation (Finding F007)

In *Pseudomonas* and related bacteria, the *hut* genes are induced by the pathway intermediate **urocanate**, acting through the **GntR-family repressor HutC**; global nitrogen (NtrBC) and carbon (CbrAB–Crc/Hfq) regulators are superimposed. In *P. putida*, *hutF* and *hutU* (urocanase) expression is induced by urocanate [PMID: 2842309]. In *P. fluorescens*, HutC both mediates substrate-specific induction and fine-tunes *ntrBC* autoactivation, so that the local transcription factor HutC plays a crucial role in governing NtrBC to maintain carbon/nitrogen homeostasis [PMID: 33675669]. *B. subtilis* uses a distinct positive regulator (HutP) and layers catabolite and amino-acid repression on top. The use of a pathway intermediate (urocanate) rather than the substrate (histidine) as the true inducer is a recurring theme.

5. Evolutionary and Cell-Biological Variation

Across evolutionary lineages

- **Bacteria** typically use the two-step HutF/HutG formate route and regulate the operon via HutC (urocanate-responsive) or HutP.
- **Protozoa** (e.g., *Trypanosoma cruzi*) use the one-step arginase-superfamily formiminoglutamase.
- **Vertebrates** use the folate-coupled FTCD route, coupling the pathway to one-carbon metabolism and folate homeostasis.

The obligatory core (HAL → urocanase → HutI/AMDHD1) is conserved across all these lineages, but the terminal step has been independently solved at least three times.

Cell-type and tissue variation in mammals (Finding F006)

HAL (histidase, EC 4.3.1.3) is a cytosolic enzyme that catalyzes the non-oxidative deamination of histidine to urocanic acid [PMID: 8530107], expressed in liver and epidermis. In liver it drives net histidine catabolism; in the **epidermis**, HAL-derived urocanic acid accumulates in the stratum corneum as a UVB-absorbing chromophore. Histidinemic (HAL-mutant) mice have reduced stratum-corneum UCA, greater UVB-induced cyclobutane-pyrimidine-dimer accumulation and apoptosis, reversible by topical UCA [PMID: 20686493]. In humans, reduced HAL activity causes **histidinemia**, historically the most frequent inborn metabolic error in Japan [PMID: 8530107]. The same enzyme thus has a catabolic role in one tissue and a photoprotective/structural role in another.

FTCD moonlighting and compartmentalization (Finding F011)

Vertebrate FTCD is a liver-enriched, cytosol/Golgi-localized activity that **moonlights** as the "58K" microtubule-binding Golgi protein (rat) and the chicken p60 that mediates trans-Golgi-network vesicle–microtubule interactions. Sequencing showed that 58K is a formiminotransferase cyclodeaminase, the bifunctional enzyme that catalyzes two consecutive steps in the modification of tetrahydrofolate [PMID: 9677387;

].

P 9677386

FTCD binds microtubules via post-translationally added polyglutamate side chains and localizes to the centrosome, more abundantly around the mother centriole, throughout the cell cycle and independent of Golgi integrity [PMID: 16534631]. This couples a folate-metabolic enzyme to cytoskeletal cell biology — a clear case where an enzyme's compartmentalized second function must be distinguished from its catabolic role.

Anaerobic diversion to imidazole propionate (Findings F010, F012)

In anaerobic and gut bacteria, urocanate can be diverted away from the canonical glutamate-producing route by **urocanate reductase (UrdA)**, an FAD-dependent flavoenzyme that reduces urocanate to imidazole propionate (ImP) and enables bacteria to use urocanate as an alternative respiratory electron acceptor [PMID: 38280427]. Stopped-flow analysis of the isolated FAD domain shows a charge-transfer complex between reduced FAD and urocanate forming within ~1 ms, with flavin oxidation at ~60 s⁻¹. ImP is a microbially produced histidine metabolite clinically and mechanistically linked to **insulin resistance and type 2 diabetes**, and independently associated with hypertension, chronic kidney disease progression, heart failure, and 5-year mortality (adjusted HR 1.85, 95% CI 1.20–2.88 for the highest quartile in a

North American cohort of 2,155); elevated ImP was a significant independent predictor of 5-year mortality [PMID: 37115134]. Magnesium supplementation reduced plasma ImP by ~40% ($P = 0.02$) [PMID: 33487303]. UrdA is present in humans only through the gut microbiome, providing a mechanistic link between microbiota and host cardiometabolic health. This is a boundary process — a competing fate of urocanate — that must be distinguished from canonical catabolism.

6. Constraints, Dependencies, and Failure Modes

Obligatory ordering

The first three reactions are strictly sequential because each enzyme's substrate is the previous enzyme's product: histidine → urocanate → imidazolonepropionate → FIGLU. No shortcut through this segment is chemically available; the imidazole ring must be deaminated, then hydrated, then hydrolytically opened in that order.

Mutually exclusive terminal routes

The three terminal solutions are mutually exclusive within a given organism: each disposes of the formimino carbon by a different chemistry (deimination + deformylation; single hydrolysis to formamide; or transfer to THF). An organism generally possesses one route.

Compartment- and condition-specific competition at the urocanate node

Whether urocanate proceeds to glutamate (via urocanase) or to imidazole propionate (via UrdA) depends on redox state and enzyme complement: UrdA activity requires anaerobic/respiratory conditions in which urocanate serves as an electron acceptor. This is the principal branch that "rules out" a single deterministic path through the system — the same intermediate has two fundamentally different fates.

Human failure modes

- **Histidinemia** — loss of HAL; elevated histidine, reduced urocanic acid; generally benign but with epidermal photoprotection consequences (F006) [PMID: 8530107;

P 20686493

].

- **Glutamate formiminotransferase deficiency (formiminoglutamic aciduria, OMIM 229100)** — biallelic loss-of-function FTCD variants; formiminotransferase-cyclodeaminase (FTCD) deficiency is caused by a bifunctional enzyme encoded by FTCD [PMID: 29178637], and is the **second most common inborn error of folate metabolism** [PMID: 12815595], with elevated urinary FIGLU and variable megaloblastic anemia and developmental delay. Newborn-screening data increasingly show a benign biochemical phenotype in many cases.

Therapeutic dependency via folate drain (Finding F013)

Because the FTCD transferase step consumes tetrahydrofolate (converting it to 5-formimino-THF/methenyl-THF), **high histidine-catabolic flux depletes reduced folate**. Histidine catabolism was identified as a major determinant of methotrexate sensitivity; methotrexate inhibits dihydrofolate reductase, anchoring the mechanistic link to folate metabolism [PMID: 29995852]. Independently, driving histidine metabolism with Liposome-His depleted the tetrahydrofolate (THF) pool, thereby impairing nucleotide synthesis, and methotrexate and folate depletion further enhanced its in vitro efficacy against drug-resistant cancer cells [PMID: 42287952]. This makes the folate-coupled branch a therapeutically exploitable metabolic dependency.

7. Controversies and Open Questions

1. **Is FTCD "part of" histidine catabolism or folate metabolism?** The literature splits along disciplinary lines. Because FTCD both terminates histidine degradation and feeds one-carbon metabolism, its classification is genuinely dual. Reviews that treat it purely as a folate enzyme obscure its catabolic role, and vice versa.
2. **Mixing organisms and assays.** Much mechanistic detail comes from bacteria (*P. putida*, *B. subtilis*), whereas the clinical and cell-biological detail comes from vertebrates. Extrapolating regulatory logic (e.g., HutC/urocanate induction) or terminal-route identity across these lineages risks overgeneralization. The three terminal routes are organism-specific and should not be assumed universal.

3. **Physiological significance of FTCD moonlighting.** Whether the microtubule/Golgi/centrosome localization of FTCD reflects a genuine second physiological function or an incidental binding property remains incompletely resolved; the evidence is strong that it binds and co-localizes, but causal roles in vesicle transport in vivo are less firmly established.
4. **Genotype–phenotype variability in FTCD deficiency.** Newborn-screening cohorts reveal many biochemically positive but clinically asymptomatic individuals, challenging the classical association with megaloblastic anemia and developmental delay and raising questions about penetrance and modifiers.
5. **Directionality and regulation of the UrdA branch in the human gut.** The quantitative determinants of when microbes route histidine/urocanate to ImP versus to glutamate — and how host diet (e.g., magnesium) modulates this — are active questions with cardiometabolic implications.
6. **Best representative for the ancestral role of expanded families.** For MIO-dependent lyases, HAL versus PAL/TAL; for the amidohydrolase superfamily, HutI among many members; for the arginase superfamily, formiminoglutamase versus arginase proper. Choosing the right family member to infer ancestral chemistry is non-trivial and affects evolutionary reconstructions.

8. Key References

Core pathway organization and enzymology

- *Cloning and sequencing of a 29 kb region of the Bacillus subtilis genome containing the hut and wapA loci.* [PMID: 7704263](#) — supports F001 (hutU=urocanase, hutI=imidazolonepropionase).
- *Organization and multiple regulation of histidine utilization genes in Pseudomonas putida.* [PMID: 2842309](#) — supports F001, F004, F007 (gene order; hutF=formiminoglutamate hydrolase; urocanate induction).
- *Cloning and expression in Escherichia coli of histidine utilization genes from Pseudomonas putida.* [PMID: 2858467](#) — hut gene organization.

Histidine ammonia-lyase / MIO family

- *Crystal structure of phenylalanine ammonia lyase.* [PMID: 15350127](#) — supports F002 (PAL/ HAL homology).
- *The structure of L-tyrosine 2,3-aminomutase.* [PMID: 17516659](#) — supports F002 (MIO cofactor dependence).
- *Molecular cloning and structural characterization of the human histidase gene (HAL).* [PMID: 8530107](#) — supports F006 (HAL enzymology, histidinemia).
- *Increased sensitivity of histidinemic mice to UVB radiation.* [PMID: 20686493](#) — supports F006 (epidermal photoprotection).

Urocanase

- *Structure and action of urocanase.* [PMID: 15313616](#) — supports F003 (homodimer, NAD⁺).
- *Mechanism of urocanase as studied by deuterium isotope effects.* [PMID: 6110440](#) — supports F003 (non-redox NAD⁺).
- *Proposal for the mechanism of action of urocanase.* [PMID: 2885981](#) — supports F003 (NAD⁺ electrophile).

Imidazolonepropionase

- *A common catalytic mechanism for proteins of the HutI family.* [PMID: 18442260](#) — supports F008 (amidohydrolase superfamily, Fe center).

Terminal routes

- *Histidine dissimilation in Streptomyces coelicolor.* [PMID: 6129283](#) — supports F004 (N-formylglutamate route).
- *Formiminoglutamase from Trypanosoma cruzi is an arginase-like manganese metalloenzyme.* [PMID: 24261485](#) — supports F009 (one-step arginase-superfamily route).
- *Dissociation of the octameric bifunctional enzyme formiminotransferase-cyclodeaminase.* [PMID: 3593701](#) — supports F005 (octameric channeling).

FTCD deficiency and moonlighting

- *The molecular basis of glutamate formiminotransferase deficiency.* [PMID: 12815595](#) — supports F005 (inborn error of folate metabolism).
- *Allelic spectrum of formiminotransferase-cyclodeaminase gene variants.* [PMID: 29178637](#) — supports F005 (bifunctional FTCD, formiminoglutamic aciduria).

- *58K, a microtubule-binding Golgi protein, is a formiminotransferase cyclodeaminase.* [PMID: 9677387](#) — supports F011.
- *A formiminotransferase cyclodeaminase isoform localized to the Golgi complex.* [PMID: 9677386](#) — supports F011.
- *Localization of Golgi 58K protein (FTCD) to the centrosome.* [PMID: 16534631](#) — supports F011.

Regulation

- *Role of a local transcription factor in governing cellular carbon/nitrogen homeostasis in *Pseudomonas fluorescens*.* [PMID: 33675669](#) — supports F007 (HutC).

Boundary processes — imidazole propionate and UrdA

- *Microbially Produced Imidazole Propionate Is Associated With Heart Failure and Mortality.* [PMID: 37115134](#) — supports F010.
- *Magnesium and imidazole propionate.* [PMID: 33487303](#) — supports F010.
- *Rapid reaction studies on the chemistry of flavin oxidation in urocanate reductase.* [PMID: 38280427](#) — supports F012 (UrdA).

Folate drain and therapeutics

- *Histidine catabolism is a major determinant of methotrexate sensitivity.* [PMID: 29995852](#) — supports F013.
- *Liposomes containing histidine overcome PARP inhibitor resistance.* [PMID: 42287952](#) — supports F013 (THF pool depletion).

Evolutionary context

- *Evolutionary convergence in the biosyntheses of the imidazole moieties of histidine and purines.* [PMID: 29698445](#) — context for molecular convergence and MIO autocatalysis.

9. Limitations and Knowledge Gaps

- **Organism mixing.** The mechanistic model synthesizes data from bacteria, a protozoan, and vertebrates. The obligatory core is genuinely conserved, but terminal-route assignments, regulation, and moonlighting functions are lineage-specific and should not be extrapolated wholesale.

- **Structural coverage is uneven.** High-resolution structural/mechanistic detail is excellent for HAL, urocanase, HutI, and formiminoglutamase, but the HutF/HutG two-step route is characterized more genetically than structurally, and the exact catalytic residues of HutF/HutG deserve firmer structural grounding.
- **Quantitative flux partitioning at the urocanate node** (glutamate vs. imidazole propionate) in complex communities such as the human gut is not resolved and depends on redox, diet, and microbial composition.
- **Penetrance of FTCD deficiency** is being revised by newborn screening; classical phenotype associations may be overstated.
- This review is a literature synthesis; no primary experimental data were generated. Citation snippets were verified against stored abstracts, but the breadth of the field means some lineage-specific variants (e.g., archaeal and fungal terminal routes) are under-sampled here.

10. Proposed Follow-up Experiments / Actions

1. **Comparative structural biology of HutF/HutG.** Solve or model structures of the two-step terminal enzymes to place them definitively within the amidohydrolase (or other) superfamily and identify catalytic residues, closing the structural gap relative to the other terminal routes.
2. **Phylogenomic census of terminal routes.** Systematically map which of the three terminal solutions occurs in each major lineage (including archaea and fungi) to test the "convergent, mutually exclusive" model and identify any organisms carrying more than one route.
3. **Quantitative flux assays at the urocanate branch point.** Use isotope tracing in defined gut-microbial consortia under varying redox and magnesium conditions to quantify partitioning between urocanase ($\rightarrow$ glutamate) and UrdA ($\rightarrow$ imidazole propionate).
4. **Test FTCD moonlighting causality.** Use separation-of-function FTCD mutants (catalytic-dead vs. microtubule-binding-dead) to determine whether the Golgi/centrosome role is physiologically required, independent of catabolic activity.
5. **Exploit the folate-drain dependency therapeutically.** Extend the histidine-catabolism/antifolate synergy (Liposome-His + methotrexate) into in vivo models and additional tumor types with defined FTCD/AMDHD1 expression, stratifying by pathway flux.

6. **Clarify FTCD-deficiency penetrance.** Correlate genotype, residual FTCD activity, and clinical phenotype in newborn-screening cohorts to refine counseling and the classical anemia/developmental-delay association.
-

Prepared as a commissioned review synthesis. Findings F001–F013 and 41 reviewed papers underpin the claims above; citation snippets were validated against stored abstracts. Uncertainty and organism-specific caveats are flagged throughout to avoid overgeneralization from any single system.

Generated by OpenScientist — Scientific Hypothesis Agent for Novel Discovery