

Glutathione Biosynthesis: A Two-Reaction Module Conserved by Function, Not by Ancestry

*A commissioned review-style synthesis for a molecular biology audience Search date: 2026-07-25.
Sources: PubMed primary literature and authoritative reviews.*

1. Executive Summary

Glutathione (GSH; γ -L-glutamyl-L-cysteinyl-glycine) is the most abundant low-molecular-weight thiol in most eukaryotes and in the cyanobacterial and proteobacterial lineages of prokaryotes. Its **de novo biosynthesis is a strikingly simple, two-reaction module**: (1) glutamate–cysteine ligase (γ -ECL; GshA in bacteria, GCLC in animals) condenses L-glutamate and L-cysteine into the unusual γ -peptide-linked dipeptide **γ -L-glutamyl-L-cysteine (γ -EC)**, and (2) glutathione synthetase (GS; GshB / GSS) adds glycine to γ -EC to give glutathione. Both steps consume ATP. The pathway is defined at the level of these two catalytic *roles* — not by a fixed gene arrangement, subunit composition, or regulatory logic — and the module reappears across life in several physical implementations that perform the same chemistry.

The central mechanistic facts are well established and mutually reinforcing across structural, kinetic, genetic, and clinical evidence. The **first step is rate-limiting, committed, and feedback-inhibited by the end product GSH**; structural work in yeast shows glutathione physically occupying the substrate sites and disrupting Mg^{2+} /ATP coordination. The **order of the two reactions is obligatory** because the second enzyme requires the γ -EC intermediate produced by the first — a dependency dramatically confirmed by human inborn errors of metabolism, where blocking the *second* enzyme (GSS) diverts accumulating γ -EC into 5-oxoproline (5-oxoprolinuria), while blocking the *first* enzyme (GCLC) causes hemolytic anemia *without* 5-oxoprolinuria. The two enzymes belong to two **unrelated ancient ligase superfamilies** — the glutamine-synthetase family (γ -ECL) and the ATP-grasp fold family (GS) — so the module's conservation reflects convergent recruitment of two catalytic activities rather than descent from a single ancestral operon.

Variation on this theme is biologically informative. Some Firmicutes and Pasteurellaceae fuse the two activities into a single **bifunctional GshF** enzyme that channels the intermediate; other bacteria dispense with synthesis entirely and **import** GSH; and many prokaryotes never adopted glutathione at all, using **mycothiol (Actinobacteria)** or **bacillithiol (Firmicutes)** as their protective thiol. In animals, the first enzyme is elaborated into a **GCLC–GCLM heterodimer** in which a non-catalytic modifier subunit tunes catalytic kinetics and feedback sensitivity, and expression of both subunits is governed by the NRF2 antioxidant program. In plants the enzymes are chloroplast-localized, consistent with a cyanobacterial/endosymbiotic origin. Finally, a key compartmental constraint: **mammalian mitochondria do not synthesize GSH** — they import cytosolic glutathione through the inner-membrane carrier SLC25A39. This report synthesizes eight confirmed findings from 43 papers into a coherent picture of how the pathway works, how it varies, where it came from, and what remains uncertain.

2. Definition and Biological Boundaries

2.1 What is inside the boundary

The system under review is the **committed, ATP-dependent conversion of L-glutamate + L-cysteine + glycine into glutathione via γ -glutamylcysteine**, comprising exactly two catalytic roles:

Step	Enzyme (roles)	Reaction	Product
1	Glutamate–cysteine ligase (γ -ECL / GshA / GCLC; EC 6.3.2.2)	$\text{L-Glu} + \text{L-Cys} + \text{ATP} \rightarrow \gamma\text{-Glu-Cys} + \text{ADP} + \text{P}_i$	γ -glutamylcysteine (γ -EC)
2	Glutathione synthetase (GS / GshB / GSS; EC 6.3.2.3)	$\gamma\text{-Glu-Cys} + \text{Gly} + \text{ATP} \rightarrow \text{GSH} + \text{ADP} + \text{P}_i$	glutathione (GSH)

The defining biochemical feature is the **γ -carboxamide (isopeptide) bond** formed in step 1 between the γ -carboxyl of glutamate and the amino group of cysteine. This non-canonical linkage is what makes GSH resistant to ordinary peptidases and is the structural signature of the pathway.

2.2 What is explicitly outside the boundary

The following processes are frequently discussed alongside glutathione but are **not** part of the biosynthetic module and should be treated as neighboring systems:

- **Glutathione reduction** ($\text{GSSG} \rightarrow 2 \text{GSH}$ by glutathione reductase) — maintains the redox pool but does not make GSH de novo.
- **Conjugation and detoxification** (glutathione S-transferases; microsomal MGST enzymes) — consumes GSH.
- **Peroxide detoxification** (glutathione peroxidases, GPX4; ferroptosis defense) — consumes GSH.
- **Degradation and the wider γ -glutamyl cycle** (γ -glutamyl transpeptidase, 5-oxoprolinase, γ -glutamyl cyclotransferase) — recovers amino acids; its involvement in disease (5-oxoproline accumulation) is a *downstream consequence* of a synthesis block, not part of synthesis itself.
- **Cysteine supply / transsulfuration and cystine import** (SLC7A11/xCT) — feeds the pathway but is upstream metabolism.
- **NRF2/KEAP1 transcriptional control** — an important regulatory layer in animals (see §5) but a *regulatory input*, not a catalytic step.


A large fraction of the literature retrieved in this investigation (ferroptosis, NRF2 pharmacology, GST/GPX enzymology) concerns these adjacent systems and must be read as context rather than as evidence about the two ligase reactions themselves.

2.3 Competing definitions

The main definitional subtlety is whether the "pathway" is a set of *genes* or a set of *reactions*. The evidence strongly favors a **reaction-role definition**: the same two-step chemistry is executed by (i) two separate monofunctional enzymes (*E. coli*, most organisms), (ii) a single bifunctional fusion protein (GshF in *Streptococcus agalactiae*, *Listeria*, *Pasteurella multocida*), or (iii) not at all, with GSH obtained by import. Any gene-centric definition fails to capture these equivalent implementations.

3. Mechanistic Overview

3.1 The best current model of the sequence of events



Step 1 (obligatory, rate-limiting, controlling). γ -ECL activates the γ -carboxyl of glutamate as an acyl-phosphate using ATP, then condenses it with the α -amino group of cysteine to form the γ -isopeptide bond. This is the **committed and flux-controlling step** of the entire pathway (Finding F001). It is directly, non-allosterically **feedback-inhibited by GSH**: crystal structures of *Saccharomyces cerevisiae* GCL show the inhibitor occupying both the glutamate- and the presumed cysteine-binding sites and disrupting the Mg^{2+} coordination in the ATP site ([PMID: 20220146](#)). The mechanism-based inhibitor L-buthionine-S-sulfoximine (BSO), widely used to deplete cellular GSH, is phosphorylated on its sulfoximine nitrogen at the same catalytic site.

Step 2 (obligatory, downstream). GS ligates glycine to the cysteinyl carboxyl of γ -EC, again through an ATP-dependent acyl-phosphate intermediate, using the two-domain **ATP-grasp fold**. This step is generally not rate-limiting and is not the primary point of feedback control.

Ordering constraint. The two reactions are strictly sequential: GS cannot act until γ -ECL has supplied γ -EC. This is not merely a kinetic preference — it is enforced by substrate specificity and is visible in human disease (§6).

3.2 Obligatory vs. conditional vs. accessory elements

- **Obligatory:** both ligation steps; ATP and Mg^{2+} ; the γ -EC intermediate; the fixed reaction order.

- **Conditional:** the GCLM modifier subunit (present in animals, absent in many bacteria and lower eukaryotes; modulates but is not required for catalysis); NRF2-driven transcriptional induction (animal-specific stress response); subcellular localization (chloroplast in plants).
 - **Accessory / alternative:** the bifunctional GshF architecture (channels the intermediate); glutathione import systems (bypass synthesis); transcriptional regulators such as LysR-family LsrB/OxyR in rhizobia.
-

4. Major Molecular Players and Active Assemblies

4.1 Two enzymes from two unrelated superfamilies (Findings F001, F007)

The single most important structural fact about this pathway is that **its two enzymes are not evolutionarily related to each other**. γ -ECL is a member of the **glutamine-synthetase family**, whereas glutathione synthetase adopts the **ATP-grasp fold** shared by "highly diverse carboxyl-to-amine/thiol ligases" ([PMID: 16339152](#)). The module is therefore conserved by *reaction role* — two independently recruited ATP-dependent amide-bond-forming activities — rather than by shared ancestry. Consistent with this, the ATP-grasp step sits within a broad ligase clade that includes cyanophycin synthetase (CphA1), whose domains are explicitly "homologous to glutathione synthetases and muramyl ligases" ([PMID: 34385683](#)). Because both belong to large, ancient ligase superfamilies, the module's near-universal presence in GSH-using organisms is a case of **conservation by function rather than by common descent**.

4.2 The controlling first enzyme (Finding F002)

γ -ECL is where flux is set. In plants, an elegant genetic demonstration comes from *Arabidopsis*: the *rax1-1* allele of **GSH1**, which encodes chloroplastic γ -glutamylcysteine synthetase, "the controlling step of glutathione biosynthesis," lowers foliar GSH (by $\geq 50\%$) and reprograms stress-defense gene expression ([PMID: 15308753](#)). This simultaneously locates the enzyme (chloroplast) and confirms its control role in a whole organism. The structural basis of its control is direct product feedback: yeast GCL structures show GSH occupying both substrate sites and disrupting Mg^{2+} /ATP coordination ([PMID: 20220146](#)).

4.3 The animal heterodimer: GCLC + GCLM (Finding F006)

In mammals, γ -ECL is a **heterodimer** of a catalytic subunit (**GCLC**) and a modifier subunit (**GCLM**). GCLM is not itself catalytic; it "change[s] the catalytic characteristics of the holoenzyme." In *Gclm*^{-/-} mice, residual GCLC shows an ≈ 2 -fold increase in K_m for glutamate and "dramatically enhanced sensitivity to GSH inhibition"; whole-animal GSH falls to 9–16% of wild-type, yet the mice remain viable and fertile (PMID: 12384496). Human enzymology reinforces the modifier role: clinical GCLC missense mutants have reduced activity that is partially rescued by co-expressing GCLM (PMID: 21657237). This heterodimeric elaboration is a lineage-specific feature absent from the monomeric single-subunit γ -ECLs of many bacteria and lower eukaryotes.

4.4 The bifunctional GshF fusion (Finding F003)

Some bacteria replace the two separate enzymes with a single **bifunctional GshF** protein carrying an N-terminal γ -proteobacterial γ -ECL-like domain fused to an ATP-grasp domain most similar to cyanophycin synthetases rather than to canonical GS. This architecture occurs in Firmicutes (*Streptococcus agalactiae*, *Listeria monocytogenes*) and Pasteurellaceae (*Pasteurella multocida*) (PMID: 16339152). Because the fused active sites are kinetically coupled, GshF produces GSH **without releasing free γ -EC** — a substrate-channeling behavior exploited biotechnologically: expressing bacterial GshF in *Pichia pastoris* enables glutathione production "without accumulation of the intermediate" (PMID: 22610434).

4.5 Transporters and compartmental players (Finding F008)

Mammalian mitochondria lack the biosynthetic enzymes and instead **import cytosolic GSH** via the inner-membrane carrier **SLC25A39**; loss of SLC25A39 depletes the mitochondrial GSH pool and impairs OXPHOS and iron homeostasis (PMID: 34707288; PMID: 35513392). This defines a hard compartmental boundary: within a single cell, one compartment can be biosynthetically self-sufficient (cytosol) while another depends entirely on transport (mitochondrion).

5. Evolutionary and Cell-Biological Variation

5.1 Distribution across lineages (Finding F004)

GSH is the dominant protective thiol in nearly all eukaryotes and in cyanobacteria and proteobacteria, but its **biosynthetic pathway has multiple origins and a patchy prokaryotic distribution**. Many bacteria that lack GSH biosynthesis use alternative thiols instead: **mycothiol** in Actinobacteria and **bacillithiol** in Firmicutes. Extensive thiol diversity "existed among prokaryotes prior to evolution of the cyanobacteria and the development of an oxidizing atmosphere" (PMID: 23075826). The eukaryotic acquisition of GSH biosynthesis is consistent with a cyanobacterial/endosymbiotic route, in line with the chloroplast localization of the plant enzymes (PMID: 15308753).

5.2 Alternative routes to the same outcome

Strategy	Representative organisms	How γ -EC is handled
Two separate enzymes (GshA + GshB)	<i>E. coli</i> , most bacteria, plants, fungi, animals	γ -EC released as a free intermediate
Bifunctional fusion (GshF)	<i>S. agalactiae</i> , <i>L. monocytogenes</i> , <i>P. multocida</i>	γ -EC channeled internally; not released
Import only	<i>Haemophilus influenzae</i> , some enterococci/streptococci; mammalian mitochondria (SLC25A39)	No synthesis; GSH acquired from environment/cytosol
No glutathione	Actinobacteria (mycothiol), many Firmicutes (bacillithiol)	Uses unrelated protective thiols

5.3 Regulatory variation

Control logic differs across lineages. In animals, both GCLC and GCLM are transcriptional targets of the **NRF2/KEAP1** antioxidant program, and much recent literature (ferroptosis defense, xenobiotic response) concerns this induction — although this is a regulatory input rather than a catalytic feature of the pathway. In α -proteobacteria such as *Sinorhizobium meliloti*, the LysR-family regulator **LsrB**, together with **OxyR**, positively regulates transcription

of *gshA* and *gshB* under free-living and symbiotic conditions (PMID: 23883684). Across all systems, however, the conserved post-translational control point is the same: **GSH feedback inhibition of the first enzyme**.

6. Constraints, Dependencies, and Failure Modes

6.1 Ordering and substrate constraints

The pathway imposes a strict order: **step 1 must precede step 2** because GS is specific for the γ -EC intermediate. This constraint is not theoretical — it is written into human genetics.

6.2 Human inborn errors define the constraint experimentally (Findings F005, F008)

The two enzyme deficiencies produce **biochemically distinct** phenotypes that pinpoint where each enzyme sits in the pathway:

Feature	First-enzyme deficiency (GCLC / γ -ECL)	Second-enzyme deficiency (GSS)
Primary phenotype	Neonatal hemolytic anemia; low erythrocyte GSH; occasional late-onset spinocerebellar ataxia	Spectrum from isolated hemolytic anemia to chronic metabolic acidosis and severe neonatal-lethal neurological disease
5-oxoprolinuria	Absent (block is upstream of γ -EC)	Present (accumulated γ -EC diverted to 5-oxoproline)
Diagnostic hallmark	Low GSH, hemolysis, genetic confirmation	Elevated urinary/amniotic 5-oxoproline (pyroglutamic acid)
Key references	PMID: 28571779, PMID: 39359943	PMID: 39221916, PMID: 34791353

When GSS is defective, accumulated γ -EC is diverted through γ -glutamyl cyclotransferase to **5-oxoproline (pyroglutamic acid)**, which builds up in body fluids and is diagnostic; clinical severity ranges "from isolated hemolytic anemia, sometimes associated with chronic metabolic acidosis and 5-oxoprolinuria, to severe neurological phenotypes with neonatal lethality" (PMID: 39221916; PMID: 34791353). Crucially, first-enzyme (GCLC) deficiency does **not** produce

5-oxoprolinuria because the block lies upstream of γ -EC (PMID: 28571779; PMID: 39359943). This asymmetry is the clearest possible in vivo confirmation of the reaction order and of the identity of the branch-point metabolite. GCLC deficiency is exceedingly rare (only ~9–12 patients from a handful of families reported worldwide; PMID: 40277844).

6.3 Compartmental failure mode

Because mitochondria depend on SLC25A39-mediated import rather than local synthesis, loss of that transporter depletes the mitochondrial GSH pool even when cytosolic biosynthesis is intact, impairing OXPHOS and iron–sulfur homeostasis (PMID: 34707288; PMID: 35513392). This is a "same-cell, different-compartment" failure mode that biosynthesis alone cannot rescue.

6.4 Substrate-analog constraint

The pathway's substrate specificity is loose enough that replacing cysteine with L-2-aminobutyrate yields **ophthalmic acid** (γ -Glu-2-aminobutyl-Gly), a natural GSH analog that can be produced in engineered *E. coli* (PMID: 29352090) — illustrating that the enzymes constrain the γ -glutamyl and glycyl positions more tightly than the central residue.

7. Controversies and Open Questions

- 1. Depth and number of evolutionary origins.** The pathway clearly has *multiple* origins and a patchy prokaryotic distribution (PMID: 23075826), but the precise phylogenetic history — how many independent recruitments of the ATP-grasp GS activity occurred, and whether GshF fusions are ancestral or derived — remains unresolved. The close similarity of the GshF ATP-grasp domain to cyanophycin synthetases rather than to canonical GS hints at independent evolutionary sampling of the same fold, but this is inferred from sequence/structure relationships rather than experimentally reconstructed ancestors.
- 2. Best representative for the ancestral GS role.** Because glutathione synthetase belongs to a large, diverse ATP-grasp clade that also contains cyanophycin synthetase and muramyl ligases (PMID: 34385683), identifying which family member best represents the ancestral glutathione-forming activity is genuinely open. Canonical prokaryotic GshB and the fused GshF domains may not descend from a single ancestral GS.

3. **Function of the GCLM modifier subunit in vivo.** GCLM clearly tunes kinetics and feedback sensitivity in vitro and in knockout mice ([PMID: 12384496](#); [PMID: 21657237](#)), yet *Gclm*^{-/-} mice survive with ~10% of normal GSH. The physiological conditions under which the modifier becomes essential (particular stresses, tissues, or developmental windows) are not fully mapped.
4. **Organism-mixing in the literature.** Much of what is "known" combines data across yeast structures, mouse genetics, human clinical cases, and bacterial biochemistry. Feedback-inhibition structures come from yeast; heterodimer kinetics from mouse; disease phenotypes from humans; fusion architecture from Firmicutes. These are broadly consistent, but caution is warranted before assuming, e.g., that mammalian feedback geometry is identical to the yeast structure or that all bacterial γ -ECLs share the mammalian regulatory behavior.
5. **Mitochondrial import mechanism.** SLC25A39 is established as necessary for mitochondrial GSH import ([PMID: 34707288](#)), but the transport mechanism, stoichiometry, and whether additional carriers contribute in specific tissues remain active questions.

Note on citation integrity: two citation snippets recorded during the investigation (for

[P 34707288](#)

and

[P 21657237](#)

were flagged as paraphrases/title-derived rather than exact abstract quotes. The underlying claims they support — SLC25A39-mediated mitochondrial GSH import, and GCLM rescue of GCLC mutants — are nonetheless corroborated by the companion primary sources cited above ([PMID: 35513392](#); [PMID: 12384496](#)).

8. Mechanistic Model — Synthesis

Bringing the eight findings together, the pathway is best understood as a **functionally conserved, structurally modular assembly of two independent ligase activities**, whose behavior is governed by three organizing principles:

1. **Committed control at the front door.** Flux is set almost entirely at step 1 by feedback inhibition of γ -ECL by the end product GSH (F002), a control point conserved from yeast structures to plant genetics to mammalian kinetics.

2. **An obligatory hand-off through γ -EC.** The intermediate γ -glutamylcysteine is the pivot of the whole system. Its fate — released and passed to a separate GS, channeled internally by a GshF fusion, or (when GS fails) shunted to 5-oxoproline — distinguishes all the pathway's variants and failure modes (F003, F005, F008).
 3. **Conservation by role, elaboration by lineage.** The two catalytic roles are ancient and near-universal among GSH-using organisms, but their physical implementation is plastic: monofunctional vs. fused enzymes, monomeric vs. heterodimeric first enzymes, cytosolic synthesis vs. mitochondrial import, and NRF2 vs. LysR-family regulation are all lineage-specific overlays on the same invariant chemistry (F001, F004, F006, F007).
-

9. Evidence Base

PMID	Contribution	Role in this review
16339152	Characterization of bifunctional GshF of <i>P. multocida</i> ; assigns γ -ECL to glutamine-synthetase family and GS to ATP-grasp fold	Supports F001, F003, F007 (two unrelated families; fusion architecture)
20220146	Crystal structures of yeast GCL with feedback/ pharmacological inhibitors	Supports F001, F002 (rate-limiting, feedback-inhibited first step)
15308753	<i>Arabidopsis rax1-1</i> GSH1 allele links biosynthesis to stress-defense gene expression	Supports F002, F004 (controlling step; chloroplast localization)
22610434	GshF expressed in <i>Pichia</i> produces GSH without intermediate accumulation	Supports F003 (intermediate channeling by fusion)
23075826	Review of glutathione, mycothiol, bacillithiol distribution and origins	Supports F004 (multiple origins; alternative thiols)
39221916	GSS-deficient fetuses; elevated 5-oxoproline; clinical severity spectrum	Supports F005 (second-enzyme block $\rightarrow$ 5-oxoprolinuria)
34791353	GC-MS diagnosis of GSS deficiency via urinary 5-oxoproline	Supports F005 (diagnostic hallmark)
12384496	<i>Gclm</i> ^{-/-} mouse; kinetics and feedback sensitivity	Supports F006 (modifier subunit role)
21657237	GCLC mutant rescue by GCLM association	Supports F006, F008 (modifier rescues catalytic subunit)
34385683	Cyanophycin synthetase structure; homology to GS and muramyl ligases	Supports F007 (ATP-grasp ligase clade)
34707288	SLC25A39 necessary for mitochondrial GSH import	Supports F008 (compartmental constraint)
35513392	CRISPR screen identifies SLC25A39 in mitochondrial GSH transport, linking iron and OXPHOS	Supports F008 (import couples GSH to OXPHOS)
28571779	Six children with GCL deficiency and hemolytic anemia	Supports F008 (first-enzyme phenotype; no 5-oxoprolinuria)
39359943		

PMID	Contribution	Role in this review
	Late-onset spinocerebellar ataxia + hemolysis from GCLC variant	Supports F008 (first-enzyme phenotype spectrum)
40277844	Rare GCLC deficiency case; ~9 patients worldwide	Context (rarity of first-enzyme deficiency)
23883684	LsrB/OxyR regulation of <i>gshA/gshB</i> in <i>S. meliloti</i>	Context (bacterial transcriptional regulation)
29352090	Ophthalmic acid production in engineered <i>E. coli</i>	Context (substrate flexibility)

10. Limitations and Knowledge Gaps

- **No primary experimental data were generated;** this is a literature synthesis. Conclusions inherit the biases of the retrieved corpus, which is heavily weighted toward downstream/adjacent topics (ferroptosis, NRF2 pharmacology) rather than the ligase reactions themselves.
- **Organism heterogeneity.** The mechanistic model stitches together yeast structural biology, mouse genetics, human clinical genetics, and bacterial biochemistry. Cross-organism extrapolation is reasonable but not proven at every node.
- **Sparse structural coverage of the mammalian holoenzyme.** Feedback-inhibition geometry is best characterized in yeast; a high-resolution mammalian GCLC–GCLM feedback-bound structure would sharpen the model.
- **Phylogenetics not reconstructed.** Claims about "multiple origins" and the ancestral GS representative rest on published reviews and structural homology, not on an ancestral-sequence reconstruction performed here.
- **Regulation outside animals under-sampled.** Beyond NRF2 (animals) and LsrB/OxyR (rhizobia), the regulatory logic in fungi, protists, and diverse bacteria is not systematically covered.

11. Proposed Follow-up Experiments / Actions

1. **Ancestral-sequence reconstruction of the ATP-grasp GS role** across GshB, GshF, cyanophycin synthetase, and muramyl ligases to test whether glutathione-forming activity arose once or multiple times, and to identify the best extant representative of the ancestral enzyme.
 2. **Comparative structural determination** of the mammalian GCLC–GCLM heterodimer in the GSH-feedback-bound state, to test whether the yeast feedback geometry ([PMID: 20220146](#)) generalizes.
 3. **Systematic metabolite profiling** (γ -EC vs. 5-oxoproline) across cell models of first- vs. second-enzyme deficiency to quantitatively map the branch-point flux that distinguishes the two disorders.
 4. **Mechanistic dissection of SLC25A39** transport (stoichiometry, driving force, tissue-specific redundancy) to close the compartmental gap in the model.
 5. **Kinetic characterization of GshF channeling** to measure the efficiency of intermediate transfer relative to two free enzymes, clarifying the selective advantage of the fusion architecture.
 6. **Broader phylogenetic survey** of GSH biosynthesis vs. mycothiol/bacillithiol use, to firm up the "cyanobacteria + rising oxygen" origin narrative with quantitative distribution data.
-

12. Key References

1. Vergauwen B. et al. *Characterization of the bifunctional γ -glutamate-cysteine ligase/ glutathione synthetase (GshF) of *Pasteurella multocida**. [PMID: 16339152](#)
2. Biterova E.I. & Barycki J.J. *Structural basis for feedback and pharmacological inhibition of *S. cerevisiae* glutamate cysteine ligase*. [PMID: 20220146](#)
3. Ball L. et al. *Evidence for a direct link between glutathione biosynthesis and stress defense gene expression in *Arabidopsis**. [PMID: 15308753](#)
4. Ge Y. et al. *Expression of bacterial GshF in *Pichia pastoris* for glutathione production*. [PMID: 22610434](#)
5. Fahey R.C. *Glutathione analogs in prokaryotes*. [PMID: 23075826](#)
6. Jury et al. *Multiple congenital anomalies in two fetuses with glutathione-synthetase deficit*. [PMID: 39221916](#)

7. Kaur et al. *GC-MS-aided diagnosis of glutathione synthetase deficiency*. PMID: 34791353
8. Yang Y. et al. *Initial characterization of the Gclm^{-/-} knockout mouse*. PMID: 12384496
9. Willis et al. *Enzymatic defects underlying hereditary glutamate cysteine ligase deficiency mitigated by subunit association*. PMID: 21657237
10. Sharon I. et al. *Structures and function of the amino acid polymerase cyanophycin synthetase*. PMID: 34385683
11. Wang Y. et al. *SLC25A39 is necessary for mitochondrial glutathione import in mammalian cells*. PMID: 34707288
12. Shi X. et al. *Combinatorial GxGxE CRISPR screen identifies SLC25A39 in mitochondrial glutathione transport*. PMID: 35513392
13. Almusafri et al. *Clinical and molecular characterization of 6 children with GCL deficiency causing hemolytic anemia*. PMID: 28571779
14. Al-Hatou et al. *GCL deficiency with late-onset spinocerebellar ataxia and hemolytic anemia*. PMID: 39359943
15. *Hemolytic anemia due to γ -glutamylcysteine synthetase deficiency: a rare novel case*. PMID: 40277844
16. Tang et al. *The Sinorhizobium meliloti LysR family regulator LsrB in glutathione biosynthesis*. PMID: 23883684
17. *Production of ophthalmic acid using engineered Escherichia coli*. PMID: 29352090

Prepared as a commissioned review synthesis. All mechanistic claims are anchored to the cited primary literature; uncertainty is flagged explicitly in §7 and §10. Evidence spanning multiple organisms is identified as such to avoid overgeneralization from any single system.
