

Homogentisate Catabolism: A Mechanistic Review

A concise, review-style synthesis of the three-reaction "lower" pathway that opens homogentisate and delivers its carbon to central metabolism.

1. Executive Summary

Homogentisate (2,5-dihydroxyphenylacetate, HGA) catabolism is a compact, evolutionarily ancient, three-enzyme module that dismantles an aromatic ring and routes its carbon skeleton into central carbon metabolism. The module comprises, in obligate order:

1. **Homogentisate 1,2-dioxygenase (HmgA / HGD)** — an Fe(II)-dependent dioxygenase that oxidatively cleaves the aromatic ring to yield **maleylacetoacetate (MAA)**.
2. **Maleylacetoacetate isomerase (HmgC / MAAI = GSTZ1)** — a glutathione-dependent *cis*→*trans* isomerase that converts MAA to **fumarylacetoacetate (FAA)**.
3. **Fumarylacetoacetate hydrolase / fumarylacetoacetase (HmgB / FAH)** — a metal-dependent C–C bond hydrolase that splits FAA into **fumarate** (a TCA-cycle intermediate) and **acetoacetate** (a ketone-body precursor).

This module is the shared *terminal* segment of tyrosine/phenylalanine catabolism in animals and of numerous microbial and fungal aromatic-degradation routes that converge on homogentisate. Its clinical and physiological importance is underscored by two human inborn errors: **alkaptonuria** (loss of HGD) and **hereditary tyrosinemia type I / HT1** (loss of FAH). Notably, the middle step (MAAI/GSTZ1) is partially redundant with a non-enzymatic glutathione-mediated isomerization, which likely explains why no unequivocal human "MAAI deficiency" disease exists, in contrast to the flanking steps.

2. Definition and Biological Boundaries

What is included. The system, *sensu stricto*, is exactly three reactions: ring cleavage (HGA → MAA), isomerization (MAA → FAA), and hydrolysis (FAA → fumarate + acetoacetate). Its **entry metabolite** is homogentisate and its **exit metabolites** are fumarate and acetoacetate. Everything between those boundaries defines the pathway.

What feeds in but is not part of it (upstream boundary). Homogentisate is a *convergence node*. In vertebrates it is produced from 4-hydroxyphenylpyruvate by **4-hydroxyphenylpyruvate dioxygenase (HPD/HPPD)**, itself downstream of tyrosine aminotransferase acting on tyrosine (and, before that, phenylalanine hydroxylase acting on phenylalanine). In microbes and fungi, homogentisate is generated from diverse aromatic substrates — e.g., an ethylbenzene → styrene → phenylacetate → homogentisate route in the fungus *Exophiala lecanii-corni*).

(P 15731901)

These upstream reactions (HPPD in particular) are frequently conflated with the pathway but are **accessory feeder reactions**, not part of the ring-opening module itself. The clinical drug **nitisinone (NTBC)**, used in HT1 and alkaptonuria, acts *upstream* by inhibiting HPPD — it does not touch any of the three core enzymes

(P 41003839)

(P 33375092)

What should be treated separately (commonly confused). - **Gentisate (2,5-dihydroxybenzoate) 1,2-dioxygenase pathways.** Gentisate catabolism is a *parallel but distinct* microbial ring-cleavage route with a similar name and similar chemistry (dioxygenolytic cleavage to maleylpyruvate, then a GSH-dependent isomerase to fumarylpyruvate, then a hydrolase). The enzymes, substrates, and gene sets differ; do not merge the two. - **GSTZ1's xenobiotic/moonlighting biology.** Because MAAI is GSTZ1, much literature discusses its role in dichloroacetate (DCA) biotransformation, redox/ferroptosis signaling (NRF2/GPX4, HMGB1/GPX4), and cancer metabolism

(P 37742772)

(P 33931597)

(P 36905252)

).

P 35562974

These are genuine activities of the same protein but are **not the isomerization step** of homogentisate catabolism and should be kept conceptually distinct. - **Pyomelanin / ochronotic pigment formation**. When the pathway is blocked, accumulated homogentisate auto-oxidizes and polymerizes into pigment (pyomelanin in microbes; ochronotic pigment in human connective tissue). This is a *downstream consequence of pathway failure*, not a pathway reaction

(

P 19028908


).

P 31020477

Competing definitions. The main terminological variation is nomenclature: the microbial locus names **hmgA/hmgB/hmgC** versus the mammalian gene names **HGD/FAH/GSTZ1(MAAI)**. Some literature draws the pathway boundary to *include* HPPD (the "homogentisate branch of tyrosine catabolism"); others, as in this brief, restrict it to the three ring-processing reactions. Both usages are defensible provided the boundary is stated.

3. Mechanistic Overview

The current best model is a strictly linear, non-branched relay in which each product is the next enzyme's substrate:



Step 1 — Ring cleavage (obligatory). HGD is a non-heme Fe(II) dioxygenase. Computational (hybrid-DFT and QM/MM) studies converge on a mechanism with three chemical stages: formation of an Fe-bound **peroxo/alkylperoxo intermediate**, **homolytic O–O cleavage**

producing an arene-oxide radical, and finally scission of the six-membered ring to the ring-opened product

([P 16332080](#))

[P 27119315](#)

The regiochemistry (intra- vs extra-diol relative to the iron-coordinating substituents) is controlled by the relative barriers for decay of the arene-oxide radical intermediate

([P 16332080](#))

Both oxygen atoms of O₂ are incorporated, defining this as an intradiol-type dioxygenase reaction on a homogentisate substrate.

Step 2 — Isomerization (obligatory chemically, but conditionally enzyme-dependent).

MAA is produced in the *cis* (maleyl) configuration; entry into the final hydrolysis requires the *trans* (fumaryl) configuration. MAAI (=GSTZ1) uses glutathione as a catalytic cofactor to rotate about the double bond. Crucially, this *cis*→*trans* isomerization can also occur **non-enzymatically via glutathione** — a bypass demonstrated directly in vitro and inferred in vivo from MAAI-knockout mice, which remain viable but accumulate FAA and succinylacetone

([P 12052898](#))

Thus the *isomerization event* is obligatory, but the *enzyme* is accessory/facultative.

Step 3 — Hydrolysis (obligatory). FAH catalyzes an unusual **C–C bond hydrolysis**, cleaving FAA into fumarate and acetoacetate. Structural work established FAH as the founding member of a novel metalloenzyme fold (a "mixed β-sandwich roll"), employing a divalent metal ion (Ca²⁺/Mg²⁺ observed), a **Glu–His catalytic dyad activating a water nucleophile**, and a charged **oxyanion hole** that stabilizes a **tetrahedral alkoxide transition-state intermediate**

([P 10508789](#))

[P 11154690](#)

Phosphorus-based transition-state mimics bind with nanomolar affinity and phenocopy the metabolic block, validating the tetrahedral-intermediate model

([P 17064256](#))

Obligatory ordering and no shortcuts. The three reactions must proceed in order because each substrate is chemically defined by the prior step (an intact ring cannot be hydrolyzed by FAH; *cis*-MAA is not the FAH substrate). No non-enzymatic bypass is known for steps 1 or 3, which is why their loss produces frank human disease.

4. Major Molecular Players and Active Assemblies

Step	Enzyme (microbial / mammalian)	EC	Cofactor / metal	Products	Assembly & structural notes
1	Homogentisate 1,2-dioxygenase — HmgA / HGD	1.13.11.5	Non-heme Fe(II) , O ₂	Maleylacetoacetate	Human HGD: 445 aa, gene on 3q13, 14 exons; oligomeric (multi-subunit) enzyme; active site with 3 His + Glu iron ligands in the modeled chemistry (P 16332080) (P 41236909)
2	Maleylacetoacetate isomerase — HmgC / MAAI = GSTZ1	5.2.1.2	Glutathione (catalytic)	Fumarylacetoacetate	Member of the cytosolic GST superfamily (zeta class); Cys16 implicated in DCA-mediated inactivation; most abundant in liver (P 21303221) (P 37742772)

Step	Enzyme (microbial / mammalian)	EC	Cofactor / metal	Products	Assembly & structural notes
3	Fumarylacetoacetate hydrolase — HmgB / FAH	3.7.1.2	Mg²⁺/Ca²⁺ , H ₂ O	Fumarate + acetoacetate	Novel α/β "FAH-superfamily" fold; N-terminal (~120 aa) + C-terminal (~300 aa) domains; catalytically active dimer in equilibrium with an inactive, aggregation-prone monomer; Glu–His–water triad + oxyanion hole (P 10508789 P 11154690 P 31300554)

Genomic organization. In many microbes and fungi the genes are clustered and co-regulated with the feeder enzymes. In *Aspergillus fumigatus*, the L-tyrosine/L-phenylalanine degradation cluster contains **hppD**, **hmgX**, **hmgA**, **fahA**, **maiA**, **hmgR** (the last a pathway-specific regulator), tying homogentisate processing to a single inducible operon-like unit

([P 31020477](#))

P 19028908

In mammals the three genes are unlinked (HGD on 3q13; FAH on 15q; GSTZ1 on 14q) and regulated independently, but co-expressed in the same organs (liver hepatocytes and kidney proximal tubule;).

P 12052898

5. Evolutionary and Cell-Biological Variation

Deep conservation. All three activities are found across bacteria, fungi, plants, and animals, reflecting an ancient role in aromatic-ring catabolism that predates the split of these lineages. FAH in particular has **no sequence homologs among other hydrolase classes** and defines its own fold/superfamily

P 10508789

(), implying an ancient and singular evolutionary invention for C–C hydrolysis. HGD belongs to the broad non-heme Fe(II) dioxygenase repertoire that underpins microbial aromatic degradation globally.

Lineage variation. - **Microbes/fungi:** homogentisate catabolism is the **central/lower funnel** for catabolism of many aromatics (tyrosine, phenylacetate, styrene, ethylbenzene, nicotinate-related aromatics), typically clustered and substrate-inducible. Blocking step 1 diverts flux to **pyomelanin**, a stress- and virulence-associated pigment

P 19028908

P 31020477

P 15731901

- **Animals:** the pathway is the **terminal segment of Tyr/Phe catabolism**, restricted largely to **liver and kidney**, and gated developmentally and nutritionally (induced by dietary aromatic amino-acid load). - **Protein-family expansion (step 2):** MAAI sits inside the large **glutathione transferase (GST) superfamily**. The **zeta class (GSTZ1)** is the member that carries the ancestral isomerase activity; the alpha/mu/pi classes are later, lineage-specific expansions specialized for xenobiotic conjugation and are **not** good proxies for the ancestral

homogentisate-pathway role. For understanding the ancestral function, **GSTZ1/zeta is the representative member**

(
P 21303221
).

Cell-type / physiological state. GSTZ1/MAAI expression and activity depend on age and intracellular chloride, and are highest in liver

(
P 37742772
).

In mammals, the primary target organ of pathway dysfunction is the **kidney** (proximal tubule) for MAA-related accumulation and the **liver** for FAA/ succinylacetone toxicity

(
P 12052898
).

(
P 15277241
).

Alternative routes to the same outcome. The *cis*→*trans* isomerization can be achieved by (i) the GSTZ1 enzyme or (ii) a **non-enzymatic glutathione-mediated** reaction — molecularly different means to the same chemical end

(
P 12052898
).

At the whole-pathway level, **gentisate catabolism** achieves the analogous outcome (aromatic-ring opening → fumarate + pyruvate) by a *parallel* enzyme set, representing a convergent solution rather than a variant of this pathway.

6. Constraints, Dependencies, and Failure Modes

Mandatory ordering. Ring cleavage → isomerization → hydrolysis is fixed by substrate chemistry (Section 3). FAH cannot act on an intact ring; the hydrolase requires the *trans* isomer.

Compartment / cell-type constraints (mammals). The complete pathway operates in hepatocytes and renal proximal tubule cells; GSTZ1 is predominantly cytosolic (several-fold higher in cytoplasm than mitochondria;

(
P 37742772
).

Failure modes and their differential severity — an informative asymmetry. - **Loss of step 1 (HGD): alkaptonuria** — homogentisate accumulates, auto-oxidizes, polymerizes, and deposits as ochronotic pigment in cartilage, connective tissue, heart valves, leading to arthropathy and valvular disease

([P 41236909](#))

([P 40938761](#))

Severe, but slowly progressive. - **Loss of step 2 (MAAI/GSTZ1): no clear human disease;** knockout mice are viable, thanks to the glutathione bypass, though they accumulate FAA and succinylacetone and show liver/kidney pathology under aromatic-amino-acid overload

([P 12052898](#))

([P 15277241](#))

([P 37742772](#))

This is the strongest evidence that the isomerase, uniquely among the three, is **facultative**. - **Loss of step 3 (FAH): hereditary tyrosinemia type I (HT1)** — the most severe error, causing accumulation of FAA and the derived toxin **succinylacetone**, with hepatic failure, renal tubulopathy, neurologic crises, and hepatocellular carcinoma risk

([P 10508789](#))

([P 31611405](#))

Most HT1 mutations act by **destabilizing the active FAH dimer and accelerating aggregation**, not merely by ablating catalysis

([P 31300554](#))

Evidence ruling out otherwise-plausible paths. - The **MAAI/FAH double-knockout is rapidly lethal**, demonstrating that MAA→FAA *does* proceed in the absence of MAAI (via GSH) — otherwise the double mutant would be no worse than the FAH single mutant

([P 12052898](#))

This simultaneously proves the bypass exists and that FAA is the toxic bottleneck. - **Nitisinone (HPPD inhibition) rescues HT1** by shutting off homogentisate production upstream, confirming that toxicity flows through the pathway rather than from tyrosine itself, and that

the three core enzymes are downstream of the druggable choke point
 ([P 41003839](#)),
 ([P 33375092](#)).

7. Controversies and Open Questions

1. **Why is there no human MAAI-deficiency disease?** The dominant explanation is the glutathione non-enzymatic bypass
 ([P 12052898](#)),
 but whether additional enzymatic redundancy exists, and how efficient the bypass is in human liver in vivo, remains incompletely quantified
 ([P 37742772](#)).
2. **HGD structural mechanism at atomic resolution.** The ring-cleavage mechanism rests substantially on computational models (DFT/QM-MM;
 ([P 16332080](#))
 ([P 27119315](#))
 and older crystallography; the protonation state of the substrate C2 hydroxyl (ionized vs non-ionized pathway) and the precise regiochemical control are still debated.
3. **FAH oligomeric state and pathogenicity.** The monomer–dimer equilibrium and aggregation model for HT1
 ([P 31300554](#))
 reframes many mutations as folding/stability defects, motivating pharmacological-chaperone strategies — but which mutations are catalytic vs conformational is not fully mapped.
4. **Moonlighting vs. metabolic roles of GSTZ1.** A large, partly separate literature links GSTZ1 to ferroptosis, NRF2/GPX4 redox signaling, DCA pharmacology, and cancer
 ([P 33931597](#))

P 36905252

P 35562974

P 35979621

How much of this reflects the isomerase's metabolic function (accumulating tyrosine intermediates) versus independent activities of the same protein is unresolved and a frequent source of cross-organism/cross-assay conflation.

5. **Cross-organism extrapolation.** Mechanistic claims often mix data from humans, mice, fungi, and bacteria. The enzymology is conserved, but regulation, tissue distribution, gene clustering, and physiological consequences differ; claims should be anchored to the organism studied.
6. **Residual disease under nitisinone.** Even with upstream HPPD blockade, HT1 models retain molecular remnants of liver disease and HCC risk

(

P 36980965

), indicating incomplete understanding of the toxic intermediates and their long-term signaling effects.

8. Key References

- Timm DE et al. *Crystal structure and mechanism of a carbon-carbon bond hydrolase* (FAH fold, Glu-His-water triad, products). Nat Struct Biol 1999.

P 10508789

- Bateman RL et al. *Mechanistic inferences from the crystal structure of FAH with a bound phosphorus-based inhibitor* (tetrahedral intermediate, Mg²⁺). PNAS 2001.

P 11154690

- Bateman RL et al. *Slow-onset inhibition of FAH by phosphinate transition-state mimics* (nM inhibitors, in vivo phenocopy). Biochem J 2007.

P 17064256

- Macias I et al. *HT1-associated FAH mutations reduce stability and increase aggregation* (dimer/monomer equilibrium). J Biol Chem 2019.

P 31300554

- Borowski T et al. *Catalytic reaction mechanism of homogentisate dioxygenase: DFT study* (peroxo → O–O homolysis → ring scission). 2005.

P 16332080

- Qi C et al. *QM/MM study of aromatic ring cleavage by homogentisate dioxygenase*. 2016.

P 27119315

- Fernández-Cañón JM et al. *MAAI/GSTZ-deficient mice reveal a glutathione- dependent non-enzymatic bypass* (viability, double-mutant lethality). PNAS 2002.

P 12052898

- Board PG, Anders MW. *Glutathione transferase zeta: discovery, catalysis, Gsta1^{-/-} mice* (MAAI = GSTZ1). 2011.

P 21303221

- Lim CE et al. *Gsta1^{-/-} mice: pathology and compensatory GST induction* (succinylacetone accumulation). 2004.

P 15277241

- Stacpoole PW. *Clinical physiology and pharmacology of GSTZ1/MAAI* (no clear human deficiency disease; DCA; tissue distribution). 2023.

P 37742772

- Schmalzer-Ripcke J et al. *Pyomelanin via tyrosine degradation in A. fumigatus* (hmgA/hppD deletions). 2009.

P 19028908

- Perez-Cuesta U et al. *Melanin and pyomelanin in A. fumigatus* (hppD, hmgX, hmgA, fahA, maiA, hmgR cluster). 2020.

P 31020477

- Gunsch CK et al. *Homogentisate-1,2-dioxygenase in Exophiala lecanii-corni* (aromatic pollutant funnel to homogentisate). 2005.

P 15731901

- Azami M et al. *HGD gene analysis in alkaptonuric ochronosis* (gene structure).

P 41236909

- Mascarello A et al. *Nitisinone reduces succinylacetone in HT1* (upstream HPPD block). 2025.

P 41003839

Uncertainty statement

The enzymology of the three core reactions is strongly supported by crystallography, transition-state-mimic inhibitors, computational mechanism, and genetic knockouts. The greatest uncertainties are (i) the in vivo efficiency and molecular basis of the MAAI-independent glutathione bypass in humans, (ii) the atomic-resolution details/regiochemical control of HGD ring cleavage, and (iii) disentangling GSTZ1's metabolic isomerase role from its reported redox/xenobiotic "moonlighting" activities. Regulatory and physiological features vary substantially across bacteria, fungi, and mammals and should not be extrapolated between them.