

GCNT1 Nuclear Speck (GO:0016607) — Function-Assignment Hypothesis Review

Gene: GCNT1 (Q02742), *Homo sapiens* — Beta-1,3-galactosyl-O-glycosyl-glycoprotein beta-1,6-N-acetylglucosaminyltransferase (core-2 GlcNAc-transferase, C2GnT-L) **Focus type:** function_assignment | **Hypothesis slug:** function-hypothesis-go-0016607 **Seed hypothesis:** "GCNT1 has nuclear speck (GO:0016607)." **Annotation under review:** GO:0016607 nuclear speck, evidence IDA, original reference GO_REF:0000052 (Human Protein Atlas immunofluorescence; UniProt cross-reference evidence code IDA:HPA).

Summary

Verdict: Weakly supported / over-annotated — do not treat as a core localization. The proposed cellular-component assignment of GCNT1 to the **nuclear speck (GO:0016607)** rests on a **single Human Protein Atlas (HPA) immunofluorescence antibody pattern**, imported automatically under evidence code IDA with reference GO_REF:0000052. It is contradicted by the strong, independent, and mechanistically coherent evidence establishing GCNT1 as a **single-pass type II Golgi membrane glycosyltransferase** whose catalytic domain faces the Golgi lumen. Such a protein has no known trafficking route into the nuclear interior and no plausible role in membraneless nuclear-speckle condensates, which are liquid–liquid phase-separated assemblies of splicing factors. No primary literature reports a nuclear pool or nuclear function of GCNT1, and the enzyme's established biochemistry — core-2 O-glycan branching using lumenal UDP-GlcNAc — is a Golgi-luminal process.

A comparative check of HPA subcellular records across twelve Golgi type II glycosyltransferases (GCNT1–4, B3GNT6, C1GALT1, ST3GAL1, ST6GALNAC1, GALNT1, MGAT1, B4GALT1, FUT8) reinforces this reading. GCNT1 is the **only** enzyme in the panel (1/12) reported with "Nuclear speckles," it is reported as the **sole main location**, and — uniquely — it carries **no Golgi call at all** in HPA immunofluorescence. Its peers with informative HPA IF data recover the Golgi as expected, and three that also show a "Nucleoplasm" background call still retain the Golgi, reflecting a recognized HPA nuclear-artifact tendency for this membrane-

enzyme class. The complete substitution of GCNT1's biochemically established Golgi compartment by an outlier speckle call is the signature of an antibody artifact, not a genuine dual localization.

Curation lead (requires curator verification): Treat GO:0016607 as **non-core** for GCNT1 — a candidate for **removal** or a **NOT qualifier** pending orthogonal validation (second independent antibody, knockout-validated stain, endogenous tagging, or fractionation). Retain the Golgi CC terms (GO:0000139 Golgi membrane, GO:0005802 trans-Golgi network, GO:0031985 Golgi cisterna) and the core-2 branching MF/BP terms as the gene product's true location and function. The nuclear-speck term should not influence the gene review as a bona fide localization.

Key Findings

Finding 1 — The nuclear-speck term is a single unvalidated HPA immunofluorescence call, at odds with GCNT1's Golgi membrane topology

GCNT1 (UniProt Q02742) is a **428-residue single-pass type II membrane protein** with a short cytoplasmic tail (residues 1–9), a transmembrane anchor (residues 10–32), and a large luminal catalytic domain (residues 33–428). Its curated UniProt subcellular location is **Golgi apparatus membrane**, and its GO cellular-component annotations are dominated by Golgi compartments: Golgi cisterna (GO:0031985, IDA), Golgi membrane (GO:0000139, EXP), and trans-Golgi network (GO:0005802, IDA). Against this backdrop, the nuclear-speck term (GO:0016607) is supported by exactly one line of evidence — **IDA:HPA**, reference **GO_REF:0000052** — i.e., an automated import of a Human Protein Atlas immunofluorescence result. In the HPA record (ENSG00000187210), "Nuclear speckles" is listed as the sole main subcellular location with an IF reliability of "Supported," while the immunohistochemistry reliability for the same gene is only "Uncertain."

The membrane topology is decisive. A type II Golgi glycosyltransferase is inserted into the Golgi membrane with its catalytic domain in the lumen; it is synthesized on the rough ER and trafficked through the secretory pathway, with no signal, mechanism, or precedent for entering the nucleus, let alone partitioning into membraneless nuclear-speckle condensates. The original molecular cloning of GCNT1 explicitly reported type II membrane topology:

"The cDNA sequence predicts a protein with type II membrane topology, as has been found for all other mammalian glycosyltransferases cloned to date." — PMID: 1329093

No primary study reports a nuclear pool or nuclear activity for GCNT1. The enzyme's biochemical function — building the core-2 branch of mucin-type O-glycans by transferring GlcNAc from UDP-GlcNAc onto Gal β 1-3GalNAc-R (PMID: 1329093; PMID: 23027862) — is intrinsically a Golgi-luminal reaction dependent on the nucleotide-sugar donor and the Golgi glycosylation machinery. There is no substrate, donor, or acceptor for this chemistry in nuclear speckles.

Finding 2 — GCNT1 is the lone Golgi glycosyltransferase HPA calls "nuclear speckles," and it uniquely misses the Golgi entirely

To test whether the nuclear-speckle call is a systematic property of Golgi glycosyltransferases or an isolated outlier, HPA subcellular records were compared across twelve Golgi type II glycosyltransferases: GCNT1, GCNT2, GCNT3, GCNT4, B3GNT6, C1GALT1, ST3GAL1, ST6GALNAC1, GALNT1, MGAT1, B4GALT1, and FUT8.

Two features single GCNT1 out:

- 1. GCNT1 is the only gene (1/12) reported with "Nuclear speckles,"** and it is reported as the **sole main location**.
- 2. GCNT1 uniquely carries no Golgi call in HPA immunofluorescence.** Enzymes in the panel with informative HPA IF data (GCNT3, ST6GALNAC1, B4GALT1, FUT8, GCNT4) show "Golgi apparatus" as expected. Three of them (GCNT4, ST6GALNAC1, FUT8) additionally show a "Nucleoplasm" background call — a known HPA nuclear-artifact pattern for membrane proteins — but all of them still recover the Golgi.

Gene	HPA main location	HPA all locations
GCNT1 (target)	Nuclear speckles	Nuclear speckles (no Golgi)
GCNT3	Golgi apparatus	Golgi apparatus, Vesicles
GCNT4	Golgi apparatus	Nucleoplasm, Golgi apparatus
ST6GALNAC1	Golgi apparatus	Nucleoplasm, Golgi apparatus
B4GALT1	Golgi apparatus	Golgi apparatus
FUT8	Golgi apparatus	Nucleoplasm, Golgi apparatus, Cytosol
GCNT2, B3GNT6, C1GALT1, ST3GAL1, GALNT1, MGAT1	(no HPA IF loc data)	—

GCNT1 is therefore doubly anomalous: it gains an implausible nuclear-speckle call **and** loses the compartment where its biochemistry is known to occur. A genuinely dual-localized protein would be expected to retain a Golgi signal alongside any secondary compartment. GCNT1's speckle call **replaces** rather than **supplements** the expected Golgi signal — the classic signature of an antibody that binds an off-target, speckle-enriched epitope while failing to detect the true Golgi target. The nucleoplasm background calls in three other Golgi GTs confirm that HPA has a recognized nuclear-artifact tendency for this membrane-enzyme class, reinforcing the artifact interpretation.

Mechanistic Model / Interpretation

The competing interpretations can be laid out compactly:

SEED HYPOTHESIS

GCNT1 -> nuclear speck
(GO:0016607)|
| single HPA IF antibody
| evidence IDA (GO_REF:0000052)

v

membraneless splicing-factor
condensate in the nucleus|
| no donor, no acceptor,
| no trafficking route,
| no primary literature

v

NOT MECHANISTICALLY VIABLE

ESTABLISHED BIOLOGY

GCNT1 = type II Golgi membrane enzyme
cytoplasmic tail (1-9)
TM anchor (10-32)
luminal catalytic domain (33-428)

|

v

core-2 O-glycan branching
UDP-GlcNAc -> Galb1-3GalNAc-R
in the GOLGI LUMEN

|

v

<a href="https://pubmed.ncbi.nlm.nih.gov/1329093/" rel="

<a href="https://pubmed.ncbi.nlm.nih.gov/9915862/" rel="

Nuclear speckles (GO:0016607) are interchromatin granule clusters enriched in pre-mRNA splicing factors (e.g., SRRM2/SC35, SR proteins). They are liquid-liquid phase-separated bodies, not membrane compartments, and their resident proteins characteristically have intrinsically disordered regions and RNA-binding capacity — for example, galectin-3 partitions into speckles via an intrinsically disordered domain and ribonucleoprotein association ([PMID: 37003559](#)). GCNT1 has none of these properties: it is a folded, membrane-anchored, luminal glycosyltransferase with no reported disordered speckle-targeting region and no RNA-binding role.

The most parsimonious explanation is that the HPA antibody produces a punctate nuclear pattern that the automated HPA pipeline classified as "nuclear speckles," while failing to detect the true Golgi pool. This yields a single-antibody, single-modality IDA annotation that propagated into GO via GO_REF:0000052. It is exactly the class of annotation that hypothesis-level review is designed to catch: technically an IDA, but biologically an uncorroborated localization that conflicts with the gene product's core molecular identity.

Evidence Base

Citation	Evidence type	Supports/ Refutes/ Qualifies	Claim tested	Key finding	Context
HPA ENSG00000187210 (GO_REF:0000052)	Localization (IF)	Supports (sole basis)	GCNT1 in nuclear speckles	Main location "Nuclear speckles"; Reliability(IF)= Supported ; Reliability(IH)= Uncertain	Cultured human cell lines, single antibody IF
UniProt Q02742 (curated)	Structural/ topology, curated CC	Refutes/ qualifies	Where does GCNT1 reside	Golgi apparatus membrane; type II single- pass TM; luminal catalytic domain (33–428); GO CC Golgi cisterna/membrane/ TGN	Human, Swiss-Prot review
PMID: 1329093	Direct assay + sequence	Refutes	Molecular nature of GCNT1	Type II membrane topology; forms core-2 O- glycan branch	Expression cloning, CHO/COS
PMID: 23027862	Enzymatic characterization	Qualifies (function only)	Catalytic activity	Basis of UniProt FUNCTION (core-2 branching); no nuclear claim	In vitro / cellular glyco-assay
PMID: 9915862	Direct assay	Qualifies (context)	C2GnT family function	C2GnT enzymes build core 2/4/I branches on O- glycans	Human cloning, expression
PMID: 8308016	Direct assay; transfection	Qualifies (context)	C2GnT O- glycan output	C2GnT directs core-2 branched O-glycans and poly-LacNAc on cell- surface mucins	CHO transfection
PMID: 37003559	Review	Qualifies (competing concept)	What resides in speckles	Speckles are LLPS bodies of SR/splicing factors;	Review

Citation	Evidence type	Supports/ Refutes/ Qualifies	Claim tested	Key finding	Context
				residents have IDRs & RNA association	

Literature narrative. The foundational cloning paper (PMID: 1329093) establishes GCNT1's type II membrane topology and core-2 branching activity, directly refuting a soluble nuclear localization. The C2GnT family characterizations (PMID: 9915862; PMID: 8308016) situate GCNT1 within a family of Golgi O-glycan branching enzymes, none of which is a nuclear/speckle protein. The speckle-biology review (PMID: 37003559) defines the physical properties of genuine speckle residents — properties GCNT1 does not have — underscoring the implausibility of the seed hypothesis.

GO Curation Implications

Lead requiring curator verification.

- **Cellular component (CC):** Retain the Golgi set as core — **GO:0031985 (Golgi cisterna)**, **GO:0000139 (Golgi membrane)**, **GO:0005802 (trans-Golgi network)** — supported by curated UniProt location and consistent with type II topology. **Do not treat GO:0016607 (nuclear speck) as core.** Recommended action: downgrade to non-core; candidate for **removal** or a **NOT qualifier**, on the grounds that a single unvalidated HPA IF pattern conflicts with strong, independent Golgi/topology evidence.
- **Molecular function (MF):** Unaffected and well-supported — core-2 β 1,6-N-acetylglucosaminyltransferase activity (UDP-GlcNAc-dependent branch formation).
- **Biological process (BP):** Unaffected — mucin-type O-glycosylation / core-2 branch biosynthesis.
- Do **not** fall back to "protein binding"; the informative, supported CC terms (Golgi membrane / trans-Golgi network) already exist and should be retained.

Final Curation Decision Table (lead — verify)

GO term	Aspect	Current evidence	Recommendation
GO:0016607 nuclear speck	CC	IDA:HPA, single antibody, GCNT1-only outlier, no Golgi in HPA, no primary lit	Non-core → remove or NOT-qualify pending orthogonal validation
GO:0000139 Golgi membrane	CC	EXP:UniProtKB	Retain as core
GO:0005802 trans-Golgi network	CC	IDA:UniProtKB	Retain as core
GO:0031985 Golgi cisterna	CC	IDA:UniProtKB	Retain
core-2 branching activity / O-glycan process	MF/BP	P 1329093 , P 23027862	Retain (primary function)

Mechanistic Scope

The immediate molecular property under test is **subcellular localization**, not catalytic activity. The seed hypothesis asserts that GCNT1 physically resides in nuclear speckles. This must be separated from:

- **Direct gene-product activity:** GCNT1's actual direct activity is Golgi-luminal core-2 O-glycan branch synthesis (core-2 β 1,6-GlcNAc branching of mucin-type O-glycans using lumenal UDP-GlcNAc). This is well established and not in dispute, and is spatially incompatible with nuclear speckles.
- **Downstream phenotypes:** O-glycan branching affects mucin structure, selectin-ligand formation, and cell-surface glycan display — all downstream of the Golgi reaction and unrelated to a nuclear location.
- **Inference from a single stain:** The nuclear-speckle claim is inferred entirely from one antibody's staining pattern, with no functional readout in the nucleus and no described

mechanism. If real, it would be a secondary observation, not the gene product's primary function or location.

Conflicts and Alternatives

- **Antibody artifact (most likely):** Punctate nuclear staining is a frequent HPA false-positive class; HPA's own IHC reliability for GCNT1 is "Uncertain." The complete absence of a Golgi call for GCNT1 — unique in the 12-gene panel — is exactly what an off-target antibody would produce.
 - **HPA nuclear-background pattern:** Several membrane glycosyltransferases in the panel (GCNT4, ST6GALNAC1, FUT8) show a "Nucleoplasm" background call while still recovering the Golgi; GCNT1's speckle call may be an extreme instance of the same phenomenon, aggravated by loss of the Golgi signal.
 - **Paralog confusion (unlikely to rescue):** GCNT1/2/3/4 and the broader C2GnT family are all Golgi type II enzymes; none is a known nuclear/speckle protein ([PMID: 9915862](#)).
 - **Isoform (no support):** No described soluble/nuclear isoform of GCNT1 lacking the signal-anchor exists.
 - **Reliability discordance:** IF "Supported" vs. IHC "Uncertain" indicates the localization is not robustly reproduced across methods.
-

Limitations and Knowledge Gaps

1. **Orthogonal validation of the HPA pattern.** Checked: HPA reliability tiers (IF "Supported," IHC "Uncertain"). Not resolved because no paired-antibody or knockdown IF is reported. Why it matters: single-antibody localizations are the weakest IDA class. Resolve with a second independent antibody plus siRNA/CRISPR knockout IF, or endogenous tagging.
2. **Any endogenous nuclear pool.** Checked: primary literature (none reported). Why it matters: an orthogonal detection of nuclear GCNT1 would change the verdict. Resolve with subcellular fractionation plus mass spec / western blot.
3. **Raw HPA image not re-examined.** The assessment relies on HPA's classified output and reliability flags rather than a fresh manual read of the primary micrographs. Resolve with expert re-annotation of the HPA IF images.

4. **GO_REF:0000052 text not directly retrieved.** The official GO_REF markdown could not be fetched programmatically (HTTP 403/404 from current.geneontology.org and go-site raw/API). However, the UniProt cross-reference for GO:0016607 is explicitly attributed as **IDA:HPA**, which unambiguously identifies the source as HPA immunofluorescence — consistent with GO_REF:0000052's known scope. No fabrication: the GO_REF text itself was not accessed.

Discriminating Tests

The following would most efficiently distinguish the seed hypothesis from the antibody-artifact alternative:

1. **Knockout/knockdown-validated immunofluorescence** with ≥ 2 independent antibodies — does the speckle signal disappear on GCNT1 loss?
2. **Endogenous fluorescent tagging** (CRISPR knock-in) to visualize native localization without antibody bias (expected: Golgi, not speckles).
3. **Subcellular fractionation** (nuclear vs. membrane) with orthogonal detection (western/MS) to quantify any nuclear GCNT1 pool.
4. **Co-localization** with a bona fide speckle marker (SRRM2/SC35) versus a Golgi marker (GM130/TGN46) — a real speckle protein co-localizes with SRRM2; GCNT1 should co-localize with the Golgi marker.
5. **Sequence/feature check** for intrinsically disordered or RNA-binding speckle-targeting determinants ([PMID: 37003559](#)); their absence argues against genuine speckle residence.

Curation Leads (require curator verification)

- **Candidate action change:** Demote or remove **GO:0016607 (nuclear speck, IDA, GO_REF:0000052)** from GCNT1, or apply a **NOT** qualifier, pending orthogonal validation. Do not let it stand as a core localization.
- **Retain as core:** Golgi CC terms (GO:0031985, GO:0000139, GO:0005802) and core-2 β 1,6-GlcNAc-transferase MF/BP terms.

- **Reference to verify (topology/identity):** [PMID: 1329093](#) — snippet: *"The cDNA sequence predicts a protein with type II membrane topology, as has been found for all other mammalian glycosyltransferases cloned to date."*
 - **References for family function:** [PMID: 9915862](#) and [PMID: 8308016](#) — establish the Golgi core-2/O-glycan branching role of the C2GnT family.
 - **Suggested curator question:** Is the HPA nuclear-speckle call supported by any orthogonal (second-antibody or knockdown) data, or is it a single-antibody pattern? If single-antibody, treat as insufficient against the strong Golgi prior.
 - **Suggested experiments:** Second-antibody + KO IF; endogenous tagging; Golgi/nuclear fractionation with MS; SRRM2 vs. GM130 co-localization.
-

Proposed Follow-up Experiments / Actions

1. **Curation:** Flag GO:0016607 for GCNT1 as non-core; recommend removal or NOT qualifier subject to curator confirmation of HPA antibody count and modality.
 2. **Validation experiment (highest value):** Second-antibody IF in GCNT1-knockout cells to test antibody specificity of the speckle signal.
 3. **Orthogonal detection:** Endogenous tagging or subcellular fractionation + quantitative MS to determine whether any nuclear GCNT1 pool exists.
 4. **Co-localization:** SRRM2/SC35 vs. Golgi-marker two-color imaging to place GCNT1 definitively in the Golgi or in speckles.
 5. **Documentation:** Record the comparative HPA outlier analysis (GCNT1 as 1/12 speckle-called, uniquely Golgi-negative) as provenance supporting the non-core recommendation.
-