



# Combinatorial Fake Discrimination Studies For Combined dE/dx and Disappearing Track Search

## Introduction

- The Standard Model is a highly accurate theory of physics
- Search for long-lived, massive, charged particles that decay into a dark matter candidate
- Interested in a way to select against background in favor of signal




Example signal Process: charginos decaying into pions and neutralinos (N. Young)




- Previous analyses have limited charginos up to masses of 850 GeV
- We seek to combine two analysis strategies to gain further sensitivity

## Disappearing Track + dE/dx



- Analysis uses data collected from the ATLAS experiment inner detector at the Large Hadron Collider


- Using a combination disappearing track and high ionization energy loss (dE/dx) signature
- Much of the background that dominated the Run 2 disappearing track analysis will be suppressed by ionization energy requirements
- Many fake tracklets are expected to remain, forming the bulk of our background



Disappearing track signature, a particle decays in between Pixel and SCT detectors. Also pictured are possible background processes that produce similar looking signatures (ATLAS)

## Clusters + dE/dx

- Tracks formed up of individual clusters
  - Grouped excitations of pixel detectors
- Ionization energy loss is recorded in these
- Comparison of the calculated dE/dx value indicates a good discriminatory power between background and signal
- Searching for a selection that maximizes the strength of a signal we would observe.




$N_c$	Cluster pattern	$n_{IBL}^{OF}$	Truncation pattern	$n_{used}$
1	X	0 or 1	X	N/A
2	X,X	0 or 1	X	1
3	C,C,X	0 or 1	C,C	2
3	C,O,X	1 or 2	C	1
3	O,X,X	1,2,3	O	1
4	C,C,C,X	0 or 1	C,C,C	3
4	C,C,O,X	1 or 2	C,C	2
4	C,O,X,X	1,2,3	C,O	2
4	O,X,X,X	1,...,4	O,X	2
≥ 5	X,X,X,X,X,...	0,...,N <sub>c</sub>	X,X,X,X,...	N <sub>c</sub> - 2



Truncation pattern lookup table for track dE/dx average. Truncation pattern is based on number of clusters and the number of IBL overflow hits a track has. C=Normal Hits, O=IBL overflow hit, X=Either (ATLAS)

- Can convert a tracks cluster's dE/dx values into a truncated dE/dx via an algorithm
- What information can we use from the full set of dE/dx values?

## Selection Parameters




- With the additional parameters we can now look to select on, want to investigate distribution of several parameters in order to see which might offer the best discriminatory power
- Looking at many different parameters
  - dE/dx of dropped hits
  - Layer in which high hit occurs
  - Layer that gets dropped during truncation
  - RMS of all cluster dE/dx values



## Preliminary Significance Studies

- We can identify an optimal parameter to select on by seeing which gives the best significance score for a selection in the high dE/dx regime

$$Z_A = \sqrt{2 \left( (s+b) \log \left( 1 + \frac{s}{b} \right) - s \right)}$$



- Preliminary results indicate that the dropped layer of a tracklet could be a potentially powerful discriminant
- This may suggest that low momentum tracklets are adding to the energy deposited in the first layers

## Conclusion/Next Steps

- Searching for long-lived, massive, charged particles in the ATLAS detector
- Identified potential selection criteria to optimize discovery for further study using cluster level information
- Want to better understand behavior of high dE/dx regime
  - How does cluster size relate to high dE/dx hits in innermost layers of detector?

## References

- ATLAS Collaboration. Search for long-lived charginos based on a disappearing-track signature using 136 fb-1 of pp collisions at  $\sqrt{s} = 13$  TeV with the ATLAS detector.
- ATLAS Collaboration. Search for heavy, long-lived, charged particles with large ionisation energy loss in  $pp$  collisions at  $\sqrt{s} = 13$  TeV using the ATLAS experiment and the full Run 2 dataset.
- Lory, Alexander. Search for new physics in signatures of soft unclustered energy patterns within the ATLAS detector
- J.J. Goodson. "Search for Supersymmetry in States with Large Missing Transverse Momentum and Three Leptons including a Z-Boson". Presented 17 Apr 2012. PhD thesis. Stony Brook University, 2012.

## Acknowledgements

I would like to thank Dr. Laura Jeanty and Nathan Young for their continued mentorship and support of me during this project. I think I am also supposed to mention that this material is based upon work supported by the U.S. Department of Energy, Office of Science, Office of High Energy Physics program under Award Number DE-SC0020244