

Notto - Category for the Working Mathematician

tkunn

August 2020

This is solutions for problems on the second edition of *Category for the Working Mathematician* by S. Mac Lane. Problems are placed in the first line of each sections. They may be modified from the original text for simplicity, clarity and some laziness.

1

1.1

$$\langle \wedge _ \wedge \rangle$$

1.2

$$\langle Q \perp Q \rangle$$

1.3

1.3.1

TODO

1.3.2

TODO

1.3.3

TODO

1.3.4

TODO

1.3.5

TODO

1.4

1.4.1

TODO

1.4.2

TODO

1.4.3

TODO

1.4.4

TODO

1.4.5

TODO

1.4.6

TODO

1.5

1.5.1

TODO

1.5.2

TODO

1.5.3

TODO

1.5.4

TODO

1.5.5

TODO

1.5.6

TODO

1.5.7

TODO

1.5.8

TODO

1.5.9

TODO

1.6

Let U be a set that satisfies following conditions:

- (i) $x \in u \in U \Rightarrow x \in U$
- (ii) $(u \in U \wedge v \in U) \Rightarrow (\{u, v\}, \langle u, v \rangle, u \times v \in U)$
- (iii) (1) $x \in U \Rightarrow \mathcal{P}x \in U$
 (2) $x \in U \Rightarrow \bigcup x \in U$
- (iv) $\omega \in U$, where $\omega = \{0, 1, 2, \dots\}$ is a set of all finite ordinal numbers.
- (v) If there exists a surjection $f : a \rightarrow b$ and $a \in U$ and $b \subset U$, then $b \in U$.

1.6.1

Let $I \in U$, $f : I \rightarrow b$ and $f_i \in U$ for all $i \in I$. Prove that $\prod_i f_i \in U$.

For all $q \in \prod_i f_i$, we can construct a bijection $r : I \rightarrow q$. $q \subset U$ because $\forall w \in q, \exists j \in I, w \in f_j \in U$. Since $I \in U$ and $q \subset U$, we can say $q \in U$. Therefore $\prod_i f_i \subset U$.

Let $|f_k| \geq |f_i|$ for all i . Then, we can construct a surjection $g : f_k^I \rightarrow \prod_i f_i$. Also, we can construct a surjection $h : X \rightarrow f_k^I$, with X is either $\mathcal{P}f_k$ or $\mathcal{P}I$. As $X \in U$, $g \circ h : X \rightarrow \prod_i f_i$ and $\prod_i f_i \subset U$, we can say $\prod_i f_i \in U$.

1.6.2

(a) Let $I \in U$, $f : I \rightarrow b$ and $f_i \in U$ for all $i \in I$. Prove $\bigcup_i f_i \in U$.

(b) Prove that (a) implies following if (i), (ii), (iii)(1), (iv) and (v) holds true:

(iii)(2) $x \in U \Rightarrow \bigcup x \in U$.

(v) If $f : a \rightarrow b$ is surjective and $a \in U$ and $b \subset U$, then $b \in U$.

(a) We can construct a bijection $g : I \rightarrow \{f_i \mid i \in I\}$, $g(i) = f_i$. As $I \in U$ and $f_i \in U$ for all i , $\{f_i \mid i \in I\} \in U$. Therefore $\bigcup_i f_i = \bigcup \{f_i \mid i \in I\} \in U$.

(b) Because $x \in U$, we have $y \in U$ for all $y \in x$. Therefore we can apply $f : x \rightarrow x$, $f(i) = i$ to (a) to get $\bigcup x \in U$.

Because $a \in U$ and $b \subset U$, we can apply f to (a) to get $\bigcup_i f_i \in U$. f is surjective, therefore $b = \bigcup_i f_i$. Hence $b \in U$.

1.7

$$\langle Q \smile Q \rangle$$

1.8



$$\phi \longrightarrow \theta$$


2

2.1



$$\phi \curvearrowright \theta$$

2.2



2.3

2.3.1

Show product of categories includes product of monoids, product of groups and product of sets.

In this section, \times_S is a product operation for sets, \times_C is for categories, \times_G is for groups and \times_M is for monoids.

Monoids. Let M, N be a monoid with object m, n respectively. The only object in $M \times_C N$ is $\langle m, n \rangle$ (TODO)

Groups. Let G, H be a group with object a, b respectively. $G \times_C H$... (TODO)

Sets. Let A, B be discrete categories. The set of all objects in $A \times_C B$ is $X = A \times_S B$. The set of all arrows in $A \times_C B$ is $\{\langle f, g \rangle \mid a \in A, b \in B, f \in A(a, a) \wedge g \in B(b, b)\} = \{\langle \text{id}_A(c_0), \text{id}_B(c_1) \rangle \mid c \in X\}$. Therefore $A \times_C B$ is a discrete category of $X = A \times_S B$.

2.3.2

Prove that product of two preorders is preorder.

Let P, Q be preorders. $\forall a \forall b \mid \text{hom}(a, b) \mid \leq 1$ for both P and Q . Therefore, for all $p_1 \in P, p_2 \in P, q_1 \in Q, q_2 \in Q$, $\mid \text{hom}_{P \times Q}(\langle p_1, q_1 \rangle, \langle p_2, q_2 \rangle) \mid = \mid \text{hom}_P(p_1, p_2) \times \text{hom}_Q(q_1, q_2) \mid \leq 1$. Hence $P \times Q$ is preorder.

2.3.3

Let $\{C_i \mid i \in I\}$ be a family of categories indexed by a set I . Show product $C = \prod_i C_i$, its projections $P_i : C \rightarrow C_i$ and universal property of these projections.

Let $\text{id}_X(a) : a \rightarrow a$ be identity in category X .

Let the set of all object in C be the product set $\prod_i C_i$ and $C(a, b) = \prod_i C_i(a_i, b_i)$. Let $\text{id}_C(c)_i = \text{id}_{C_i}(c_i)$ for all $c \in C$. Now, we prove C has a universal property:

1. For every i there is a functor $P_i : C \rightarrow C_i$.

2. For every category B such that a functor $G_i : B \rightarrow C_i$ presents for every C_i , there is a functor $F : B \rightarrow C$, which makes the following diagram commute.

$$\begin{array}{ccc}
 B & & \\
 \downarrow F & \searrow G_i & \\
 C & \xrightarrow{P_i} & C_i
 \end{array}$$

First, we prove $P_i : C \rightarrow C_i$ exists. Let the object function be $P_i(x) = x_i$. Let the arrow function be $P_i(f) = f_i$. For all object $c \in C$, $P_i(\text{id}_C(c)) = \text{id}_C(c)_i = \text{id}_{C_i}(c_i) = \text{id}_{C_i}(P_i(c))$. For all arrow f, g in C , $P_i(g \circ f) = (g \circ f)_i = g_i \circ f_i = P_i(g) \circ P_i(f)$. Therefore P_i is a functor.

Second, we prove $F : B \rightarrow C$ exists. Let the object function be $F(x)_i = G_i(x)$. Let the arrow function be $F(f)_i = G_i(f)$. For all object $b \in B$, $F(\text{id}_B(b))_i = G_i(\text{id}_B(b)) = \text{id}_{C_i}(G_i(b)) = \text{id}_{C_i}(F(b)_i)$. Thus $F(\text{id}_B(b)) = \text{id}_C(F(b))$. For all arrow f, g in B , $F(f \circ g)_i = G_i(f \circ g) = G_i(f) \circ G_i(g)$. Thus $F(f \circ g) = F(f) \circ F(g)$. Therefore F is a functor.

2.3.4

Show opposite of \mathbf{Matr}_K .

In \mathbf{Matr}_K , the object set is all positive integers $\{1, 2, 3, \dots\} = \omega \setminus \{0\}$. $\mathbf{Matr}_K(n, m)$ is all rectangular matrix on K of shape $m \times n$. Therefore $\mathbf{Matr}_K^{\text{op}}$ has the same objects $\omega \setminus \{0\}$ and $\mathbf{Matr}_K^{\text{op}}(n, m)$ is all rectangular matrix on K of shape $n \times m$.

2.3.5

Show that the mapping between a topological space and the ring of real continuous functions on it is the object function of a contravariant functor on \mathbf{Top} to \mathbf{Rng} .

Let $R_T \subseteq (T \rightarrow \mathbb{R})$ be a ring whose elements are continuous functions from a topological space T to real number. We construct R_T as follows:

1. Additive identity. $0_{R_T} : x \mapsto 0$.
2. Multiplicative identity. $1_{R_T} : x \mapsto 1$.
3. Addition. $f + g : x \mapsto f(x) + g(x)$.
4. Multiplication. $f \times g : x \mapsto f(x) \times g(x)$.

Let X and Y be any topological spaces. If we have a continuous function $f : Y \rightarrow X$, we can construct a ring homomorphism $H(f) = h : R_X \rightarrow R_Y$. We define $h(r) = r \circ f$. Then $h(0_{R_X}) = 0_{R_Y}$, $h(1_{R_X}) = 1_{R_Y}$, $(h(s + t))(x) =$

$(s+t)(f(x)) = s(f(x)) + t(f(x))$, $(h(s \times t))(x) = (s \times t)(f(x)) = s(f(x)) \times t(f(x))$. Therefore $H(f)$ is a ring homomorphism.

Now we construct a functor $F : \mathbf{Top}^{\text{op}} \rightarrow \mathbf{Rng}$. Let the object function be $F(A) = R_A$ and the arrow function be $F(g) = H(g^{\text{op}})$. For all arrow a, b in \mathbf{Top}^{op} , $F(b \circ a) = H((b \circ a)^{\text{op}}) = H(a^{\text{op}} \circ b^{\text{op}}) = H(b^{\text{op}}) \circ H(a^{\text{op}}) = F(b) \circ F(a)$. For all topological space $T \in \mathbf{Top}^{\text{op}}$, $F(\text{id}(T)) = H(\text{id}(T)) = \text{id}(R_T)$. Therefore F is a functor and \bar{F} is a contravariant functor on \mathbf{Top} to \mathbf{Rng} .

2.4

2.4.1

Show that for any ring R , $R\text{-Mod}$ is a full subcategory of \mathbf{Ab}^R .

TODO

2.4.2

For a finite discrete category X , describe B^X .

For any functor $T : X \rightarrow B$, if its object function is $T(a) = b$, its arrow function maps $\text{id}(a)$ to $\text{id}(b)$. Such a functor T is an object of B^X .

Let $R, S : X \rightarrow B$ be functors and τ be a map on an object of X to an arrow in B . $(\tau : R \rightarrow S) \Leftrightarrow (\forall x \in X, \tau_x(R(x)) = S(x))$. Therefore $\text{hom}(R, S) = \{\tau \mid \forall x \in X, \tau_x(R(x)) = S(x)\}$. Therefore an arrow on R to S exists iff $(\forall x, y \in X, e_R(x, y) \rightarrow e_S(x, y)) \wedge (\forall x \in X, \text{hom}(R(x), S(x)) \neq \emptyset)$, where $e_T(x, y) \Leftrightarrow (\exists a \in B, \{x, y\} \subseteq \{w \mid a = T(w)\})$.

2.4.3

Let \mathbb{N} be a discrete category of natural numbers. Describe $\mathbf{Ab}^{\mathbb{N}}$.

An object of $\mathbf{Ab}^{\mathbb{N}}$ is a map on \mathbb{N} to \mathbf{Ab} . Same as above, $\text{hom}(R, S) = \{\tau \mid \forall n \in \mathbb{N}, \tau_n(R(n)) = S(n)\}$. In other words, a map $\tau : \mathbb{N} \rightarrow (\mathbf{Ab} \rightarrow \mathbf{Ab})$ is an arrow iff, for every $n \in \mathbb{N}$, there is a corresponding group homomorphism τ_n on $R(n)$ to $S(n)$, and, for every $m \in \mathbb{N}$ such that $R(n) = R(m)$, $S(n) = S(m)$.

2.4.4

Let P and Q be preorders. Describe Q^P and show it is a preorder.

Let $R, S : P \rightarrow Q$. Then R and S are objects of Q^P . Let τ be a natural transform $\tau : R \rightarrow S$. τ is an arrow on R to S in Q^P . Since τ is natural and P is preorder, following diagram is commute for every pair of objects $p, p' \in P$. $a = f(p, p')$, where $f(p, p')$ is the only arrow on p to p' . Since Q is preorder, $\tau p = g(Rp, Sp)$, where $g(Rp, Sp)$ is the only arrow on Rp to Sp .

$$\begin{array}{ccc} Rp & \xrightarrow{\tau p} & Sp \\ Ra \downarrow & & \downarrow Sa \\ Rp' & \xrightarrow{\tau p'} & Sp' \end{array}$$

From the two downward arrows in the diagram, we can say that $\text{Im}(R)$ and $\text{Im}(S)$ contain the preorder structure of P . There are two functors $P \rightarrow \text{Im}(R)$ and $P \rightarrow \text{Im}(S)$, where $\text{Im}(T)$ is a category from the image of the object function of T and all arrows between any two pairs in the image.

As explained above, $\sigma p = g(Rp, Sp)$ for all $\sigma : R \rightarrowtail S$. Thus $|\text{hom}(R, S)| \leq |\{\sigma \mid \forall p \in P, \sigma p = g(Rp, Sp)\}| = 1$. Therefore Q^P is preorder.

2.4.5

Let \mathbf{Fin} be a category of all finite sets and G be a finite group. Describe \mathbf{Fin}^G .

Let \mathbf{Fin} be a category of all finite sets. The object is every finite set and the arrow is every mapping between every pair of finite sets.

Let G be a finite group. G is a category of only one object. Every arrow a in G has its inverse a^{-1} such that $a \circ a^{-1} = a^{-1} \circ a = \text{id}$.

\mathbf{Fin}^G is a category that have any functor on G to \mathbf{Fin} as objects and any natural transform between two objects as arrows. The group G has only one object $x \in G$, thus any functor $T \in \mathbf{Fin}^G$ map x to a finite set $T(x) \in \mathbf{Fin}$, and endomorphisms of x to endomorphisms of $T(x)$. For any arrow a, b in G , $T(b \circ a) = T(b) \circ T(a)$. Also, $\text{id} = T(\text{id}) = T(a \circ a^{-1}) = T(a) \circ T(a^{-1})$. Thus any element in the image of the arrow function of T is invertible. Therefore T is a permutation representation of G .

Now, let $\tau : R \rightarrowtail S$ be an arrow in \mathbf{Fin}^G . Then, the following diagram commutes for any arrow a in G :

$$\begin{array}{ccc} Rx & \xrightarrow{\tau x} & Sx \\ Ra \downarrow & & \downarrow Sa \\ Rx & \xrightarrow{\tau x} & Sx \end{array}$$

Therefore, $\text{hom}(R, S) = \{\tau \mid \forall a, \tau x \circ Ra = Sa \circ \tau x\} \dots$ what does it mean????
TODO

2.4.6

TODO

2.4.7

TODO

2.4.8

TODO

2.5

2.5.1

Let \mathbf{A} , \mathbf{B} and \mathbf{C} be small categories. Prove that $\mathbf{Cat}(A \times B, C) \cong \mathbf{Cat}(A, C^B)$ and that they are natural.

First, we prove there is a bijection $f : \mathbf{Cat}(A \times B, C) \rightarrow \mathbf{Cat}(A, C^B)$. Let $f(F) = G$, where $F : A \times B \rightarrow C$ and $G : A \rightarrow C^B$.

We define the object function of G . Let $G(a) = H$, where $H : B \rightarrow C$ and $a \in A$. Let $H(b) = F(\langle a, b \rangle)$ for objects and $H(b) = F(\langle \text{id}(a), b \rangle)$ for arrows. Then for any $b \in B$, $H(\text{id}(b)) = F(\langle \text{id}(a), \text{id}(b) \rangle) = \text{id}(H(b))$. For any arrow x, y in A , $H(x \circ y) = F(\langle \text{id}(a), x \circ y \rangle) = F(\langle \text{id}(a), x \rangle) \circ F(\langle \text{id}(a), y \rangle) = H(x) \circ H(y)$. Therefore H is a functor.

We define the arrow function of G . Let $G(a) = \tau$, where a is an arrow in A and $\tau : B \rightarrow C$. Let $\tau(b) = F(\langle a, \text{id}(b) \rangle)$.

TODO

2.5.2

Let \mathbf{A} , \mathbf{B} and \mathbf{C} be categories. Prove that $(A \times B)^C \cong A^C \times B^C$ and $C^{A \times B} \cong (C^B)^A$.

For any $x \in (A \times B)^C$, there are two functions $G(c) = x(c)_0$ and $H(c) = x(c)_1$. Let the object function be $F(x) = \langle G, H \rangle$, where $F : (A \times B)^C \rightarrow A^C \times B^C$.

For any natural transform $\tau : (A \times B) \rightarrow C$, there are two natural transforms $\tau_0 : A \rightarrow C$ and $\tau_1 : B \rightarrow C$. Let $\tau_0(x) =$ and $\tau_1(x) =$.

Thus we can construct the arrow function $F(\tau) = \langle \tau_0, \tau_1 \rangle$.

2.5.3

2.5.4

2.5.5

2.5.6

2.5.7

2.5.8

2.6

2.7

2.8

3

4

5

6

7

8

9

10

11

12