

We consider a simple two reagent reaction diffusion problem

$$\frac{dQ}{dt} = \gamma f(Q, R) + \nabla^2 Q \quad (1)$$

$$\frac{dR}{dt} = \gamma g(Q, R) + D \nabla^2 R \quad (2)$$

To practice using Deal.ii, our goal will be to simulate this system. The first step is to determine the weak form of the problem, and decide how we will update in time. To keep things simple, let's consider a forward Euler scheme in time which we will eventually adapt into a full IMEX scheme.

$$\frac{Q^{n+1} - Q^n}{\Delta t} = \gamma f(Q^n, R^n) + \nabla^2 Q^n \quad (3)$$

$$\frac{R^{n+1} - R^n}{\Delta t} = \gamma g(Q^n, R^n) + D \nabla^2 R^n \quad (4)$$

We can easily update the diffusion terms implicitly to help with the stability of the method. Therefore, we consider the following expression with  $n$  shifted down by one:

$$\frac{Q^n - Q^{n-1}}{\Delta t} - \nabla^2 Q^n = \gamma f(Q^{n-1}, R^{n-1}) \quad (5)$$

$$\frac{R^n - R^{n-1}}{\Delta t} - D \nabla^2 R^n = \gamma g(Q^{n-1}, R^{n-1}) \quad (6)$$

Now let's project this into the trial space:

$$(\phi_i^n, Q^n) - \Delta t (\phi_i^n, \nabla^2 Q^n) = (\phi_i^n, Q^{n-1}) + \gamma \Delta t f(Q^{n-1}, R^{n-1}) \quad (7)$$

$$(\phi_i^n, R) - D \Delta t (\phi_i^n, \nabla^2 R^n) = (\phi_i^n, R^{n-1}) + \gamma \Delta t g(Q^{n-1}, R^{n-1}) \quad (8)$$

If  $Q$  and  $R$  are approximately equal to their projections in the trial space, then

$$(\phi_i^n, q_j^n \phi_j^n) - \Delta t (\phi_i^n, \nabla^2 q_j^n \phi_j^n) = (\phi_i^n, q_j^{n-1} \phi_j^{n-1}) + \gamma \Delta t f(Q^{n-1}, R^{n-1}) \quad (9)$$

$$(\phi_i^n, r_j^n \phi_j^n) - D \Delta t (\phi_i^n, \nabla^2 r_j^n \phi_j^n) = (\phi_i^n, r_j^{n-1} \phi_j^{n-1}) + \gamma \Delta t g(Q^{n-1}, R^{n-1}) \quad (10)$$

We will assume homogeneous Dirichlet conditions so that after integrating by parts the system reduces to

$$(\phi_i^n, q_j^n \phi_j^n) - \Delta t (\nabla \phi_i^n, \nabla q_j^n \phi_j^n) = (\phi_i^n, q_j^{n-1} \phi_j^{n-1}) + \gamma \Delta t f(Q^{n-1}, R^{n-1}) \quad (11)$$

$$(\phi_i^n, r_j^n \phi_j^n) - D \Delta t (\nabla \phi_i^n, \nabla r_j^n \phi_j^n) = (\phi_i^n, r_j^{n-1} \phi_j^{n-1}) + \gamma \Delta t g(Q^{n-1}, R^{n-1}) \quad (12)$$

or

$$Mq^n - \Delta t A q^n = Mq^{n-1} + \gamma \Delta t f(Q^{n-1}, R^{n-1}) \quad (13)$$

$$Mr^n - D \Delta t A r^n = Mr^{n-1} + \gamma \Delta t g(Q^{n-1}, R^{n-1}) \quad (14)$$