Class 18: Pertussis Mini Project

Aiden Quisneberry (PID:A19083392)

Table of contents

Background 1

CMI-PB Project **8**

Focus on "PT" Pertussis Toxin antigen 17

Background

Pertussis (a.k.a, Whooping Cough) is a common lung infection caused by the bacteria Bordatella Pertussis (a.k.a, B. Pertussis)

This disease mostly effects only humans and can infect all ages, the symptoms and long-term effects however are most severe for ages under 1 year and can lead to chronic illnes or even death.

The CDC track the number of reported cases in US: We can "scrape" this data with the **datapasta** package.

```
cdc <- data.frame(
  year = c(
    1922L,
    1923L,
    1924L,
    1925L,
    1926L,
    1927L,
    1928L,
    1929L,
    1930L,
    1931L,
    1932L,
```

```
    1933L,
    1934L,
    1935L,
    1936L,
    1937L,
    1938L,
    1939L,
    1940L,
    1941L,
    1942L,
    1943L,
    1944L,
    1945L,
    1946L,
    1947L,
    1948L,
    1949L,
    1950L,
    1951L,
    1952L,
    1953L,
    1954L,
    1955L,
    1956L,
    1957L,
    1958L,
    1959L,
    1960L,
    1961L,
    1962L,
    1963L,
    1964L,
    1965L,
    1966L,
    1967L,
    1968L,
    1969L,
    1970L,
    1971L,
    1972L,
    1973L,
    1974L,
```

```
    1975L,
    1976L,
    1977L,
    1978L,
    1979L,
    1980L,
    1981L,
    1982L,
    1983L,
    1984L,
    1985L,
    1986L,
    1987L,
    1988L,
    1989L,
    1990L,
    1991L,
    1992L,
    1993L,
    1994L,
    1995L,
    1996L,
    1997L,
    1998L,
    1999L,
    2000L,
    2001L,
    2002L,
    2003L,
    2004L,
    2005L,
    2006L,
    2007L,
    2008L,
    2009L,
    2010L,
    2011L,
    2012L,
    2013L,
    2014L,
    2015L,
    2016L,
```

```
    2017L,
    2018L,
    2019L,
    2020L,
    2021L,
    2022L,
    2023L,
    2024L,
    2025
  ), cases = c(
    107473,
    164191,
    165418,
    152003,
    202210,
    181411,
    161799,
    197371,
    166914,
    172559,
    215343,
    179135,
    265269,
    180518,
    147237,
    214652,
    227319,
    103188,
    183866,
    222202,
    191383,
    191890,
    109873,
    133792,
    109860,
    156517,
    74715,
    69479,
    120718,
    68687,
    45030,
    37129,
```

```
    60886,
    62786,
    31732,
    28295,
    32148,
    40005,
    14809,
    11468,
    17749,
    17135,
    13005,
    6799,
    7717,
    9718,
    4810,
    3285,
    4249,
    3036,
    3287,
    1759,
    2402,
    1738,
    1010,
    2177,
    2063,
    1623,
    1730,
    1248,
    1895,
    2463,
    2276,
    3589,
    4195,
    2823,
    3450,
    4157,
    4570,
    2719,
    4083,
    6586,
    4617,
    5137,
```

```
    7796,
    6564,
    7405,
    7298,
    7867,
    7580,
    9771,
    11647,
    25827,
    25616,
    15632,
    10454,
    13278,
    16858,
    27550,
    18719,
    48277,
    28639,
    32971,
    20762,
    17972,
    18975,
    15609,
    18617,
    6124,
    2116,
    3044,
    7063,
    22538,
    21996
  )
)
```

Q. Make a Plot of year vs cases

```
library(ggplot2)
ggplot(cdc, aes(x = year, y = cases)) +
  geom_point() +
  geom_line() +
  labs( title = "Pertussis Cases by Year", x = "Year", y = "No. of Cases") +
  theme_minimal() +
  scale_y_continuous(labels = scales::label_comma())
```



Q. Add some major milestones including the first wP vaccine roll-out (1946), the switch to the newer aP vaccine (1996), the COVID years (2020)

```
library(ggplot2)
ggplot(cdc, aes(x = year, y = cases)) +
  geom_point() +
  geom_line() +
  labs( title = "Pertussis Cases by Year in the US", x = "Year", y = "No. of Cases") +
  theme_minimal() +
  scale_y_continuous(labels = scales::label_comma()) +
  geom_vline(xintercept = c(1946, 1996, 2020), col=c("blue","red","yellow"), lty = 2 )
```



There were high case numbers in the pre 1940s, then in 1946 after the first wP vaccine was introduced the number of cases began to drastically reduce.

In 1996 the wP vaccine which required 1 shot was switched to an aP vaccine which required a booster every 10 years. Approximately 10 years later, there was a massive spike in cases.

In 2020, the worldwide COVID pandemic caused a large drop in pertussis cases. This is likely caused by the social distancing and health regulations imposed to address the worldwide pandemic. During this time vaccines got a lot of controversy publicly.

We understand that aP induced protection wanes faster than the wP induced protection. Pulling information from other countries that switched vaccines, they also saw a correlated rise in cases approximately 10 years after its switch.

Why is this vaccine-preventable disease on the upswing? To answer this question we need to investigate the mechanisms underlying waning protection against pertussis. This requires evaluation of pertussis-specific immune responses over time in wP and aP vaccinated individuals.

CMI-PB Project

Computation Models of Immunity - Pertussis Boost project aims to procide the scientific community with this very information.

They make their data available via JSON format APIs. We can read this is R with the `read_json()` function from the **jsonlite** package:

```
library(jsonlite)

subject <- read_json("http://cmi-pb.org/api/v5_1/subject", simplifyVector = TRUE)

head(subject)
```

```
  subject_id infancy_vac biological_sex              ethnicity  race
1          1          wP         Female Not Hispanic or Latino White
2          2          wP         Female Not Hispanic or Latino White
3          3          wP         Female                Unknown White
4          4          wP           Male Not Hispanic or Latino Asian
5          5          wP           Male Not Hispanic or Latino Asian
6          6          wP         Female Not Hispanic or Latino White
  year_of_birth date_of_boost      dataset
1    1986-01-01    2016-09-12 2020_dataset
2    1968-01-01    2019-01-28 2020_dataset
3    1983-01-01    2016-10-10 2020_dataset
4    1988-01-01    2016-08-29 2020_dataset
5    1991-01-01    2016-08-29 2020_dataset
6    1988-01-01    2016-10-10 2020_dataset
```

Q. How many wP and aP individuals are in this `subject` table

```
sum(subject$infancy_vac=="wP")
```

```
[1] 85
```

```
sum(subject$infancy_vac=="aP")
```

```
[1] 87
```

```
table(subject$infancy_vac)
```

```
aP wP
87 85
```

Q. What is the biological sex breakdown?

```r
table(subject$biological_sex)
```

```
Female   Male
   112     60
```

Q. In terms of race and gender, is this dataset representative of the US population?

```r
table(subject$race, subject$biological_sex)
```

```
                                           Female Male
  American Indian/Alaska Native                 0    1
  Asian                                        32   12
  Black or African American                     2    3
  More Than One Race                           15    4
  Native Hawaiian or Other Pacific Islander     1    1
  Unknown or Not Reported                      14    7
  White                                        48   32
```

Let's read some more database tables:

```r
specimen <- read_json("http://cmi-pb.org/api/v5_1/specimen", simplifyVector = TRUE)
ab_titer <- read_json("http://cmi-pb.org/api/v5_1/plasma_ab_titer", simplifyVector = TRUE)
```

```r
head(specimen)
```

```
  specimen_id subject_id actual_day_relative_to_boost
1           1          1                           -3
2           2          1                            1
3           3          1                            3
4           4          1                            7
5           5          1                           11
6           6          1                           32
  planned_day_relative_to_boost specimen_type visit
1                             0         Blood     1
2                             1         Blood     2
3                             3         Blood     3
4                             7         Blood     4
5                            14         Blood     5
6                            30         Blood     6
```

```r
head(ab_titer)
```

```
  specimen_id isotype is_antigen_specific antigen        MFI MFI_normalised
1           1     IgE               FALSE   Total 1110.21154       2.493425
2           1     IgE               FALSE   Total 2708.91616       2.493425
3           1     IgG                TRUE      PT   68.56614       3.736992
4           1     IgG                TRUE     PRN  332.12718       2.602350
5           1     IgG                TRUE     FHA 1887.12263      34.050956
6           1     IgE                TRUE     ACT    0.10000       1.000000
   unit lower_limit_of_detection
1 UG/ML                 2.096133
2 IU/ML                29.170000
3 IU/ML                 0.530000
4 IU/ML                 6.205949
5 IU/ML                 4.679535
6 IU/ML                 2.816431
```

To analyze this data we need to first "join" the different tables so we have all the data in one place and not spread across multiple tables.

We can use the `*_join` family of functions from **dplyr** to do this

```r
library(dplyr)
```

```
Attaching package: 'dplyr'

The following objects are masked from 'package:stats':

    filter, lag

The following objects are masked from 'package:base':

    intersect, setdiff, setequal, union
```

```r
meta <- inner_join(subject, specimen)
```

```
Joining with `by = join_by(subject_id)`
```

```r
head(meta)
```

```
  subject_id infancy_vac biological_sex              ethnicity  race
1          1          wP         Female Not Hispanic or Latino White
2          1          wP         Female Not Hispanic or Latino White
3          1          wP         Female Not Hispanic or Latino White
4          1          wP         Female Not Hispanic or Latino White
5          1          wP         Female Not Hispanic or Latino White
6          1          wP         Female Not Hispanic or Latino White
  year_of_birth date_of_boost      dataset specimen_id
1    1986-01-01    2016-09-12 2020_dataset           1
2    1986-01-01    2016-09-12 2020_dataset           2
3    1986-01-01    2016-09-12 2020_dataset           3
4    1986-01-01    2016-09-12 2020_dataset           4
5    1986-01-01    2016-09-12 2020_dataset           5
6    1986-01-01    2016-09-12 2020_dataset           6
  actual_day_relative_to_boost planned_day_relative_to_boost specimen_type
1                           -3                             0         Blood
2                            1                             1         Blood
3                            3                             3         Blood
4                            7                             7         Blood
5                           11                            14         Blood
6                           32                            30         Blood
  visit
1     1
2     2
3     3
4     4
5     5
6     6
```

```r
abdata <- inner_join(ab_titer, meta)
```

```
Joining with `by = join_by(specimen_id)`
```

```r
head(abdata)
```

```
  specimen_id isotype is_antigen_specific antigen        MFI MFI_normalised
1           1     IgE               FALSE   Total 1110.21154       2.493425
2           1     IgE               FALSE   Total 2708.91616       2.493425
```

```
3           1     IgG                TRUE      PT   68.56614       3.736992
4           1     IgG                TRUE     PRN  332.12718       2.602350
5           1     IgG                TRUE     FHA 1887.12263      34.050956
6           1     IgE                TRUE     ACT    0.10000       1.000000
   unit lower_limit_of_detection subject_id infancy_vac biological_sex
1 UG/ML                 2.096133          1          wP         Female
2 IU/ML                29.170000          1          wP         Female
3 IU/ML                 0.530000          1          wP         Female
4 IU/ML                 6.205949          1          wP         Female
5 IU/ML                 4.679535          1          wP         Female
6 IU/ML                 2.816431          1          wP         Female
               ethnicity  race year_of_birth date_of_boost      dataset
1 Not Hispanic or Latino White    1986-01-01    2016-09-12 2020_dataset
2 Not Hispanic or Latino White    1986-01-01    2016-09-12 2020_dataset
3 Not Hispanic or Latino White    1986-01-01    2016-09-12 2020_dataset
4 Not Hispanic or Latino White    1986-01-01    2016-09-12 2020_dataset
5 Not Hispanic or Latino White    1986-01-01    2016-09-12 2020_dataset
6 Not Hispanic or Latino White    1986-01-01    2016-09-12 2020_dataset
  actual_day_relative_to_boost planned_day_relative_to_boost specimen_type
1                           -3                             0         Blood
2                           -3                             0         Blood
3                           -3                             0         Blood
4                           -3                             0         Blood
5                           -3                             0         Blood
6                           -3                             0         Blood
  visit
1     1
2     1
3     1
4     1
5     1
6     1
```

Q. What Antibody isotypes are measured for these patrients?

```r
table(abdata$isotype)
```

```
  IgE   IgG  IgG1  IgG2  IgG3  IgG4 
 6698  7265 11993 12000 12000 12000 
```

Q. What antigens are reported?

```r
table(abdata$antigen)
```

```
    ACT   BETV1      DT   FELD1     FHA  FIM2/3   LOLP1     LOS Measles     OVA 
   1970    1970    6318    1970    6712    6318    1970    1970    1970    6318 
    PD1     PRN      PT     PTM   Total      TT 
   1970    6712    6712    1970     788    6318 
```

Let's focus on IgG isotype and make a plot of MFI_normalized for all antigens.

```r
igg  <- abdata |>
  filter(isotype == "IgG")
head(igg)
```

```
  specimen_id isotype is_antigen_specific antigen        MFI MFI_normalised
1           1     IgG                TRUE      PT   68.56614       3.736992
2           1     IgG                TRUE     PRN  332.12718       2.602350
3           1     IgG                TRUE     FHA 1887.12263      34.050956
4          19     IgG                TRUE      PT   20.11607       1.096366
5          19     IgG                TRUE     PRN  976.67419       7.652635
6          19     IgG                TRUE     FHA   60.76626       1.096457
   unit lower_limit_of_detection subject_id infancy_vac biological_sex
1 IU/ML                 0.530000          1          wP         Female
2 IU/ML                 6.205949          1          wP         Female
3 IU/ML                 4.679535          1          wP         Female
4 IU/ML                 0.530000          3          wP         Female
5 IU/ML                 6.205949          3          wP         Female
6 IU/ML                 4.679535          3          wP         Female
               ethnicity  race year_of_birth date_of_boost      dataset
1 Not Hispanic or Latino White    1986-01-01    2016-09-12 2020_dataset
2 Not Hispanic or Latino White    1986-01-01    2016-09-12 2020_dataset
3 Not Hispanic or Latino White    1986-01-01    2016-09-12 2020_dataset
4                Unknown White    1983-01-01    2016-10-10 2020_dataset
5                Unknown White    1983-01-01    2016-10-10 2020_dataset
6                Unknown White    1983-01-01    2016-10-10 2020_dataset
  actual_day_relative_to_boost planned_day_relative_to_boost specimen_type
1                           -3                             0         Blood
2                           -3                             0         Blood
3                           -3                             0         Blood
4                           -3                             0         Blood
5                           -3                             0         Blood
```

```
6                           -3                              0          Blood
  visit
1     1
2     1
3     1
4     1
5     1
6     1
```

```
ggplot(igg) +
  aes(MFI_normalised, antigen) +
  geom_boxplot()
```



Q. Is there a difference for aP vs wP individuals with these values?

```
ggplot(igg) +
  aes(MFI_normalised, antigen) +
  geom_boxplot() +
  facet_wrap(~infancy_vac)
```




```
ggplot(igg) +
  aes(MFI_normalised, antigen, col=infancy_vac) +
  geom_boxplot()
```



Q. Is there a temporal response - i.e do values increase or decrease over time?

```
ggplot(igg) +
  aes(MFI_normalised, antigen,col=infancy_vac) +
  geom_boxplot() +
  facet_wrap(~visit)
```



Focus on "PT" Pertussis Toxin antigen

```
pt.igg.21 <-  igg |> filter(antigen == "PT",
                            dataset =="2021_dataset")
```

```
ggplot(pt.igg.21) +
  aes(planned_day_relative_to_boost,
      MFI_normalised,
      col=infancy_vac,
      group = subject_id) +
  geom_point() +
  geom_line() +
  geom_vline(xintercept = 14, lty = 2)
```




```
ggplot(pt.igg.21) +
  aes(planned_day_relative_to_boost,
      MFI_normalised,
      col=infancy_vac,
      group = subject_id) +
  geom_point() +
  geom_line(alpha = .3) +
  geom_smooth(aes(group = infancy_vac),
              se = F,
              span = .4) +
  geom_vline(xintercept = 14, lty = 2)
```

```
`geom_smooth()` using method = 'loess' and formula = 'y ~ x'
```


MFI_normalised
15
10
5
0
0
25
50
75
100
125
planned_day_relative_to_boost
infancy_vac
aP
wP