Copyright Notice

These slides are distributed under the Creative Commons License.

DeepLearning.ai makes these slides available for educational purposes. You may not use or distribute these slides for commercial purposes. You may make copies of these slides and use or distribute them for educational purposes as long as you cite DeepLearning.AI as the source of the slides.

For the rest of the details of the license, see https://creativecommons.org/licenses/by-sa/2.0/legalcode

W3 Lesson 1

DeepLearning.AI

Vectors and Linear Transformations

Machine Learning motivation

Neural Networks - AI generated images

AI-generated human faces.

- Generative learning: Generating realistic looking images.

Text-to-image and image-to-text generation

DeepLearning.AI

Vectors and Linear Transformations

Vectors and their properties

Vectors

How to get from point A to point B?

How to get from point A to point B?

How to get from point A to point B?

Helicopter distance $= \sqrt{3^2 + 4^2} = 5$

Pythagorean Theorem

Norms

L1-norm $= \|(a,b)\|_1 = |a| + |b|$

L2-norm $= \|(a,b)\|_2 = \sqrt{a^2 + b^2}$

Norm of a vector

$$\sqrt{4^2 + 3^2} = \sqrt{25} = 5$$

Direction of a vector

$$\tan(\theta) = \frac{3}{4}$$

$$\theta = \arctan(3/4) = 0.64 \ = 36.87^\circ$$

DeepLearning.AI

Vectors and Linear Transformations

Sum and difference of vectors

Sum of vectors

$$u + v = (4 + 1, 1 + 3) = (5,4)$$

Difference of vectors

$$u - v = (4 - 1, 1 - 3) = (3, -2)$$

General definition: sum and difference

Same number of components

$$x = (x_1 \quad x_2 \quad \dots \quad x_n) \qquad y = (y_1 \quad y_2 \quad \dots \quad y_n)$$

Sum

$$x + y = (x_1 + y_1 \quad x_2 + y_2 \quad \dots \quad x_n + y_n)$$

Sum component by component

Difference

$$x - y = (x_1 - y_1 \quad x_2 - y_2 \quad \dots \quad x_n - y_n)$$

Subtract component by component

DeepLearning.AI

Vectors and Linear Transformations

Distance between vectors

Distances

L1-distance

$$\|u - v\|_1 = |5| + |-3| = 8$$

L2-distance

$$\|u - v\|_2 = \sqrt{5^2 + 3^2} = 5.83$$

Distances

L1-distance

$$\|u - v\|_1 = |5| + |-3| = 8$$

L2-distance

$$\|u - v\|_2 = \sqrt{5^2 + 3^2} = 5.83$$

Cosine distance

$$\cos(\theta)$$

DeepLearning.AI

Vectors and Linear Transformations

Multiplying a vector by a scalar

Multiplying a vector by a scalar

$$u = (1,2)$$

$$\lambda = 3$$

$$\lambda u = (3,6)$$

If the scalar is negative

$$u = (1,2)$$

$$\lambda = -2$$

$$\lambda u = (-2, -4)$$

General definition

Multiplication by a Scalar

$$x = (x_1 \quad x_2 \quad \dots \quad x_n)$$

$$\lambda x = \left(\lambda x_1 \quad \lambda x_2 \quad \dots \quad \lambda x_n\right)$$

Video 6: The dot product

DeepLearning.AI

Vectors and Linear Transformations

The dot product

A shortcut for linear operations

Quantities
2 apples
4 bananas
1 cherry

Prices
apples: $3
bananas: $5
cherries: $2

Total price
$28

$$6 + 20 + 2 = 28$$

The dot product

The dot product

$$(2 \times 3) + (4 \times 5) + (1 \times 2) = 28$$

The dot product

$$(2 \times 3) + (4 \times 5) + (1 \times 2) = 28$$

The dot product

$$\begin{bmatrix} 2 & 4 & 1 \end{bmatrix} \cdot \begin{bmatrix} 3 \\ 5 \\ 2 \end{bmatrix} = 28$$

$$(2 \times 3) + (4 \times 5) + (1 \times 2) = 28$$

Norm of a vector using dot product

$$\sqrt{4^2 + 3^2} = \sqrt{25} = 5$$

$$\begin{bmatrix} 4 & 3 \end{bmatrix} \begin{bmatrix} 4 \\ 3 \end{bmatrix} = 25$$

$$L2 - norm = \sqrt{dot\ \ product(u, u)}$$

$$\|u\|_2 = \sqrt{\langle u, u \rangle}$$

Vector Transpose

$$\begin{bmatrix} 2 \\ 4 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 3 \\ 5 \\ 2 \end{bmatrix} = 28$$

$$(2 \times 3) + (4 \times 5) + (1 \times 2) = 28$$

Vector Transpose

Transpose: convert columns to rows

$$\begin{bmatrix} 2 \\ 4 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 3 \\ 5 \\ 2 \end{bmatrix} = 28$$

$$\begin{bmatrix} 2 & 4 & 1 \end{bmatrix} \cdot \begin{bmatrix} 3 \\ 5 \\ 2 \end{bmatrix} = 28$$

$$2 \cdot 3 + 4 \cdot 5 + 1 \cdot 2 = 28$$

$$2 \cdot 3 + 4 \cdot 5 + 1 \cdot 2 = 28$$

Vector Transpose

$$\begin{bmatrix} 2 \\ 4 \\ 1 \end{bmatrix}^T =$$

Vector Transpose

$$\begin{bmatrix} 2 \\ 4 \\ 1 \end{bmatrix}^T = \begin{bmatrix} 2 & 4 & 1 \end{bmatrix}$$

Vector Transpose

$$\begin{bmatrix} 2 \\ 4 \\ 1 \end{bmatrix}^T = \begin{bmatrix} 2 & 4 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 2 & 4 & 1 \end{bmatrix}^T =$$

Vector Transpose

$$\begin{bmatrix} 2 \\ 4 \\ 1 \end{bmatrix}^T = \begin{bmatrix} 2 & 4 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 2 & 4 & 1 \end{bmatrix}^T = \begin{bmatrix} 2 \\ 4 \\ 1 \end{bmatrix}$$

Matrix Transpose

$$\begin{bmatrix} 2 & 5 \\ 4 & 7 \\ 1 & 3 \end{bmatrix}^T =$$

Transpose the columns!

Matrix Transpose

$$\begin{bmatrix} 2 & 5 \\ 4 & 7 \\ 1 & 3 \end{bmatrix}^T = \begin{bmatrix} 2 & 4 & 1 \end{bmatrix}$$

Matrix Transpose

$$\begin{bmatrix} 2 & 5 \\ 4 & 7 \\ 1 & 3 \end{bmatrix}^T = \begin{bmatrix} 2 & 4 & 1 \\ 5 & 7 & 3 \end{bmatrix}$$

3×2 $\qquad$ 3×2

Columns → Rows

General definition: dot product

Same number of components

$$x = (x_1 \quad x_2 \quad \dots \quad x_n) \qquad\qquad y = (y_1 \quad y_2 \quad \dots \quad y_n)$$

$$x \cdot y = (x_1 \times y_1) + (x_2 \times y_2) + \dots + (x_n \times y_n)$$

$\langle x, y \rangle$ is another notation for the dot product

$$x \cdot y^T = (x_1 \quad x_2 \quad \dots \quad x_n) \cdot \begin{pmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{pmatrix}$$

DeepLearning.AI

Vectors and Linear Transformations

Geometric dot product

Orthogonal vectors have dot product 0

6	2	-1	=	0
		3		

$$\langle u, v \rangle = 0$$

The dot product

$$\langle u, u \rangle = \|u\|^2$$

$$\langle u, v \rangle = 0$$

$$\langle u, v \rangle = ?$$

The dot product

$$\langle u, u \rangle = \|u\|^2 = \|u\| \cdot \|u\|$$

$$\langle u, v \rangle = \|u\| \cdot \|v\|$$

$$\langle u, v \rangle = \|u'\| \cdot \|v\|$$

$$= \|u\| \|v\| \cos(\theta)$$

Geometric dot product

6	2	2 / 4	=	20	Positive
6	2	-1 / 3	=	0	
6	2	-4 / 1	=	-22	Negative

Geometric dot product

$$\langle u, v \rangle > 0$$

$$\langle u, v \rangle = 0$$

$$\langle u, v \rangle < 0$$

DeepLearning.AI

Vectors and Linear Transformations

Multiplying a matrix by a vector

Equations as dot product

$$2a + 4b + c = 28$$

Equations as dot product

$a + b + c = 10$

$a + 2b + c = 15$

$a + b + 2c = 12$

Equations as dot product

$a + b + c = 10$

$a + 2b + c = 15$

$a + b + 2c = 12$

Equations as dot product

$a + b + c = 10$

$a + 2b + c = 15$

$a + b + 2c = 12$

Equations as dot product

System of equations

$$a + b + c = 10$$

$$a + 2b + c = 15$$

$$a + b + 2c = 12$$

Matrix product

Equations as dot product

System of equations

$$a + b + c = 10$$

$$a + 2b + c = 15$$

$$a + b + 2c = 12$$

Matrix product

$$\begin{bmatrix} 1 & 1 & 1 \\ 1 & 2 & 1 \\ 1 & 1 & 2 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} 10 \\ 15 \\ 12 \end{bmatrix}$$

Equations as dot product

Matrix product

$$\begin{bmatrix} 1 & 1 & 1 \\ 1 & 2 & 1 \\ 1 & 1 & 2 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} 10 \\ 15 \\ 12 \end{bmatrix}$$

$3 \times \boxed{3}$ Length $\boxed{3}$

Matrix can be rectangular

columns = length of vector

Equations as dot product

Matrix product

$$\begin{bmatrix} 1 & 1 & 1 \\ 1 & 2 & 1 \\ 1 & 1 & 2 \\ 2 & 1 & 1 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} 10 \\ 15 \\ 12 \\ 13 \end{bmatrix}$$

$\boxed{4} \times \boxed{3}$ Length $\boxed{3}$ Length $\boxed{4}$

Matrix can be rectangular

columns = length of vector

W3 Lesson 2

DeepLearning.AI

Vectors and Linear Transformations

Matrices as linear transformations

Matrices as linear transformations

$$\begin{bmatrix} 3 & 1 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 4 \\ 3 \end{bmatrix}$$

$(0,0) \rightarrow (0,0)$

$(1,0) \rightarrow (3,1)$

$(0,1) \rightarrow (1,2)$

$(1,1) \rightarrow (4,3)$

Matrices as linear transformations

3	1	-2
1	2	3

=

-3
4

Systems of equations as linear transformations

$$\begin{bmatrix} 3 & 1 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 4 \\ 3 \end{bmatrix}$$

First day: $3a + b$
Second day: $a + 2b$

(1,1) → (4,3)

DeepLearning.AI

Vectors and Linear Transformations

Linear transformations as matrices

Linear transformations as matrices

?	?
?	?

(0,0) → (0,0)
(1,0) → (3,-1)
(0,1) → (2,3)
(1,1) → (5,2)

Linear transformations as matrices

$$\begin{bmatrix} ? & ? & 0 \\ ? & ? & 1 \end{bmatrix} = \begin{bmatrix} 2 \\ 3 \end{bmatrix}$$

$$\begin{bmatrix} ? & ? & 1 \\ ? & ? & 0 \end{bmatrix} = \begin{bmatrix} 3 \\ -1 \end{bmatrix}$$

(1,0) → (3,-1)

(0,1) → (2,3)

DeepLearning.AI

Vectors and Linear Transformations

Matrix multiplication

Combining linear transformations

$$\begin{bmatrix} 3 & 1 & 1 \\ 1 & 2 & 0 \end{bmatrix} = \begin{bmatrix} 3 \\ 1 \end{bmatrix}$$

$$\begin{bmatrix} 3 & 1 & 0 \\ 1 & 2 & 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$$

Combining linear transformations

$$\begin{bmatrix} 3 & 1 & 1 \\ 1 & 2 & 0 \end{bmatrix} = \begin{bmatrix} 3 \\ 1 \end{bmatrix}$$

$$\begin{bmatrix} 3 & 1 & 0 \\ 1 & 2 & 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$$

Combining linear transformations

$$\begin{bmatrix} 2 & -1 & 3 \\ 0 & 2 & 1 \end{bmatrix} = \begin{bmatrix} 5 \\ 2 \end{bmatrix}$$

$$\begin{bmatrix} 2 & -1 & 1 \\ 0 & 2 & 2 \end{bmatrix} = \begin{bmatrix} 0 \\ 4 \end{bmatrix}$$

Combining linear transformations

$$\begin{bmatrix} 2 & -1 & 3 \\ 0 & 2 & 1 \end{bmatrix} = \begin{bmatrix} 5 \\ 2 \end{bmatrix}$$

$$\begin{bmatrix} 2 & -1 & 1 \\ 0 & 2 & 2 \end{bmatrix} = \begin{bmatrix} 0 \\ 4 \end{bmatrix}$$

Combining linear transformations

Combining linear transformations

Combining linear transformations

Second First

$$\begin{bmatrix} 2 & -1 \\ 0 & 2 \end{bmatrix} \cdot \begin{bmatrix} 3 & 1 \\ 1 & 2 \end{bmatrix} = \begin{bmatrix} 5 & 0 \\ 2 & 4 \end{bmatrix}$$

Dimension of the matrices

Dimension of the matrices

$$\begin{bmatrix} 2 & -1 \\ 0 & 2 \end{bmatrix} \cdot \begin{bmatrix} 3 & 1 \\ 1 & 2 \end{bmatrix} = \begin{bmatrix} 5 & 0 \\ 2 & 4 \end{bmatrix}$$

2×2 $\quad$ 2×2 $\quad$ 2×2

Dimension of the matrices

$$\begin{bmatrix} 3 & 1 & 4 \\ 2 & -1 & 2 \end{bmatrix} \cdot \begin{bmatrix} 3 & 0 & 1 & -2 \\ 1 & 5 & 2 & 0 \\ -2 & 1 & 4 & 0 \end{bmatrix} = \begin{bmatrix} & & & \\ ? & & & \end{bmatrix}$$

2×3 $\quad$ 3×4 $\quad$ 2×4

Dimension of the matrices

3	1	4
2	-1	2

2×3

.

3	0	1	-2
1	5	2	0
-2	1	4	0

3×4

=

2	9	21	-6
?	-3	8	-4

2×4

- Columns of first matrix must match rows of second (numbers in red match)
- Result takes number of rows from first matrix (numbers in blue match)
- Result takes number of columns from second matrix (numbers in purple match)

DeepLearning.AI

Vectors and Linear Transformations

The identity matrix

The identity matrix

The identity matrix

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x \\ y \end{bmatrix}$$

(0,0) → (0,0)

(1,0) → (1,0)

(0,1) → (0,1)

(1,1) → (1,1)

DeepLearning.AI

Vectors and Linear Transformations

Matrix inverse

Matrix inverses

Multiplying matrices

$$\begin{bmatrix} 3 & 1 \\ 1 & 2 \end{bmatrix} \cdot \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 3 & 1 \\ 1 & 2 \end{bmatrix}^{-1} = \begin{bmatrix} 2/5 & -1/5 \\ -1/5 & 3/5 \end{bmatrix}$$

How to find an inverse?

$$\begin{bmatrix} 3 & 1 \\ 1 & 2 \end{bmatrix} \cdot \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 3 & 1 \end{bmatrix} \begin{bmatrix} a \\ c \end{bmatrix} = \begin{bmatrix} 1 \end{bmatrix} \qquad 3a + 1c = 1 \qquad a = \frac{2}{5}$$

$$\begin{bmatrix} 3 & 1 \end{bmatrix} \begin{bmatrix} b \\ d \end{bmatrix} = \begin{bmatrix} 0 \end{bmatrix} \qquad 3b + 1d = 0 \qquad b = -\frac{1}{5}$$

$$\begin{bmatrix} 1 & 2 \end{bmatrix} \begin{bmatrix} a \\ c \end{bmatrix} = \begin{bmatrix} 0 \end{bmatrix} \qquad 1a + 2c = 0 \qquad c = -\frac{1}{5}$$

$$\begin{bmatrix} 1 & 2 \end{bmatrix} \begin{bmatrix} b \\ d \end{bmatrix} = \begin{bmatrix} 1 \end{bmatrix} \qquad 1b + 2d = 1 \qquad d = \frac{3}{5}$$

Quiz

- Find the inverse of the following matrix. If you find that the task is impossible, feel free to click on “I couldn’t find it”

5	2
1	2

Solution

- By solving the corresponding system of linear equations, we get the following.

$$\begin{bmatrix} 5 & 2 \\ 1 & 2 \end{bmatrix} \cdot \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 5 & 2 \end{bmatrix} \begin{bmatrix} a \\ c \end{bmatrix} = \begin{bmatrix} 1 \end{bmatrix}$$

$$\begin{bmatrix} 5 & 2 \end{bmatrix} \begin{bmatrix} b \\ d \end{bmatrix} = \begin{bmatrix} 0 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 2 \end{bmatrix} \begin{bmatrix} a \\ c \end{bmatrix} = \begin{bmatrix} 0 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 2 \end{bmatrix} \begin{bmatrix} b \\ d \end{bmatrix} = \begin{bmatrix} 1 \end{bmatrix}$$

- **$5a + 2c = 1$**
- **$5b + 2d = 0$**
- **$a + 2c = 0$**
- **$b + 2d = 1$**

- **$a = 1/4$**
- **$b = -1/4$**
- **$c = -1/8$**
- **$d = 5/8$**

Quiz

- Find the inverse of the following matrix. If you find that the task is impossible, feel free to click on “I’m reaching a dead end”

1	1
2	2

Solutions

- The inverse doesn't exist!

We need to solve the following system of linear equations:

$$\begin{bmatrix} 1 & 1 \\ 2 & 2 \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$a + c = 1$
$2b + 2d = 1$
$2a + 2c = 0$
$b + d = 0$
This is clearly a contradiction, since equation 1 says $a+c=1$, and equation 3 says $2a+2c=0$.

DeepLearning.AI

Vectors and Linear Transformations

Which matrices have an inverse?

Which matrices have inverses?

$$5^{-1} = 0.2$$

$$\begin{bmatrix} 3 & 1 \\ 1 & 2 \end{bmatrix}^{-1} = \begin{bmatrix} 0.4 & -0.2 \\ -0.2 & 0.6 \end{bmatrix}$$

Non-singular matrix
Invertible

Det = 5

$$8^{-1} = 0.125$$

$$\begin{bmatrix} 5 & 2 \\ 1 & 2 \end{bmatrix}^{-1} = \begin{bmatrix} 0.25 & -0.25 \\ -0.125 & 0.625 \end{bmatrix}$$

Non-singular matrix
Invertible

Det = 8

Non-zero determinants

$$0^{-1} = ???$$

$$\begin{bmatrix} 1 & 1 \\ 2 & 2 \end{bmatrix} = \begin{bmatrix} ? & ? \\ ? & ? \end{bmatrix}$$

Singular matrix
Non-invertible

Det = 0

Zero determinant

DeepLearning.AI

Vectors and Linear Transformations

Neural networks and matrices

Quiz: Natural language processing

Spam	Lottery	Win
Yes	1	1
Yes	2	1
No	0	0
Yes	0	2
No	0	1
No	1	0
Yes	2	2
Yes	2	0
Yes	1	2

Scores:

Lottery: ___ points

Win: ___ points

Examples

Lottery: 3 point
Win: 2 points
“Win, win the lottery!” : 7points

Rule:
If the number of points of the sentence is bigger than ____, then the email is spam.

Goal: Find the best points and threshold
Lottery: ___ point
Win: ___ point
Threshold: ___ points

Quiz: Natural language processing

Spam	Lottery	Win
Yes	1	1
Yes	2	1
No	0	0
Yes	0	2
No	0	1
No	1	0
Yes	2	2
Yes	2	0
Yes	1	2

Score	> 1.5?
2	Yes
3	Yes
0	No
2	Yes
1	No
1	No
4	Yes
2	Yes
3	Yes

Solution:
Lottery: 1 point
Win: 1 point
Threshold: 1.5 points

Graphical natural language processing

Spam	Lottery	Win
Yes	1	1
Yes	2	1
No	0	0
Yes	0	2
No	0	1
No	1	0
Yes	2	2
Yes	2	0
Yes	1	2

Line:
$1 \cdot \text{Win} + 1 \cdot \text{Lottery} = 1.5$

Positive zone:
$1 \cdot \text{Win} + 1 \cdot \text{Lottery} > 1.5$

Negative zone:
$1 \cdot \text{Win} + 1 \cdot \text{Lottery} < 1.5$

Graphical natural language processing

Spam	Lottery	Win
Yes	1	1
Yes	2	1
No	0	0
Yes	0	2
No	0	1
No	1	0
Yes	2	2
Yes	2	0
Yes	1	2

Model
1
1

= 3

Check: > 1.5?

Spam

Dot product between vectors

Spam	Lottery	Win
Yes	1	1
Yes	2	1
No	0	0
Yes	0	2
No	0	1
No	1	0
Yes	2	2
Yes	2	0
Yes	1	2

Model
1
1

= 1

Check: > 1.5?

✗

Not spam

Matrix multiplication

Spam	Lottery	Win
Yes	1	1
Yes	2	1
No	0	0
Yes	0	2
No	0	1
No	1	0
Yes	2	2
Yes	2	0
Yes	1	2

Model
1
1

=

Prod
2
3
0
2
1
1
4
2
3

Check: >1.5?

Check
Yes
Yes
No
Yes
No
No
Yes
Yes
Yes

Perceptrons

Spam	Word1	Word2	…	WordN
Yes				
Yes				
No				
Yes				
No				
No				
Yes				
Yes				
Yes				

Model
…

=

Prod

Check:

Check
Yes
Yes
No
Yes
No
No
Yes
Yes
Yes

Threshold and bias

Spam	Lottery	Win	Bias
Yes	1	1	1
Yes	2	1	1
No	0	0	1
Yes	0	2	1
No	0	1	1
No	1	0	1
Yes	2	2	1
Yes	2	0	1
Yes	1	2	1

Check

$$1 \cdot \text{Win} + 1 \cdot \text{Lottery} - 1.5 > 0$$

$$1 \cdot \text{Win} + 1 \cdot \text{Lottery} - 1.5 > 0$$

Threshold

Bias

Check: > 0?

Model
1
1
-1.5

Bias

The AND operator

AND	x	y
No	0	0
No	1	0
No	0	1
Yes	1	1

Model
1
1

=

Dot prod
0
1
1
2

Check: >1.5?

Check
No
No
No
Yes

The AND operator

AND	x	y
No	0	0
No	1	0
No	0	1
Yes	1	1

$$1 \cdot x + 1 \cdot y = 1.5$$

The perceptron

x_1
w_1
x_2
w_2
b
$z = Wx + b$
U
1
blue
0
red
2
1.5
1
0.5
-5
0
5
10

DeepLearning.AI

Vectors and Linear Transformations

Conclusion