


Conic Sections and Eccentricity



The parabola, circle, ellipse and hyperbola are called conic sections because they can all be formed as cross sections of a cone (or pair of cones) as indicated in the diagrams above.

Each is understood to be a set of points in a plane satisfying a certain condition.


The conditions are usually stated as follows:

Parabola - equidistant from a fixed point (the focus) and a fixed line (the directrix)

Circle - at a fixed distance (radius) from a fixed point (center)

Ellipse - the sum of the distances from two fixed points (foci) is constant

Hyperbola – the difference of the distances from two fixed points (foci) is constant.



Eccentricity provides us with a different perspective on these curves and unifies the definitions.

Let F be a fixed point and \mathcal{L} a fixed line. Let $d(P, \mathcal{L})$ denote the distance from point P to line \mathcal{L} , measured along a perpendicular drawn from P to \mathcal{L} .

Consider the points P in the plane for which $PF = e \cdot d(P, \mathcal{L})$.

$$e = 1 \Rightarrow \text{parabola} \quad / \quad 0 < e < 1 \Rightarrow \text{ellipse} \quad / \quad e > 1 \Rightarrow \text{hyperbola}$$

The Circle: As e approaches 1, the ellipse becomes more and more elongated; as it approaches zero it becomes more and more circular.

For the ellipse and hyperbola, the distance from the center to a vertex is referred to as a and the distance from the center to a focus as c . For the ellipse and hyperbola, the eccentricity e has a

value of $\frac{c}{a}$.