

**MASSACHUSETTS MATHEMATICS LEAGUE
CONTEST 4 - JANUARY 2013 SOLUTION KEY**

Team Round - continued

B) The first complex fraction simplifies to $\frac{1}{x + \frac{1}{x + \frac{1}{2}}} = \frac{1}{x + \frac{2}{2x+1}} = \frac{2x+1}{2x^2+x+2}$.

The second complex fraction simplifies to $\frac{1}{2 + \frac{1}{2 + \frac{1}{x}}} = \frac{1}{2 + \frac{x}{2x+1}} = \frac{2x+1}{5x+2}$.

Thus, we require that $\frac{5x+2}{2x^2+x+2} = -1$

$$\Leftrightarrow 2x^2 + x + 2 = -5x - 2$$

$$\Leftrightarrow 2x^2 + 6x + 4 = 2(x+1)(x+2) = 0$$

$$\Leftrightarrow x = \underline{-1, -2}$$

C) $\sin^2(t) + \cos^2(t) = 1 \Rightarrow 1 + \cot^2(t) = \csc^2(t)$

$x^2 = 25\cot^2(t)$ and $y^2 = 9\csc^2(t)$ Subtracting, $\frac{x^2}{25} - \frac{y^2}{9} = \cot^2 t - \csc^2 t = -1$

Thus, $9x^2 - 25y^2 = -225$ or $9x^2 - 25y^2 + 225 = 0 \Rightarrow (A, C, D, E, F) = \underline{(9, -25, 0, 0, 225)}$

D) $(2x-3)(Bx-1) = 5 \Leftrightarrow 2Bx^2 - (2+3B)x - 2 = 0$

To insure exactly one root, we set the discriminant equal to zero.

$$b^2 - 4ac = (-(2+3B))^2 - 4(2B)(-2) = 0 \Rightarrow 9B^2 + 28B + 4 = 0$$

$$\Rightarrow B = \frac{-28 \pm \sqrt{28^2 - 4(36)}}{18} = \frac{-28 \pm \sqrt{4^2(7^2 - 9)}}{18} = \frac{-28 \pm 8\sqrt{10}}{18} = \frac{-14 \pm 4\sqrt{10}}{9}$$

The larger of the two values is $\frac{-14 + 4\sqrt{10}}{9}$. Substituting, 3 for $\sqrt{10}$, we have

$$\frac{-14 + 4(3)}{9} = \frac{-2}{9} \Rightarrow (P, Q) = \underline{(-2, 9)}.$$

The original Team D) question was simply “Approximate the larger of these two values to the nearest hundredth.” How could you tackle this question without a calculator?

One possibility is outlined at the end of this solution key and is followed by a discussion of an algorithm for computation of square root without a calculator.