

# Monte Carlo Simulation of Spins

---

Aiichiro Nakano

*Collaboratory for Advanced Computing & Simulations*

*Department of Computer Science*


*Department of Physics & Astronomy*

*Department of Chemical Engineering & Materials Science*

*Department of Quantitative & Computational Biology*

*University of Southern California*

Email: [anakano@usc.edu](mailto:anakano@usc.edu)



# Markov Chains for Complex Dynamics

- Hidden Markov model—Viterbi algorithm
- Perron-Frobenius eigenvalue cluster analysis

## DNA Sequencing via Quantum Mechanics and Machine Learning

*Int'l J. Comput. Sci.* **4**, 352 ('10)

Henry Yuen<sup>1</sup>, Fuyuki Shimojo<sup>1,2</sup>, Kevin J. Zhang<sup>1,3</sup>, Ken-ichi Nomura<sup>1</sup>, Rajiv K. Kalia<sup>1</sup>, Aiichiro Nakano<sup>1\*</sup>, Priya Vashishta<sup>1</sup>



## A Tutorial on Hidden Markov Models and Selected Applications in Speech Recognition

LAWRENCE R. RABINER, FELLOW, IEEE

*Proc. IEEE*, **77**, 257 ('89)

## Implementing the Viterbi Algorithm

*Fundamentals and real-time issues for processor designers*

HUI-LING LOU

*IEEE Signal Processing Mag.*, **12(5)**, 42 ('95)



ELSEVIER

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)




Linear Algebra and its Applications 398 (2005) 161–184

LINEAR ALGEBRA  
AND ITS  
APPLICATIONS

[www.elsevier.com/locate/laa](http://www.elsevier.com/locate/laa)

Robust Perron cluster analysis in conformation dynamics <sup>☆</sup>

Peter Deuflhard, Marcus Weber\*



USC **Viterbi**  
School of Engineering


UNIVERSITY OF SOUTHERN CALIFORNIA  
1880  
1920  
1945  
1950  
1960  
1970  
1980  
1990  
2000  
2010  
**USC**

# Ising Model


- Ising model: A collection of spins on a lattice,  $\{s_k \mid s_k = \pm 1\}$
- Potential energy

$$V(s^N) = -J \sum_{(k,l)} s_k s_l - H \sum_k s_k, \quad s_k = \pm 1$$

where  $J$  is the exchange coupling,  $H$  is the magnetic field, &  $(k, l)$  are nearest-neighbor pairs of lattice sites




Ernst Ising



- Periodic boundary condition: Wrapping around the lattice

Curie temperature



# Exotic Magnets


nature

Vol 451 | 3 January 2008 | doi:10.1038/nature06433

LETTERS

## Magnetic monopoles in spin ice

C. Castelnovo<sup>1</sup>, R. Moessner<sup>1,2</sup> & S. L. Sondhi<sup>3</sup>



$$H = \frac{J}{3} \sum_{\langle ij \rangle} S_i S_j + D a^3 \sum_{\langle ij \rangle} \left[ \frac{\hat{e}_i \cdot \hat{e}_j}{|\mathbf{r}_{ij}|^3} - \frac{3(\hat{e}_i \cdot \mathbf{r}_{ij})(\hat{e}_j \cdot \mathbf{r}_{ij})}{|\mathbf{r}_{ij}|^5} \right] S_i S_j$$



# 3D Ising Problem Is NP-Complete

PHYSICAL REVIEW VOLUME 65, NUMBERS 3 AND 4 FEBRUARY 1 AND 15, 1944

## Crystal Statistics. I. A Two-Dimensional Model with an Order-Disorder Transition

LARS ONSAGER

Sterling Chemistry Laboratory, Yale University, New Haven, Connecticut

(Received October 4, 1943)

The partition function of a two-dimensional "ferromagnetic" with scalar "spins" (Ising model) is computed rigorously for the case of vanishing field. The eigenwert problem involved in the corresponding computation for a long strip crystal of finite width ( $n$  atoms), joined straight to itself around a cylinder, is solved by direct product decomposition; in the special case  $n = \infty$  an integral replaces a sum. The choice of different interaction energies ( $\pm J, \pm J'$ ) in the  $(0\ 1)$  and  $(1\ 0)$  directions does not complicate the problem. The two-way infinite crystal has an order-disorder transition at a temperature  $T = T_c$  given by the condition

$$\sinh(2J/kT_c) \sinh(2J'/kT_c) = 1.$$

The energy is a continuous function of  $T$ ; but the specific heat becomes infinite as  $-\log |T - T_c|$ . For strips of finite width, the maximum of the specific heat increases linearly with  $\log n$ . The order-converting dual transformation invented by Kramers and Wannier effects a simple automorphism of the basis of the quaternion algebra which is natural to the problem in hand. In addition to the thermodynamic properties of the massive crystal, the free energy of a  $(0\ 1)$  boundary between areas of opposite order is computed; on this basis the mean ordered length of a strip crystal is

$$(\exp(2J/kT) \tanh(2J'/kT))^n.$$

## Statistical Mechanics, Three-Dimensionality and NP-completeness \*

### I. Universality of Intractability for the Partition Function of the Ising Model Across Non-Planar Lattices

[Extended Abstract]

Sorin Istrail

Sandia National Laboratories  
Applied Mathematics Department, MS 1110  
Albuquerque, NM 87185-1110  
[scistra@cs.sandia.gov](mailto:scistra@cs.sandia.gov)

This paper is authored by an employee(s) of the [U.S.] Government and is in the public domain.

STOC 2000 Portland Oregon USA  
1-58113-184-4/00/5

Nobel laureate Richard Feynman wrote in 1972 of the three-dimensional Ising model that "the exact solution for three dimensions has not yet been found."


Other researchers who have tried read like a roll call of famous names in science and mathematics: Onsager, Kac, Feynman, Fisher, Kasteleyn, Temperley, Green, Hurst, and more recently Barahona.

Says Istrail, "What these brilliant mathematicians and physicists failed to do, indeed cannot be done."

# Ising Model in Machine Learning

Physically very appealing methods for unsupervised learning are the so-called **Boltzmann machines** (BM). A BM is basically an *inverse Ising model* where the data samples are seen as samples from a Boltzmann distribution of a pairwise interacting Ising model. The goal is to learn the values of the interactions and magnetic fields so that the likelihood (probability in the Boltzmann measure) of the observed data is large.


Machine Learning and the Physical Sciences  
G. Carleo et al., *Rev. Mod. Phys.* **91**, 045002 ('19)



Quantum-neural nexus?

Quantum-annealing Boltzmann machine  
J. Liu et al., *Comput. Mater. Sci.* **173**, 109429 ('20)

Hybrid BM-quantum circuit  
G. Torlai et al., *Phys. Rev. Lett.* **123**, 230504 ('19)



# MC Algorithm for 2D Ising Model

```

initialize the spins, s[i][j] (0 ≤ i,j ≤ L-1)
Sum_A = 0
for step = 1 to maximum_step
    randomly select a grid point, (i,j)
    compute the change in potential energy, dV, with a single spin
    flip, si,j → -si,j
    if dV ≤ 0 accept the flip, si,j ← -si,j
    else if random() ≤ exp(-dV/kBT) then // 0 < random() < 1
        accept the flip, si,j ← -si,j
    endif
    Sum_A = Sum_A + A(sN) // Sample physical quantity A(sN)
endfor
Average_A = Sum_A/maximum_step

```

$$\begin{aligned}
 \delta V &= V(\dots, s_k', \dots) - V(\dots, s_k, \dots) & \frac{P(\dots, s_k', \dots)}{P(\dots, s_k, \dots)} &= \frac{e^{-V(\dots, s_k', \dots)/k_B T}}{e^{-V(\dots, s_k, \dots)/k_B T}} = \exp\left(-\frac{\overbrace{V(\dots, s_k', \dots) - V(\dots, s_k, \dots)}^{\delta V}}{k_B T}\right) \\
 &= -J \sum_{l \in n.n.(k)} (s_k' - s_k) s_l - H(s_k' - s_k) & i &= L * (\text{rand}() / (\text{double}) \text{RAND\_MAX}); \\
 &= -2s_k' \left( J \sum_{l \in n.n.(k)} s_l + H \right) & j &= L * (\text{rand}() / (\text{double}) \text{RAND\_MAX}); \\
 & & i &= \text{rand}() \% L; \\
 & & j &= \text{rand}() \% L;
 \end{aligned}$$

$V(s^N) = -J \sum_{(k,l)} s_k s_l - H \sum_k s_k, \quad s_k = \pm 1$

# State Transition

Spin flip:  $s_k \rightarrow s'_k = -s_k$


- double exp\_dV[2][5]:  $\exp(-\delta V / k_B T) = \exp\left(2s'_k \left(\frac{J}{k_B T} \sum_{l \in n.n.(k)} s_l + \frac{H}{k_B T}\right)\right)$

Central spin,  $s'_k$

-1


1

Neighbor spins				Sum
-1	-1	-1	-1	-4
-1	-1	-1	1	-2
-1	-1	1	1	0
-1	1	1	1	2
1	1	1	1	4




# Now, How to Code?

---



**Life of a computational  
physicist: Concept to  
algebra to program (=  
data structure—define &  
algorithm—pseudocode)**



# Data Structures

- `#define L 20 //Lattice size  
int s[L][L]; //Spins s[i][j] = ± 1`
- **Periodic boundary condition:** The west, east, south, north neighbors of site  $(i, j)$  are  $(im, j)$ ,  $(ip, j)$ ,  $(i, jm)$ ,  $(i, jp)$ , where  
 $im = (i + L - 1) \% L$   
 $ip = (i + 1) \% L$   
 $jm = (j + L - 1) \% L$   
 $jp = (j + 1) \% L$
- **Transition probability:** `double exp_dV[ 2 ][ 5 ]`

$$\exp_{-dV}[k][l] = \exp\left(-\frac{\delta V}{k_B T}\right) = \exp\left(2s\left[\frac{J}{k_B T} \sum_{s' \in n.n.(s)} s' + \frac{H}{k_B T}\right]\right)$$

where


$$k = (1 + s)/2 \quad (k = 0, 1; s = -1, 1)$$

$$l = (4 + S)/2 \quad (l = 0, 1, 2, 3, 4; S = \sum_{\text{neighbor}} s' = -4, -2, 0, 2, 4)$$

- **Ising model parameters**

$$\text{double JdivT} = J/k_B T$$

$$\text{double HdivT} = H/k_B T$$



$k=(1+s)/2$	$s$
0	-1
1	1

$l=(4+S)/2$	$S$
0	-4
1	-2
2	0
3	2
4	4

# Physical Quantities

---

- double runM;
  - > **Running value of magnetization**,  $M = \sum_k s_k$
  - > **To update**,  $M += 2s'_k$
- double sumM = 0.0, sumM2 = 0.0;
  - > **To calculate the mean & variance of the magnetization at the end**
  - > **After each MC step**

```
    sumM += runM;
    sumM2 += runM*runM;
```
- int Sta\_step;
  - > **# of MC steps to be performed**
  - > **Mean & standard deviation of the magnetization**



$$avgM = sumM/Sta\_step$$

$$sigM = \sqrt{\frac{sumM2}{Sta\_step} - avgM^2}$$

```
double avgM, sigM;
```


# Magnetization & Its Fluctuation

$L = 20$



Histogram for 2 million MC steps

# Energy & Its Fluctuation





**Potential energy:**  $V(s^N) = -J \sum_{(k,l)} s_k s_l - H \sum_k s_k, \quad s_k = \pm 1$

See notes on (1) [fluctuation-dissipation theorem](#), (2) [unitary time propagation](#) & (3) [linear response](#)

# Cluster MC Algorithms

- **Cluster MC:** Speed up the convergence of MC simulations by introducing collective motions of the degrees-of-freedom, e.g., flipping a cluster of spins at a time
- **Correlation time,  $\tau$ :** The number of MC steps before two states become uncorrelated
- **Dynamic critical exponent,  $z$ :** Near the critical temperature for magnetic-to-nonmagnetic phase transition,  $\tau \sim L^z$  ( $L$ : system size)
  - >  $z = 2.125$  for 2D Ising model
  - >  $z = 0.35$  for Swendsen-Wang cluster MC ('87)
  - >  $z \sim 0$  for multigrid (hierarchical cluster) MC (Kandel *et al.*, '88)



# Cluster MC Algorithm—Wolff

---

## Algorithm: Single step of Wolff's cluster flip

1. Choose a seed spin at random from the lattice.
2. Look in turn at each of the neighbors of that spin. If they are pointing in the same direction as the seed spin, add them to the cluster with probability  $P_{\text{add}} = 1 - \exp(-2J/k_B T)$ .
3. For each spin that was added in step 2, examine each of its neighbors to find the ones pointing in the same direction & add each of them to the cluster with probability  $P_{\text{add}}$ . Repeat this step as many times as necessary until there are no spins left in the cluster whose neighbors have not been considered for inclusion in the cluster.
4. Flip the cluster

U. Wolff, *Phys. Rev. Lett.* **62**, 361 ('89)

<https://aiichironakano.github.io/phys516/Phys516-MCred.pdf>


# Detailed Balance in Wolff Algorithm

- Key to detailed balance,  $\pi_{mn}\rho_n = \pi_{nm}\rho_m$ :  $P_{\text{add}} = 1 - \exp(-2J/k_B T)$
- Detailed balance for a cluster flip:  
 $\mu \rightarrow \nu$  ( **$m$  bonds broken**);  $\nu \rightarrow \mu$  ( **$n$  bonds broken**)

$$(1 - P_{\text{add}})^m P(\mu) = (1 - P_{\text{add}})^n P(\nu)$$

or

$$(1 - P_{\text{add}})^{m-n} = P(\nu)/P(\mu)$$




$$(1 - P_{\text{add}})^{m-n} = \exp(-2J[m-n]/k_B T) = \exp(-[E_\nu - E_\mu]/k_B T) = P(\nu)/P(\mu)$$

# Wolff Cluster MC Program

## Initiate a cluster-flip step using a stack

```
/* Put a random seed spin site onto a stack */
i = rand()%L; j = rand()%L;
stack[0] = i*L + j; // 1D site index put in a stack
sp = 1; // Stack pointer

/* Flip the seed and remember the old & new spins */
oldspin = s[i][j]; newspin = -s[i][j];
s[i][j] = newspin;
```



# Cluster Flip

```

while (sp) {
    /* Pop a site off the stack */
    current = stack[--sp]; i = current/L; j = current%L;

    /* Check the neighbors */
    if ((nn=i+1) >= L) nn -= L; /* South neighbor */
    if (s[nn][j] == oldspin)
        if (rand()/(double)RAND_MAX < padd) {
            stack[sp++] = nn*L + j; s[nn][j] = newspin;
        }

    if ((nn=i-1) < 0) nn += L; /* North neighbor */
    if (s[nn][j] == oldspin)
        if (rand()/(double)RAND_MAX < padd) {
            stack[sp++] = nn*L + j; s[nn][j] = newspin;
        }

    if ((nn=j+1) >= L) nn -= L; /* East neighbor */
    if (s[i][nn] == oldspin)
        if (rand()/(double)RAND_MAX < padd) {
            stack[sp++] = i*L + nn; s[i][nn] = newspin;
        }


    if ((nn=j-1) < 0) nn += L; /* West neighbor */
    if (s[i][nn] == oldspin)
        if (rand()/(double)RAND_MAX < padd) {
            stack[sp++] = i*L + nn; s[i][nn] = newspin;
        }
} /* End while stack is not empty */

```

$$padd = 1 - \exp(-2J/k_B T)$$

	j			
i	0	1	2	3
0	0	1	2	3
1	4	5	6	7
2	8	9	10	11
3	12	13	14	15

stack



# Sample Run

