

class10 structural bioinformatics p1

amy (pid A16962111)

the PDB database

Here we examine the size and composition of the main database of biomolecular structures - the PDB.

Get a CSV file from the PDB database and read it into R.

```
pdbstats <- read.csv("Data Export Summary.csv", row.names=1)
pdbstats
```

	X.ray	EM	NMR	Multiple.methods	Neutron	Other
Protein (only)	161,663	12,592	12,337	200	74	32
Protein/Oligosaccharide	9,348	2,167	34	8	2	0
Protein/NA	8,404	3,924	286	7	0	0
Nucleic acid (only)	2,758	125	1,477	14	3	1
Other	164	9	33	0	0	0
Oligosaccharide (only)	11	0	6	1	0	4
Total						
Protein (only)	186,898					
Protein/Oligosaccharide	11,559					
Protein/NA	12,621					
Nucleic acid (only)	4,378					
Other	206					
Oligosaccharide (only)	22					

- **Q1:** What percentage of structures in the PDB are solved by X-Ray and Electron Microscopy.

My pdbstats dataframe has numbers with commas in them. This may cause us problems.

```
sum(as.numeric(pdbstats$X.ray, pdbstats$EM))
```

Warning: NAs introduced by coercion

[1] NA

We found a function called `gsub()` . I can use this to make a function that I can use for every column in the table.

```
commasum <- function(x) {  
  sum(as.numeric(gsub(",", "", x)))  
}  
  
totals <- apply(pdbstats, 2, commasum)  
totals
```

X.ray	EM	NMR	Multiple.methods
182348	18817	14173	230
Neutron	Other	Total	
79	37	215684	

```
round(totals/totals["Total"]*100, 2)
```

X.ray	EM	NMR	Multiple.methods
84.54	8.72	6.57	0.11
Neutron	Other	Total	
0.04	0.02	100.00	

85% were solved by x-ray and 9% by EM.

- **Q2:** What proportion of structures in the PDB are protein?

```
commasum(pdbstats["Protein (only)", "Total"])/totals["Total"]*100
```

```
Total  
86.65362
```

87%

- **Q.** What fraction of Uniprot structures are represented in the PDB?

0.09%

```
215684/249751891*100
```

[1] 0.08635931

visualizing protein structure

We will learn the basics of Mol* (mol-star) homepage: <https://molstar.org/viewer/>

We will play with PDB 1HSG:




Figure 1: HIV-Pr with a bound inhibitor

back to R and working with PDB structures

```
library(bio3d)
hiv <- read.pdb("1hsg")
```

Note: Accessing on-line PDB file

```
hiv
```

Call: `read.pdb(file = "1hsg")`



Figure 2: with ASP25 residues represented as ball-and-stick figures

Total Models#: 1

Total Atoms#: 1686, XYZs#: 5058 Chains#: 2 (values: A B)

Protein Atoms#: 1514 (residues/Calpha atoms#: 198)

Nucleic acid Atoms#: 0 (residues/phosphate atoms#: 0)

Non-protein/nucleic Atoms#: 172 (residues: 128)

Non-protein/nucleic resid values: [HOH (127), MK1 (1)]

Protein sequence:

PQITLWQRPLVTIKIGGQLKEALLDTGADDTVLEEMSLPGRWKPKMIGGIGGFIKVRQYD
 QILIEICGHKAIGTVLVGPTPVNIIGRNLLTQIGCTLNFPQITLWQRPLVTIKIGGQLKE
 ALLDTGADDTVLEEMSLPGRWKPKMIGGIGGFIKVRQYDQILIEICGHKAIGTVLVGPTP
 VNIIGRNLLTQIGCTLNF

+ attr: atom, xyz, seqres, helix, sheet,
 calpha, remark, call

```
head(hiv$atom)
```

	type	eleno	elety	alt	resid	chain	resno	insert	x	y	z	o	b
1	ATOM	1	N	<NA>	PRO	A	1	<NA>	29.361	39.686	5.862	1	38.10
2	ATOM	2	CA	<NA>	PRO	A	1	<NA>	30.307	38.663	5.319	1	40.62
3	ATOM	3	C	<NA>	PRO	A	1	<NA>	29.760	38.071	4.022	1	42.64
4	ATOM	4	O	<NA>	PRO	A	1	<NA>	28.600	38.302	3.676	1	43.40
5	ATOM	5	CB	<NA>	PRO	A	1	<NA>	30.508	37.541	6.342	1	37.87
6	ATOM	6	CG	<NA>	PRO	A	1	<NA>	29.296	37.591	7.162	1	38.40

	segid	elesy	charge
1	<NA>	N	<NA>
2	<NA>	C	<NA>
3	<NA>	C	<NA>
4	<NA>	O	<NA>
5	<NA>	C	<NA>
6	<NA>	C	<NA>

```
pdbseq(hiv)
```

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
"P"	"Q"	"I"	"T"	"L"	"W"	"Q"	"R"	"P"	"L"	"V"	"T"	"I"	"K"	"I"	"G"	"G"	"Q"	"L"	"K"
21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40
"E"	"A"	"L"	"L"	"D"	"T"	"G"	"A"	"D"	"D"	"T"	"V"	"L"	"E"	"E"	"M"	"S"	"L"	"P"	"G"
41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60
"R"	"W"	"K"	"P"	"K"	"M"	"I"	"G"	"G"	"I"	"G"	"G"	"F"	"I"	"K"	"V"	"R"	"Q"	"Y"	"D"
61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80
"Q"	"I"	"L"	"I"	"E"	"I"	"C"	"G"	"H"	"K"	"A"	"I"	"G"	"T"	"V"	"L"	"V"	"G"	"P"	"T"
81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	1
"P"	"V"	"N"	"I"	"I"	"G"	"R"	"N"	"L"	"L"	"T"	"Q"	"I"	"G"	"C"	"T"	"L"	"N"	"F"	"P"
2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21
"Q"	"I"	"T"	"L"	"W"	"Q"	"R"	"P"	"L"	"V"	"T"	"I"	"K"	"I"	"G"	"G"	"Q"	"L"	"K"	"E"
22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41
"A"	"L"	"L"	"D"	"T"	"G"	"A"	"D"	"D"	"T"	"V"	"L"	"E"	"E"	"M"	"S"	"L"	"P"	"G"	"R"
42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61
"W"	"K"	"P"	"K"	"M"	"I"	"G"	"G"	"I"	"G"	"G"	"F"	"I"	"K"	"V"	"R"	"Q"	"Y"	"D"	"Q"
62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81
"I"	"L"	"I"	"E"	"I"	"C"	"G"	"H"	"K"	"A"	"I"	"G"	"T"	"V"	"L"	"V"	"G"	"P"	"T"	"P"
82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99		
"V"	"N"	"I"	"I"	"G"	"R"	"N"	"L"	"L"	"T"	"Q"	"I"	"G"	"C"	"T"	"L"	"N"	"F"		

here we will do a normal mode analysis (NMA) to predict functional motions of a kinase protein

```
adk <- read.pdb("6s36")
```

Note: Accessing on-line PDB file

PDB has ALT records, taking A only, rm.alt=TRUE

```
adk
```

```
Call: read.pdb(file = "6s36")
```

```
Total Models#: 1
```

```
Total Atoms#: 1898, XYZs#: 5694 Chains#: 1 (values: A)
```

```
Protein Atoms#: 1654 (residues/Calpha atoms#: 214)
```

```
Nucleic acid Atoms#: 0 (residues/phosphate atoms#: 0)
```

```
Non-protein/nucleic Atoms#: 244 (residues: 244)
```

```
Non-protein/nucleic resid values: [ CL (3), HOH (238), MG (2), NA (1) ]
```

```
Protein sequence:
```

```
MRIILLGAPGAGKGTQAQFIMEKYGIPQISTGDMRLRAAVKSGSELGKQAKDIMDAGKLV  
TDELVIALVKERIAQEDCRNGFLLDGFPRTPQADAMKEAGINVDYVLEFDVPDELIVDKI  
VGRRVHAPSGRVYHVKNPPKVEGKDDVTGEELTTRKDDQEETVRKRLVEYHQMTAPLIG  
YYSKEAEAGNTKYAKVDGTPVAEVRADLEKILG
```

```
+ attr: atom, xyz, seqres, helix, sheet,  
      calpha, remark, call
```

```
modes <- nma(adk)
```

```
Building Hessian... Done in 0.046 seconds.
```

```
Diagonalizing Hessian... Done in 0.526 seconds.
```

```
plot(modes)
```