

Precise asymptotics for subgroups of the symmetric group

The critical model, section capacity, and global reduction

Abstract

Let s_n be the number of actual subgroups of S_n , and let G_r count the subspaces of $\mathbb{F}_2^r$. We prove

$$s_n = n! G_{\lfloor n/2 \rfloor} [y^{\lfloor n/2 \rfloor}] (1 + y/6)^{n \bmod 2} \exp(y/2 + y^2/6 + y^4/384) (1 + O(2^{-cn}))$$

for a constant $c > 0$, common to both parities. The critical family includes all central lifts and the two odd-degree markers. The upper bound uses an annihilator-aware section capacity theorem, joint-source quotient moments, and complete weighted recurrences that establish boundedness before using it. We prove the necessary orbit-action exhaustion and give explicit finite certificates for its bounded cases. The split $c = 1$ case is a separate application and a check on the retained joint data. A saddle-point formula has relative error $O(n^{-1})$; an elementary expression has four residue-class constants and a first correction of order $n^{-1/4}$.

Contents

1	The counting problem and its normalization	3
1.1	Actual actions and recovered profiles	4
1.2	Algebraic inputs	5
2	The complete critical family	5
3	Exact terminal attachment	8
4	A complete small-support estimate	9
5	Annihilator-aware section capacity	11
5.1	Permutation-module bounds	11
5.2	A coupled socle estimate	13
5.3	The retained annihilator and the central cut	13
6	Same-source lifts and original-weight fusion	14
6.1	The binary permutation rank bound	14
6.2	The fractional Schur cost	15
6.3	A complete finite-menu estimate	16
7	Complete quotient moments and uniform transfer	19
8	Faithful character tuples on a common source	22
8.1	Whole-socle feasibility	22
8.2	Joint kernel counting	23

9	Relative heads of actual transitive actions	24
10	Complete chief-layer fibres on one source	30
11	Uniform capacity of affine local components	33
12	Explicit affine block counts	35
13	Nonaffine primitive components and compressed quotients	38
14	Complete primitive-action consumers	39
15	The bounded ternary block counts	43
15.1	Signed modules on a bounded number of triples	44
16	The bounded affine exceptions	48
17	The exceptional small pair actions	52
17.1	The finite inputs and what they certify	53
17.2	The twelve-pair actions	54
17.3	The sixteen-pair actions	57
18	Repeated natural heads and the smaller pair counts	63
18.1	The three exceptional five-point block counts	67
18.2	Three pairs and the complete degree-sixteen slice	69
19	The two zero-ternary frame exceptions	70
20	Exhaustion of the nonbinary orbit actions	73
21	Binary sections and joint character incidence	74
22	Reversible simultaneous quotient transport	77
23	The complete critical, cyclic-four, and carrier mixture	79
23.1	Three simultaneous parameters on a full projection	79
23.2	The finite rows and the recurrence	86
23.3	Adding every cyclic-four orbit	88
24	Complete binary counting	91
24.1	The original degree-thirty-two boundary	91
24.2	Uniformity over unbounded binary widths	92
24.3	Finite witness coverage and its universal interpretation	95
24.4	Small widths and the complete recurrence	97
25	Direct fusion for original binary orbits	98
25.1	An original cut-or-character alternative	98
25.2	The ordinary forward recurrence for binary orbits	100

26 A joint cyclic-four character moment	101
26.1 The retained normal columns	101
26.2 The abelian packet in the same tableau	102
26.3 Excessive quotients and bounded binary words	103
26.4 The weighted generation reduction	104
27 Markers, binary moments, and the one-triple audit	105
27.1 Exact natural- S_3 transport	105
27.2 A character moment on the complete binary error	105
27.3 The split one-triple incidence	106
28 The complete marker sum	107
29 The first split one-triple application	111
30 The complete macroscopic alternating packet	120
31 The small natural alternating packet	122
32 Exhaustive counting and boundedness-first assembly	123
33 Evaluation of the benchmark	125
33.1 The binary Gaussian sum	125
33.2 One positive saddle	125
33.3 An elementary four-periodic expression	126
A Finite evidence and reproduction	128
B The finite carrier reduction and its two rational cones	129

1 The counting problem and its normalization

We count subgroups as subsets of S_n , rather than conjugacy classes or abstract isomorphism classes. All unspecified logarithms in counting estimates are to base two; $\log_e$ denotes the natural logarithm. Roney-Dougal and Tracey [RDT25] prove

$$s_n \leq 2^{n^2/16+O(n^{3/2})}. \quad (1.1)$$

Together with the elementary binary construction, this identifies the quadratic exponential scale. The more precise benchmark considered here also records the labelled orbit decomposition.

Put $R = \lfloor n/2 \rfloor$ and $\varepsilon = n - 2R$, and define

$$P(y) = \frac{y}{2} + \frac{y^2}{6} + \frac{y^4}{384}, \quad c_R = [y^R]e^{P(y)}, \quad p_n = c_R + \frac{\varepsilon}{6}c_{R-1}, \quad L_n = n!G_R p_n. \quad (1.2)$$

Here $c_j = 0$ for $j < 0$, $G_R = \sum_k \begin{bmatrix} R \\ k \end{bmatrix}_2$, and $G_0 = 1$. In particular p_n is a coefficient, whereas the normalized number of subgroups in the critical family will be denoted C_n^{crit} . The identity $P'(y)e^{P(y)} = (e^{P(y)})'$ gives

$$Rc_R = \frac{1}{2}c_{R-1} + \frac{1}{3}c_{R-2} + \frac{1}{96}c_{R-4}. \quad (1.3)$$

All coefficients are positive. Useful crude consequences include $c_{R-1}/c_R \leq 2R$ and $c_{R-h}/c_R \leq (2R)^h$ for $0 \leq h \leq R$.

Theorem 1.1 (Critical-family asymptotic). *Consider the subgroups whose even orbits are the four actions in Table 1. In odd degree allow, in addition, either one fixed point or one natural S_3 orbit. Require full projection onto each displayed action, and include every subgroup with these orbit projections. Their number is $L_n(1+O(2^{-an}))$ for some $a > 0$. Consequently $s_n \geq L_n(1-O(2^{-an}))$.*

Theorem 1.2 (Section capacity). *Let $T \leq S_s$ be faithful, transitive, and not a 2-group, where $s \geq 24$ is even. Let A be any section of its binary permutation module, with action factoring through a specified quotient B of T . For a finite $\mathbb{F}_2 B$ -module M , put*

$$r(M) = \max_X \frac{m_X(\text{Soc } M)}{\dim_{\text{End}_B X} X}, \quad r(0) = 0,$$

where X runs over simple modules. There is a subspace $C \leq A^B$ such that

$$2 \dim C + 4r(A/C) \leq \frac{47s}{48}. \quad (1.4)$$

The statement concerns the actual quotient A/C , including all socle constituents exposed by the quotient.

The critical model and the capacity estimate are proved in Sections 2 and 5. The complete counting argument gives the following theorem.

Theorem 1.3 (Precise subgroup asymptotic). *There exist $c, C > 0$ and n_0 such that*

$$\left| \frac{s_n}{L_n} - 1 \right| \leq C 2^{-cn} \quad (n \geq n_0).$$

The constants are common to both parities. No boundedness assumption on s_n/L_n is needed.

We first prove complete counting estimates on one complementary source, with the normalizer of each original permutation action retained. Sections 20 and 24 give the exhaustive nonbinary action reduction and the independent binary recurrence. The natural C_3/A_4 packets and exact marker collapse complete the disjoint partition in Section 32. The binary recurrence is transported into the ambient ordinary row, and one boundedness-first induction then proves Theorem 1.3. The split $c = 1$ calculation in Section 29 is a required application of the joint-source method, with its shared-character and inverse-complement controls retained. Section 33 evaluates L_n and gives the saddle-point and elementary forms of the theorem.

1.1 Actual actions and recovered profiles

For a transitive action $U \leq S_w$ write $a(U) = |N_{S_w}(U)|$. If a profile contains m_U copies of each action type, then the number of labelled orbit data is

$$\frac{n!}{\prod_U a(U)^{m_U} m_U!}. \quad (1.5)$$

Indeed, partition the points into the prescribed unordered blocks and choose a conjugate of U on each block. The factors $w!$ cancel. A subgroup full on each block recovers both its actual orbits and their permutation actions, so there is no further overcount.

For later use, let $b = n - w$. Pointing an original orbit of type U has placement factor $n!/(b!a(U))$. If its complete complement is $J \leq S_b$, a fibre bound $z_U(J)$ therefore gives the normalized bound

$$\frac{1}{b!a(U)p_n G_R} \sum_{J \leq S_b} z_U(J). \quad (1.6)$$

In particular a uniform fibre envelope $z_U(b)$ produces the coefficient

$$\Delta(n, b) \frac{z_U(b)}{a(U)}, \quad \Delta(n, b) = \frac{p_b G_{\lfloor b/2 \rfloor}}{p_n G_R} \quad (1.7)$$

on the complete normalized count s_b/L_b . An auxiliary quotient cover does not change $a(U)$.

1.2 Algebraic inputs

We use Gaussian coefficients, Goursat's lemma, Burnside's basis theorem, the inflation–restriction sequence, the degree-two Künneth formula, and classification of group extensions by H^2 . For permutation groups we use the abelian-section bound of Kovács and Praeger [KP89]: an elementary abelian p -section of a subgroup of S_b has rank at most b/p . Its form for p -groups is also recorded as [RDT25, Theorem 2.8]. The module-generator estimates used in Section 5 come from [Tra18, Definition 1.5, Theorem 1.6 and Corollary 3.12]. The finite algebraic constants and the distinction between literal witness verification and exhaustive coverage are specified in Appendix A.

2 The complete critical family

The four even actions used in the model are listed below. The indicated kernel is trivial in the first two cases and is the centre in the last two. The resulting quotient is elementary abelian of the indicated rank. The extraspecial action is the faithful transitive action of plus type on eight points, denoted E_8 . The normalizer and quadratic-form calculations are literal finite calculations;

Action	Degree	Quotient rank	$a(U)$
C_2	2	1	2
regular V_4	4	2	24
natural D_8	4	2	8
$E_8 \cong 2_+^{1+4}$	8	4	384

Table 1: The critical even actions.

Appendix A describes their witnesses. The theorem about this family does not require the additional assertion that these exhaust equality in a permutation-rank bound.

For a fixed word D in these actions, let K be the product of the specified kernels and put $V = D/K = \mathbb{F}_2^R$. If $t = \dim K$, the squaring map is a vector-valued quadratic map $q : V \rightarrow K$. Its polarization is the commutator map. Write

$$\phi = \prod_{i \geq 1} (1 - 2^{-i}) > 0.$$

The Gaussian product formula implies

$$2^{k(R-k)} \leq \begin{bmatrix} R \\ k \end{bmatrix}_2 \leq \phi^{-1} 2^{k(R-k)}, \quad G_R \asymp 2^{\lfloor R^2/4 \rfloor}. \quad (2.1)$$

All constants in this section are absolute.

Lemma 2.1 (The canonical subgroups). *Uniformly in the critical word, the number of full-projecting subspaces $U \leq V$ is $G_R(1 + O(R2^{-R/2}))$.*

Proof. Failure to project onto a quotient coordinate of dimension at most four places U in one of its at most 15 pullback hyperplanes. There are at most R coordinates. Each hyperplane contains G_{R-1} subspaces, and (2.1) gives $G_{R-1}/G_R = O(2^{-R/2})$. The full preimage of a full-projecting U contains K . Its projection to each factor therefore contains the specified kernel and surjects onto the quotient, so equals that factor. Different U give different full preimages. $\square$

Lemma 2.2 (All lifts of a subspace). *Let $U \leq V$ have dimension k , and set $Q(U) = \langle q(u) : u \in U \rangle \leq K$. Put $\mathcal{A}(U) = Q(U)^\perp \leq K^*$. The number of subgroups of D whose image is exactly U is*

$$\sum_{B \leq \mathcal{A}(U)} 2^{k \dim B} = \sum_{j=0}^{\dim \mathcal{A}(U)} \begin{bmatrix} \dim \mathcal{A}(U) \\ j \end{bmatrix}_2 2^{kj}. \quad (2.2)$$

If U is full-projecting, all these subgroups project onto the factors.

Proof. For such a subgroup H , put $W = H \cap K$. Necessarily $q(U) \subseteq W$. Conversely every element of $\pi^{-1}(U)/W$ then has square one. Such a group is abelian, since $(xy)^{-1} = y^{-1}x^{-1}$ and every element is its own inverse; hence the quotient is elementary abelian. The subgroups with intersection W are the complements to K/W in this elementary abelian extension. They form a torsor under $\text{Hom}(U, K/W)$ and number $2^{k \dim_K W}$. Annihilator duality $B = W^\perp$ gives (2.2). Fullness follows again from Burnside's basis theorem on each factor. $\square$

Lemma 2.3 (Quadratic realization). *Fix a quadratic form Q on a binary vector space. If q_0 is a nonsingular plus-type form of dimension $h \in \{2, 4\}$, the surjections f satisfying $q_0 \circ f = Q$ form either the empty set or a torsor for $O^+(h, 2)$. In particular there are at most 72 such maps.*

Proof. If a map exists, the radical of the polar form of Q is exactly $\ker f$, since the target polar form is nonsingular and f is onto. Thus every such map has the same kernel and induces an isometry from the same quotient to the target. Any two differ by a unique target isometry. The elementary orders are $|O^+(2, 2)| = 2$ and $|O^+(4, 2)| = 72$. $\square$

Proposition 2.4 (Uniform exceptional-lift estimate). *Set $d = R/2 - t \geq 0$. For the fixed critical word, the total number of noncanonical full subgroups is $O(G_R 2^{-d})$.*

Proof. Fix an ℓ -space $B \leq K^*$, where $\ell \geq 1$. The restrictions to B of the nonabelian coordinate functionals span B^* ; choose ℓ of them as a basis. These are the pivot coordinates. Count ordered maps $\mathbb{F}_2^k \rightarrow V$ which are injective and full on each coordinate and which satisfy $B \leq \mathcal{A}(U)$. Relations dual to the chosen basis isolate the pivot coordinates. Thus fixing the nonpivot maps fixes every pivot quadratic form jointly. Lemma 2.3 allows at most 72 choices for its surjective coordinate map. Each pivot has target dimension at least two. Thus the ordered-map count is at most $72^\ell 2^{(R-2\ell)k}$.

Division by $|\text{GL}(k, 2)|$, compared with the unrestricted ordered injection count at least $\phi 2^{Rk}$, gives

$$\#\{U : \dim U = k, B \leq \mathcal{A}(U), U \text{ full}\} \leq \phi^{-1} \begin{bmatrix} R \\ k \end{bmatrix}_2 72^\ell 2^{-2\ell k}.$$

There are at most $\phi^{-1}2^{\ell(t-\ell)}$ choices for B . Multiplying by its lift weight $2^{k\ell}$ in (2.2), the exceptional mass at dimension k is at most

$$\phi^{-2} \begin{bmatrix} R \\ k \end{bmatrix}_2 \sum_{\ell \geq 1} 72^\ell 2^{-\ell(k-t+\ell)}. \quad (2.3)$$

Put $x = k - R/2$. Dividing by G_R introduces a factor at most $C2^{-x^2}$ by (2.1). The exponent depending on x and ℓ becomes

$$-x^2 - \ell x - \ell d - \ell^2 = -(x + \ell/2)^2 - \ell d - 3\ell^2/4.$$

The Gaussian sum over either lattice is uniformly bounded. Also,

$$\sum_{\ell \geq 1} 72^\ell 2^{-3\ell^2/4} < \infty.$$

Since $d \geq 0$, the result is $O(2^{-d})$. □

Proof of Theorem 1.1. Summing (1.5) over critical profiles of rank R gives $n!c_R$ in even degree, because the two rank-two actions have combined weight $1/24 + 1/8 = 1/6$. The canonical count therefore supplies the main term, with the uniform error of Lemma 2.1.

For multiplicities (a, v, b, e) of the four actions,

$$R = a + 2v + 2b + 4e, \quad t = b + e, \quad d = a/2 + v + e.$$

Inserting the weight 2^{-d} of Proposition 2.4 changes the exponential generating polynomial to

$$\tilde{P}(y) = \frac{y}{2\sqrt{2}} + \frac{7y^2}{48} + \frac{y^4}{768}. \quad (2.4)$$

Write $\tilde{c}_R = [y^R]e^{\tilde{P}(y)}$ and use the original positive saddle $\rho = \rho_R$ from Section 33. Positivity and the independent saddle estimate of Theorem 33.2 give

$$\frac{\tilde{c}_R}{c_R} \ll \sqrt{R} \exp(\tilde{P}(\rho) - P(\rho)) \ll \sqrt{R} e^{-R/16} = O(2^{-R/32}). \quad (2.5)$$

Indeed $\tilde{c}_R \rho^R \leq e^{\tilde{P}(\rho)}$, whereas $c_R \rho^R \gg e^{P(\rho)}/\sqrt{R}$. Also $\tilde{P}(\rho) \leq P(\rho) - \rho^4/768$ and $\rho^4 \geq 48R$ for all sufficiently large R , by the saddle equation. The final bound absorbs the square-root factor. No second saddle estimate is needed; the rate is not optimized.

In odd degree the fixed-point option contributes c_R . The natural S_3 option contributes $c_{R-1}/6$. Any subgroup full on its S_3 coordinate and on the binary coordinates contains A_3 on the first coordinate alone: raise an element projecting to a 3-cycle to a sufficiently large power of two. Quotienting this forced subgroup turns that coordinate into one elementary binary coordinate. Its full lift analysis is consequently the same one, with exceptional marker $y/(6\sqrt{2})$. Apply (2.5) at ranks R and $R-1$ and divide their positive weighted sum by $c_R + c_{R-1}/6$. The additional factor $1/\sqrt{2}$ only improves the bound, and the fixed shift changes the constant. This gives the same exponential conclusion. The two options have different actual orbit profiles and are disjoint. □

3 Exact terminal attachment

For a critical product D , let $N_D(T)$ count subgroups of $D \times T$ full on T and on each displayed critical factor. The group T is the complete exterior projection, with all of its internal couplings. Write $d_2(T) = \dim \text{Hom}(T, C_2)$ and $A_2(T) = T/(T'T^2)$.

Lemma 3.1 (Binary contraction). *If $O^2(T)$ is the intersection of the normal subgroups of T with 2-group quotient, image and full preimage induce a bijection*

$$N_D(T) = N_D(T/O^2(T)).$$

Proof. Goursat's lemma expresses a subgroup above its two projections by a common quotient. The projection inside D is a 2-group, so the common quotient of T is a 2-group. Its kernel contains $O^2(T)$. Every subgroup being counted therefore contains $1 \times O^2(T)$. $\square$

Lemma 3.2 (Central extension fibre). *Let $1 \rightarrow K \rightarrow X \rightarrow Y \rightarrow 1$ be central with K elementary abelian of exponent two, and let $\xi \in H^2(Y, K)$ be its class. Define*

$$\text{Ann}(\xi) = \{\lambda \in K^* : \lambda_* \xi = 0 \text{ in } H^2(Y, \mathbb{F}_2)\}.$$

The number of subgroups of X mapping onto Y equals

$$\sum_{L \leq \text{Ann}(\xi)} 2^{d_2(Y) \dim L}. \quad (3.1)$$

Proof. The annihilator is the image of restriction $\text{Hom}(X, C_2) \rightarrow K^*$: a normalized section identifies extension of a character with its scalar defect being a coboundary. Fix $W = H \cap K$. Taking kernels bijects these subgroups with homomorphisms $r : X \rightarrow K/W$ restricting to the quotient map on K . Such an r exists exactly when every scalar coordinate extends, or $W^\perp \leq \text{Ann}(\xi)$. Differences of extensions factor through Y , so this fibre is a torsor under $\text{Hom}(Y, K/W)$, of size $2^{d_2(Y) \text{codim } W}$. Different intersections are disjoint. $\square$

Let $V = D/K = \mathbb{F}_2^r$, $c = \dim K$ and $\delta = r/2 - c$. For an arbitrary finite group T define

$$d = d_2(T), \quad \tau = \dim \ker(H^2(A_2(T), \mathbb{F}_2) \rightarrow H^2(T, \mathbb{F}_2)), \quad X_r(d) = \sum_j \begin{bmatrix} r \\ j \end{bmatrix}_2 2^{dj}. \quad (3.2)$$

The kernel in this definition is retained as a subspace, not merely replaced by the number τ , until a positive upper bound is taken.

Proposition 3.3 (Exact quotient records). *For $0 \leq u \leq r$, let $\mathcal{P}_u$ be the linear maps $p : \mathbb{F}_2^u \oplus A_2(T) \rightarrow V$ injective on the first summand and surjective onto each displayed critical quotient coordinate. Pull back the critical extension to $C_2^u \times T$, and call its class ξ_p . Then*

$$N_D(T) = \sum_{u=0}^r \frac{1}{|\text{GL}(u, 2)| 2^{ud}} \sum_{p \in \mathcal{P}_u} \sum_{L \leq \text{Ann}(\xi_p)} 2^{(u+d) \dim L}. \quad (3.3)$$

Proof. The image $Y \leq V \times T$ of a subgroup being counted is onto T . Put $U = Y \cap V$. Its graph map $T \rightarrow V/U$ factors through $A_2(T)$. A linear lift of this map identifies Y with $U \times T$. There are exactly $|\text{GL}(u, 2)| 2^{ud}$ representations by a choice of ordered basis for U and a linear lift. Lemma 3.2 counts every subgroup above Y ; surjectivity modulo the critical Frattini kernels gives the required full projections. $\square$

Proposition 3.4 (Uniform terminal bound). *With the notation above,*

$$N_D(T) \leq \phi^{-2} \sum_{j=0}^r \sum_{\ell=0}^c \left[\begin{matrix} c \\ \ell \end{matrix} \right]_2 72^\ell 2^{\tau\ell} 2^{(j-\ell)(r-j+d)}. \quad (3.4)$$

Consequently

$$\frac{N_D(T)}{X_r(d)} \leq (r+2)^{O(1)} \left(1 + \sum_{\ell \geq 1} 2^{-\ell(\delta+d/2-\tau-\log 72)-3\ell^2/4} \right). \quad (3.5)$$

In particular, if $\tau = 0$ then $N_D(T) \leq C(r+2)^k X_r(d)$ for absolute C, k .

Proof. Put $u = r - j$ and $k = u + d$. For an ℓ -dimensional space of splitting characters in K^* use row-echelon coordinates as in Proposition 2.4. The original product section has a bilinear scalar defect with diagonal q_λ . If its pullback to $U \times T$ is a coboundary, evaluation on a vertical involution, comparison on commuting vertical and exterior elements, and restriction to T give, respectively,

$$q_\lambda|_U = 0, \quad \beta_\lambda(U, f(A_2(T))) = 0, \quad q_\lambda \circ f \in \ker(H^2(A_2(T)) \rightarrow H^2(T)).$$

Diagonal evaluation $c(a, a) - c(0, 0)$ descends to a linear map on actual $H^2(A_2(T), \mathbb{F}_2)$, since it vanishes on coboundaries. The first two conditions force the whole quadratic outcome to be the exterior pullback of the diagonal of a class in the retained inflation kernel. This allowed function space has dimension at most τ . Modulo that space, the joint row relations determine every pivot outcome. Each outcome has at most 2^τ representatives, and Lemma 2.3 leaves at most 72 surjective maps for each representative on the whole k -dimensional domain. Dropping the other restrictions leaves at most $72^\ell 2^{\tau\ell} 2^{(r-2\ell)k}$ ordered maps. The record divisor is at least $\phi 2^{uk}$. Multiplication by the fibre size $2^{k\ell}$ and summation over L prove (3.4), even with one fewer factor ϕ^{-1} .

Use $\left[\begin{matrix} c \\ \ell \end{matrix} \right]_2 \leq \phi^{-1} 2^{\ell(c-\ell)}$ and write $A = r + d$. The exponent apart from this fixed constant is

$$j(A - j) + \ell(c + \tau + \log 72 - \ell - A + j).$$

If $d \leq r$, its unconstrained maximum in j is

$$A^2/4 - \ell(\delta + d/2 - \tau - \log 72) - 3\ell^2/4,$$

whereas an integer j nearest $A/2$ gives $X_r(d) \geq 2^{A^2/4 - O(1)}$. If $d \geq r$, the exponent is increasing on $0 \leq j \leq r$; its maximum is at $j = r$. Divide by the literal term $2^{rd} \leq X_r(d)$. The remaining exponent is $\ell(r/2 - \delta + \tau + \log 72 - d - \ell)$, which is smaller than the one in (3.5) by $\ell(d - r)/2 + \ell^2/4$. The $r + 1$ choices of j are a polynomial factor. Finally $\delta, d \geq 0$, so at $\tau = 0$ the series is bounded by $\sum_{\ell \geq 0} 72^\ell 2^{-3\ell^2/4} < \infty$. $\square$

Remark 3.5. The module-extension annihilator used below is different from the group-extension annihilator in (3.1). No argument identifies the two. In particular a scalar quadratic form which is nonzero on $A_2(T)$ may become zero after inflation to T .

4 A complete small-support estimate

In even degree write $n = m + b$, with m, b even. Let $M_{n,b}$ count subgroups whose union of noncritical orbits has degree exactly b ; all other orbits are from Table 1. This definition includes all couplings within the noncritical support, as well as every critical lift.

Lemma 4.1. *For $J \leq S_b$ one has $\tau_2(J) \leq \lfloor b/2 \rfloor$. If b is even and some actual orbit of J is noncritical, then $d_2(J) \leq b/2 - 1$.*

Proof. Let $U = J'J^2$. In the five-term sequence for $1 \rightarrow U \rightarrow J \rightarrow A_2(J) \rightarrow 1$, inflation on H^1 is an isomorphism. Thus transgression identifies the inflation kernel in degree two with

$$H^1(U, \mathbb{F}_2)^J \cong \text{Hom}(U/(U'[J, U]U^2), \mathbb{F}_2).$$

The latter is dual to an elementary abelian section of $U \leq S_b$, so has dimension at most $b/2$.

The same abelian quotient bound gives $d_2(J) \leq b/2$. If equality holds, the moved-support equality theorem of [KP89] applies with the Sylow 2-subgroup moving all b points. Its transitive equality constituents are C_2 , regular C_4 , regular V_4 , D_8 , the degree-eight central product of two D_8 groups, and the stated odd affine constituents. The odd affine constituents have positive moved-point defect; the regular C_4 has elementary rank one instead of two. Equality in elementary rank therefore leaves precisely the four actions of Table 1 on every orbit. A noncritical orbit excludes equality, and integrality gives the claimed gap. This is a deduction from the equality classification, not an assertion of independent rank additivity for arbitrary subdirect products. $\square$

Theorem 4.2 (Square-root support). *Fix $0 < \varepsilon < 2$. Uniformly for even $b \geq 2$ with $b \leq (2 - \varepsilon)\sqrt{m}$,*

$$\frac{M_{n,b}}{L_n} \leq 2^{-\varepsilon(4-\varepsilon)m/16+o_\varepsilon(m)}.$$

The sum over this entire range is exponentially small relative to L_n .

Proof. Put $R = m/2$, $N = n/2$. For a critical word with multiplicities (a, v, d, e) its central defect is $\delta = a/2 + v + e$, and its weight is

$$W = \frac{1}{2^a a! 24^v v! 8^d d! 384^e e!}.$$

Let $c_R^{(<L)}$ be the sum of these weights with $\delta < L$. Then $a + 2v + 4e < 4L$, so $d > R/2 - 2L$. There are at most $(R + 1)^4$ tuples and each weight is at most $1/d!$. Conversely one term in c_N uses $\lfloor N/4 \rfloor$ extraspecial factors and at most one each of C_2, D_8 to fill the remainder. Hence

$$c_R^{(<L)} \leq \frac{(R + 1)^4}{\max(0, \lfloor R/2 - 2L \rfloor)!}, \quad c_N \geq \frac{1}{16 \cdot 384^{\lfloor N/4 \rfloor} \lfloor N/4 \rfloor!}.$$

Stirling's inequalities imply, for $L = o(R)$ and $N - R = o(R)$,

$$\log \frac{c_R^{(<L)}}{c_N} \leq -\frac{R}{4} \log R + O(R + L \log(R + 2) + (N - R) \log(R + 2)). \quad (4.1)$$

Let Y_b count subgroups on b points all of whose orbit projections are noncritical. Recovering the noncritical support and its complete projection gives

$$\frac{M_{n,b}}{n!} \leq \frac{Y_b}{b!} \sum_{\gamma: \text{rank } \gamma = R} W(\gamma) \max_{J \in \mathcal{Y}_b} N_{D_\gamma}(J). \quad (4.2)$$

Here $Y_b \leq s_b \leq 2^{b^2/16+O(b^{3/2})}$ by (1.1). There is no extra support-choice factor: it is already in the exact labelled weights of (4.2).

Choose $L = b + \log 72 + \log(R + 2)$. For $\delta \geq L$, Lemma 4.1 and Proposition 3.4 give $N_D(J) \leq (R + 2)^{O(1)} X_R(b/2 - 1)$. The quantity $X_R(t)$ counts subspaces of $\mathbb{F}_2^{R+t}$ transverse to a fixed t -space: choose the projection in $\mathbb{F}_2^R$ and its graph map. Thus

$$\frac{X_R(b/2 - 1)}{G_N} \leq \frac{G_{N-1}}{G_N} \leq 2^{-N/2 + O(1)}.$$

Also (1.3) gives $c_R/c_N \leq (2N)^{b/2}$. The logarithm of the normalized large-defect part is consequently at most

$$-N/2 + b^2/16 + O(b^{3/2} + (b + 1) \log(n + 2)).$$

Since $b \leq (2 - \varepsilon)\sqrt{m}$, $N/2 - b^2/16 \geq \varepsilon(4 - \varepsilon)m/16 + b/4$. The error is uniformly $o_\varepsilon(m)$.

For $\delta < L$, maximize the ℓ exponent in (3.5) using $\tau \leq b/2$. The additional factor is

$$2^{b^2/12 + O(b + \log(R + 2))}.$$

The remaining term satisfies $X_R(b/2 - 1) \leq G_N$. Equation (4.1) bounds its normalized contribution by

$$2^{-R \log R/4 + O(R + b^2 + b^{3/2} + (b + \log(R + 2)) \log(R + 2))}.$$

For $b = O(\sqrt{R})$ this is $2^{-\Omega(R \log R)}$. Finally only $O(\sqrt{n})$ support sizes occur, which changes the logarithm by $O(\log n)$. $\square$

5 Annihilator-aware section capacity

All sections in this section are sections of one specified permutation module. This restriction supplies the capacity bound; odd order alone does not supply it.

5.1 Permutation-module bounds

For a transitive group of degree u , write u_2 for the binary part of u , and set $\text{lpp}(1) = 1$. The induced-module theorem of [Tra18] bounds the number of generators of any submodule of the binary permutation module by

$$E(u, 2) = \min \left\{ \left\lfloor \sqrt{2/\pi} \frac{u}{\sqrt{\log u_2}} \right\rfloor, \frac{u}{\text{lpp}(u/u_2)} \right\}$$

when no soluble transitive subgroup exists. The first entry is infinite if $u_2 = 1$. When such a subgroup exists the applicable bound is

$$E_{\text{sol}}(u, 2) = \min \left\{ \frac{u}{2^{K(u)}} \binom{K(u)}{\lfloor K(u)/2 \rfloor}, u_2 \right\}, \quad K(u) = \sum_{p^a \parallel u} a(p - 1).$$

Both formulas imply the common integer bound $d_T(L) \leq E(u, 2)$: use $K(u) \geq \log u_2$, the central-binomial estimate, $u_2 \leq u/\text{lpp}(u/u_2)$, and integrality of generator number. The sharper soluble formula will be needed as well.

Directly from these formulas,

$$E(u, 2) \leq u/2 \ (u \geq 2), \quad E(u, 2) \leq 3u/8 \ (u \geq 8), \quad E(u, 2) \leq 11u/32 \ (u \geq 24 \text{ even}). \quad (5.1)$$

For a nontrivial odd part use $E \leq u/3$. For powers of two the degrees 2, 4, 8, 16, 32 give respectively 1, 2, 3, 6, 11; from 64 onward the square-root expression is less than $u/3$.

Lemma 5.1 (Orbit filtration). *Let $H \triangleleft T$ have k orbits of length $u \geq 2$ in a transitive action of degree $s = ku$. Every trivial H -section of $\mathbb{F}_2^s$ has dimension at most $kE(u, 2)$.*

Proof. Filter a submodule by its intersections with successive sums of the actual orbit modules. Each successive section embeds in one orbit module. Right exactness of coinvariants bounds the dimension of the coinvariants of the original submodule by the sum of the corresponding dimensions of these sections. Each is bounded by its module-generator number and hence by $E(u, 2)$. A trivial quotient factors through coinvariants. This filtration does not require independent orbit images. $\square$

Lemma 5.2 (A sharper fixed-section bound). *If T is transitive of even degree $s \geq 24$, every submodule L of $\mathbb{F}_2^s$ satisfies $d_T(L) \leq 5s/16$. Consequently every section A satisfies $\dim A^T \leq \lfloor 5s/16 \rfloor$.*

Proof. Write $s = 2^a o$ with o odd. If $o \notin \{1, 3\}$, then $\text{lpp}(o) \geq 5$, and the two appropriate bounds are respectively $E \leq s/5$ and $E_{\text{sol}} \leq s/o \leq s/5$. If $o = 1$, a Sylow 2-subgroup is transitive: for a point x its stabilizer in a Sylow subgroup has order at most $|T_x|_2 = |T|_2/s$. Here $a \geq 5$. If $o = 3$ and a soluble transitive subgroup exists, then $a \geq 3$ and $K(s) = a + 2 \geq 5$. In both cases the central-binomial proportion is at most $\binom{5}{2}/2^5 = 5/16$; it is nonincreasing.

It remains to consider $s = 3 \cdot 2^a$ without a soluble transitive subgroup. Apply [Tra18, Corollary 3.12] to an actual minimally transitive subgroup. It supplies a soluble subgroup with, for $0 \leq j \leq e$, exactly $\binom{e}{j} 2^{u_1}$ orbits of length $3p^j 2^{u-u_1}$, where $p = 2^h - 1$ is a Mersenne prime, $h \geq 3$, $e \geq 1$, $a = eh + u$, and $u \geq u_1 \geq 0$. Filter by these actual orbit modules. The soluble bound on each is at most its binary part, giving

$$d_T(L) \leq 2^{e+u_1} 2^{u-u_1} = 2^{e+u} \leq s/12.$$

Restriction can only increase the required generator number, and generator number is subadditive on the filtration. Finally lift A^T to a submodule of $\mathbb{F}_2^s$; its trivial quotient requires $\dim A^T$ generators. $\square$

Lemma 5.3 (Unipotent sections). *Suppose that $T \leq S_s$ is faithful and transitive and is not a 2-group. If every composition factor of a section P of $\mathbb{F}_2^s$ is trivial, then $\dim P \leq 3s/8$.*

Proof. The action on P is conjugate to a group of unitriangular matrices, so $O^2(T)$ acts trivially. This is a nontrivial normal subgroup. It is generated by odd-order elements, as are its images on its orbits; each nontrivial transitive orbit image is therefore nonbinary. All orbit lengths are equal, say u . For $u \geq 8$, use Lemma 5.1. For $u = 3, 5, 6, 7$, choose an odd prime dividing u . Its Sylow subgroup has no fixed points, since the p -part of a point stabilizer is strictly smaller than the p -part of the group. Exactness of invariants for this odd group bounds any trivial section by the number of its orbits, at most $u/p \leq 3u/8$.

For $u = 4$, a nonbinary transitive subgroup contains a 3-cycle with fixed point x . Its invariant subspace has dimension two. A trivial section of dimension two would force the numerator to contain this entire invariant subspace, including the coordinate vector at x . Transitivity would force the numerator to be the whole permutation module, whose largest trivial quotient has dimension one. This is a contradiction. Thus the bound holds at $u = 4$ too. Sum over the actual orbit filtration as in Lemma 5.1. $\square$

5.2 A coupled socle estimate

Proposition 5.4. *For every section M of the binary permutation module of a faithful transitive group T of even degree $s \geq 24$, put $t = \dim M^T$ and $r_{\text{nt}}(M) = \max_{X \neq 1} m_X(\text{Soc } M) / \dim_{\text{End}_T X} X$. Then*

$$2t + 4r_{\text{nt}}(M) \leq 47s/48. \quad (5.2)$$

No splitting condition on M^T is required.

Proof. We have $t \leq 11s/32$ by (5.1). If there is a nontrivial simple type, choose X attaining the maximum and write $h = \dim_{\mathbb{F}_2} X$, $\text{End}_T X = \mathbb{F}_{2^f}$, $d = h/f$, $m = m_X$, and $H = \ker(T \rightarrow \text{GL}(X))$. The mixed section $1^t \oplus X^m$ has dimension $L = t + hm$.

If $H \neq 1$, it has orbit length at least two, and Lemma 5.1 gives $L \leq s/2$. For $hd \geq 3$,

$$2t + 4m/d = \frac{4}{hd}L + \left(2 - \frac{4}{hd}\right)t \leq \frac{11s}{16} + \frac{5s}{8hd} \leq \frac{43s}{48}.$$

The only remaining nontrivial possibility is $h = 2, d = 1$. The image is then C_3 , acting as nonzero scalars of $\mathbb{F}_4$. There are one or three H -orbits, each of length at least eight. Thus $L \leq 3s/8$, and the required cost is $2L \leq 3s/4$.

If $H = 1$, the group embeds in $\text{GL}_d(2^f)$. Since its degree is even, $d \neq 1$. The values $hd = fd^2 < 9$ are 4 and 8. The first would give $|T| \leq 6$, impossible. In the second case $T \leq \text{GL}_2(4)$, of order 180. Since $s \mid |T|$ and $s \geq 24$, the odd part of s has largest prime-power divisor at least five. Thus $t \leq s/5$, and $L \leq s$ gives $2t + 4m/d \leq s/2 + 3t/2 \leq 4s/5$. For $hd \geq 9$, the same dimension bound gives

$$2t + 4m/d \leq \frac{11s}{16} + \frac{21s}{8hd} \leq \frac{47s}{48}.$$

If there is no nontrivial type, the initial bound on t suffices. $\square$

5.3 The retained annihilator and the central cut

For a finite $\mathbb{F}_2 B$ -module A , put $S = A^B$ and, for each simple X , define the actual module-extension annihilator

$$\mathcal{L}_X = \ker(\text{Ext}_B^1(X, S) \longrightarrow \text{Ext}_B^1(X, A)).$$

Every $C \leq S$ has a module complement in S , so $\text{Ext}_B^1(X, C)$ embeds as a specified subspace of $\text{Ext}_B^1(X, S)$. The long exact sequence gives

$$m_X(A/C) = m_X(A) - \mathbf{1}_{X=1} \dim C + \dim_{\text{End}_B X}(\mathcal{L}_X \cap \text{Ext}_B^1(X, C)). \quad (5.3)$$

Here multiplicity always means socle multiplicity. In particular

$$\dim(A/C)^B = \dim A^B - \dim C + \dim \ker(H^1(B, C) \rightarrow H^1(B, A)). \quad (5.4)$$

These identities explicitly account for newly exposed simple constituents.

Proof of Theorem 1.2. Let $t = \dim S$, $\mathcal{L}_1 = \ker(H^1(B, S) \rightarrow H^1(B, A))$, and $\ell = \dim \mathcal{L}_1$. The invariants sequence identifies $\mathcal{L}_1$ with $(A/S)^B$. Its preimage in A is a section of dimension $t + \ell$ with only trivial composition factors. Hence

$$t \leq \lfloor 5s/16 \rfloor, \quad t + \ell \leq 3s/8 \quad (5.5)$$

by Lemmas 5.2 and 5.3.

Choose bases in S and $H^1(B, \mathbb{F}_2)$, so that $H^1(B, S) = H^1(B, \mathbb{F}_2) \otimes S$. Row-reduce $\mathcal{L}_1$ in these coordinates. Its ℓ pivot positions meet at most ℓ coordinate blocks indexed by basis vectors of S . Let $Z \leq S$ be spanned by the other basis vectors. Then

$$\dim Z \geq \max(t - \ell, 0), \quad \mathcal{L}_1 \cap H^1(B, Z) = 0.$$

A nonzero vector in a row-reduced row space has a nonzero pivot coordinate, which proves the second assertion even when nonpivot supports overlap. Choose $C \leq Z$ of dimension

$$c = \min\{\lfloor t/2 \rfloor, \max(t - \ell, 0)\}.$$

Equation (5.4) gives $\dim(A/C)^B = t - c$. Apply Proposition 5.4 to this actual quotient:

$$2c + 4r_{\text{nt}}(A/C) \leq 47s/48 - 2t + 4c \leq 47s/48.$$

For its trivial type the cost is

$$2c + 4(t - c) = 4t - 2c \leq \max\{3t + 1, 2(t + \ell)\}.$$

The inequality follows by considering which entry attains the minimum defining c , including $\ell \geq t$. By (5.5),

$$3t + 1 \leq 15s/16 + 1 \leq 47s/48, \quad 2(t + \ell) \leq 3s/4.$$

The first inequality uses $s \geq 24$. Taking the maximum over the trivial and nontrivial types proves (1.4) with one and the same C . $\square$

Remark 5.5 (Why the permutation section matters). On $\mathbb{F}_2^t \oplus \mathbb{F}_2^t$ with trivial C_3 action, every linear graph from one summand to the other is invariant, giving 2^{t^2} graphs. Thus odd order supplies no strict quadratic deficit in an arbitrary trivial representation. The proof above instead uses (5.5), the actual annihilator, and a quotient cut. It does not assert a strict deficit for the ranks of the uncut trivial axes.

6 Same-source lifts and original-weight fusion

6.1 The binary permutation rank bound

The binary rank input needed below has a direct degree-induction proof. Write $d_2(P) = \dim_{\mathbb{F}_2} \text{Hom}(P, C_2)$; for a finite 2-group this is also the dimension of its Frattini quotient.

Lemma 6.1. *If a finite 2-group P acts faithfully on b points, then $d_2(P) \leq \lfloor b/2 \rfloor$.*

Proof. Induct on b , with degrees zero and one immediate. For an intransitive action, restrict to a nonempty proper orbit of size a . Its actual image acts faithfully on that orbit, and the restriction kernel acts faithfully on the original complement. Character restriction and induction give

$$d_2(P) \leq d_2(\text{image}) + d_2(\text{kernel}) \leq \lfloor a/2 \rfloor + \lfloor (b - a)/2 \rfloor \leq \lfloor b/2 \rfloor.$$

For a transitive action, a cover of the original point stabilizer has relative index two. Its actual blocks are pairs, so $b = 2m$. Let Q be the faithful action on the m pairs and K its original kernel.

The flip coordinates identify K with a P -invariant subspace of $\mathbb{F}_2^m$, retaining all correlations among the flips. Restriction gives

$$d_2(P) \leq d_2(Q) + \dim \operatorname{Hom}(K, \mathbb{F}_2)^P.$$

If $m = 1$, the top is trivial and $\dim K \leq 1$. Otherwise the transitive 2-group Q has a fixed-point-free central involution: any nonidentity central element of order two is semiregular in a faithful transitive action. Choose a lift $g \in P$. Every invariant form on K annihilates $(g - 1)K$, hence rank-nullity gives

$$\dim \operatorname{Hom}(K, \mathbb{F}_2)^P \leq \dim K / (g - 1)K = \dim \ker((g - 1)|_K) \leq \dim \ker((g - 1)|_{\mathbb{F}_2^m}) = m/2.$$

Induction bounds $d_2(Q)$ by $\lfloor m/2 \rfloor$, so the total is at most $m = b/2$. No splitting of the original extension is used. $\square$

6.2 The fractional Schur cost

The following lemma applies to nonsplit group extensions and to nonsemisimple modules.

Lemma 6.2 (Conditional cocycle bound). *Let $1 \rightarrow A \rightarrow Q \rightarrow B \rightarrow 1$, with A elementary abelian of exponent p , and let $r(A) = \max_X m_X(\operatorname{Soc}_B A) / \dim_{\operatorname{End}_B X} X$. For $J \leq S_b$ and a fixed epimorphism $\beta : J \rightarrow B$, the number of epimorphisms $J \rightarrow Q$ above β is at most*

$$|A| |H^1(B, A)| p^{(b/p)r(A)}. \quad (6.1)$$

Proof. If a lift exists, choose one actual lift as origin. All homomorphic lifts are a torsor for $Z^1(J, A_\beta)$; onto lifts are a subset. Let $F = \ker \beta$. Inflation–restriction and the bound on coboundaries give

$$|Z^1(J, A_\beta)| \leq |A| |H^1(B, A)| |\operatorname{Hom}(F, A)^J|.$$

Choose a Sylow p -subgroup $P \leq F$ and put $D = N_J(P)$. The Frattini argument gives $J = FD$, so $D \rightarrow B$ is onto. Restriction embeds

$$\operatorname{Hom}(F, A)^J \hookrightarrow \operatorname{Hom}_D(P/\Phi(P), A).$$

Indeed P maps onto every p -group image of F . The source here has the actual D -action; an action of B on it is not assumed. Filter $W = P/\Phi(P)$ by a composition series. Left exactness of $\operatorname{Hom}(-, A)$ gives

$$\dim \operatorname{Hom}_D(W, A) \leq \sum_X [W : X] m_X \dim_{\mathbb{F}_p} \operatorname{End}_D X \leq r(A) \dim W.$$

Types not inflated from B contribute zero. Since $D \rightarrow B$ is onto, the remaining simple dimensions and endomorphism fields are precisely those used in $r(A)$. Finally $\dim W \leq b/p$ by the permutation p -group abelianization bound, whose binary case is Lemma 6.1. This proves (6.1). $\square$

Proposition 6.3 (Central cut and graph moment). *Let $1 \rightarrow A \rightarrow Q \rightarrow B \rightarrow 1$ have elementary binary kernel and let $C \leq A^B$, $c = \dim C$. Suppose $T \leq S_s$ is a specified permutation cover $\pi : T \rightarrow B$. Put*

$$D_C = |A/C| |H^1(B, A/C)|, \quad \lambda = \frac{1}{2}r(A/C), \quad v = s + 2c, \quad \Phi(J) = |\operatorname{Epi}(J, B)| 2^{cd_2(J)}.$$

For $A/C = 0$ take $D_C = 1$. Then, for every b and integer $q \geq 1$,

$$|\operatorname{Epi}(J, Q)| \leq D_C 2^{\lambda b} \Phi(J), \quad \sum_{J \leq S_b} \Phi(J)^q \leq s_{b+qv}. \quad (6.2)$$

Proof. Since $C \leq Z(Q)$, each actual map $J \rightarrow Q/C$ has either no lift or a torsor of homomorphic lifts under $\text{Hom}(J, C)$. Apply Lemma 6.2 to the actual extension $1 \rightarrow A/C \rightarrow Q/C \rightarrow B \rightarrow 1$ and sum over the same maps to B .

A term $\Phi(J)^q$ selects q pairs of literal maps $(\beta_i : J \rightarrow B, \chi_i : J \rightarrow C_2^c)$. Encode them by the subgroup

$$\{(t_1, e_1, \dots, t_q, e_q, j) : \pi(t_i) = \beta_i(j), e_i = \chi_i(j) \text{ for all } i\} \leq (T \times C_2^c)^q \times J.$$

Realize each C_2^c on c disjoint pairs. Projection onto the fixed coordinate sets recovers J and every selected map. Thus the encoding is injective into the subgroups of S_{b+qv} . Repeated maps, dependent characters, and proper joint subdirect images are allowed. $\square$

Proposition 6.4 (A target-order alternative). *Let Q be a finite 2-group of order 2^a . For every finite group J and all integers $b, q \geq 0$,*

$$|\text{Hom}(J, Q)| \leq 2^{ad_2(J)}, \quad \sum_{J \leq S_b} (2^{ad_2(J)})^q \leq s_{b+2aq}.$$

Consequently this supplies the fusion parameters $D = 1$, $\lambda = 0$, $v = 2a$ whenever the original physical width satisfies $2a < w$.

Proof. Every nontrivial finite 2-group has a central subgroup C of order two. Each nonempty fibre of $\text{Hom}(J, Q) \rightarrow \text{Hom}(J, Q/C)$ is a torsor under $\text{Hom}(J, C)$: the pointwise difference of two lifts is a homomorphism because C is central. Its cardinality is $2^{d_2(J)}$. Induction on a proves the first inequality, including empty fibres and nonsplit extensions. For the second, encode the aq binary characters together as a graph over the same literal J , realizing each binary coordinate on a pair of new points. $\square$

This argument counts actual maps; it takes no quotient by target automorphisms and does not require a faithful action of Q on $2a$ points. Its width condition is distinct from the character criterion used later.

6.3 A complete finite-menu estimate

Theorem 6.5 (Original-weight fusion). *Fix a finite collection of pairs (U, N) , where $U \leq S_w$ is transitive, w is even, and $N \triangleleft U$. For each pair suppose there are fixed D, v, λ and a nonnegative function Φ such that*

$$|\text{Epi}(J, U/N)| \leq D2^{\lambda b} \Phi(J), \quad \sum_{J \leq S_b} \Phi(J)^q \leq s_{b+qv} \quad (q \geq 1), \quad g = w - v - 8\lambda > 0.$$

Let $\mathcal{U}_n$ be any family of actual subgroups of S_n each having an orbit of type U whose intersection axis is one of the specified N . Then

$$\frac{|\mathcal{U}_n|}{L_n} \leq \eta(n) + \sum_{b < n} K(n, b) \frac{s_b}{L_b}, \quad \eta(n) = O(2^{-an^2}), \quad \sum_b K(n, b) = O(2^{-a'n}) \quad (6.3)$$

for some $a, a' > 0$. The weights are those of the original actions. The result is uniform in ambient parity, but its constants may depend on the fixed finite menu.

Proof. For each U , first close the selected literal axes under $N_{S_w}(U)$. This finite enlargement leaves the existential physical orbit event unchanged. Choose one certificate on each resulting normalizer orbit and transport it through the induced quotient isomorphisms; thus D, v, λ, Φ can be constant on that orbit. All sums below refer to this closed literal-axis menu.

On a pointed orbit the subgroup has exact Goursat form

$$\{(u, j) \in U \times J : uN = \beta(j)\}$$

for a unique literal epimorphism $\beta : J \rightarrow U/N$, once the quotient map is fixed. This counts actual maps, so there is no additional division by $|\text{Aut}(U/N)|$. First apply the pointing factor $n!/(b!a(U))$ to the total normalizer-stable sum over literal axes and complete sources. A single axis need not be fixed by the action normalizer. Only after this division do we split the numerical sum entry by entry; the envelopes need not be conjugacy invariant. For an entry put $e = g/16$, $a_0 = v/8 + e$, and call its complete source hot when $\Phi(J) > 2^{a_0 b}$. If $v = 0$, the moment bound has the same finite right-hand side s_b for every q . Hence $\Phi(J) \leq 1$ for every J , and the strict hot set is empty. We may therefore assume $v > 0$ in the hot estimate below.

For a hot source and any integer $q \geq 1$,

$$\sum_{J \text{ mhot}} |\text{Epi}(J, U/N)| \leq D 2^{\lambda b - (q-1)a_0 b} s_{b+qv}.$$

Choose $q = \max(1, \lceil 8eb/v^2 \rceil)$. The exact identity

$$\begin{aligned} & \frac{(b+qv)^2}{16} - (q-1)a_0 b + \lambda b - \frac{(b+w)^2}{16} \\ &= -\frac{4e^2 b^2}{v^2} - eb - \frac{w^2}{16} + \frac{v^2}{16} \left(q - \frac{8eb}{v^2} \right)^2 \end{aligned} \quad (6.4)$$

has a strictly negative quadratic term. The all-pairs monomial gives $c_R \geq 1/(2^R R!)$. Thus, writing $R = \lfloor n/2 \rfloor$ and using $b \geq w$ so that $R \leq b$,

$$\frac{n!}{b!L_n} \leq \phi^{-1} 2^{-R^2/4 + R + 1/4} \leq \phi^{-1} 2^{-n^2/16 + 5n/8 + 1/4}.$$

Together with (1.1), this contributes only $O(n^{3/2} + n)$ to the normalized logarithm. No saddle estimate is needed for this normalization. As all entry parameters are fixed, the complete finite hot union is $O(2^{-an^2})$.

On its complement, $|\text{Epi}(J, U/N)| \leq D 2^{(w/8 - g/16)b}$. The cold kernel is bounded by the sum over entries of

$$\Delta(n, b) \frac{D}{a(U)} 2^{(w/8 - g/16)b}, \quad b = n - w. \quad (6.5)$$

For $w = 2h$, (1.3) and the Gaussian bounds give the explicit estimate

$$\Delta(n, b) \leq \phi^{-2} 2^{(h+1)/4} (n+1)^{h+1} 2^{-hb/4 - h^2/4}.$$

The guarded odd coefficient obeys the same backward-shift bound, including rank zero, so the estimate holds in both parities. Each summand in (6.5) is therefore $2^{-gb/16 + O_w(\log(n+2))}$. The finite sum proves the row estimate.

The hot and cold contributions partition the local numerical sum after pointing. Summing over the finite action menu may overcount, but only by the positive sum already bounded. Restriction by any earlier disjoint family can only reduce it. The actual complement J includes every later orbit and every terminal attachment, as required by (1.6). $\square$

Proposition 6.6 (Direct first-moment fusion). *Fix a finite original action/normal-axis menu as above. For each entry assume the local envelope and only the first moment*

$$|\text{Epi}(J, U/N)| \leq D2^{\lambda b} \Phi(J), \quad \sum_{J \leq S_b} \Phi(J) \leq s_{b+v}.$$

Suppose $w = 2h$, $v = 2r < w$, and $g = w - v - 8\lambda > 0$. For the same actual orbit family, with any retained survival predicates invariant under the relabellings in the pointing formula, there is a nonnegative forward kernel satisfying

$$\frac{|\mathcal{U}_n|}{L_n} \leq \sum_{m < n} K_{\text{dir}}(n, m) \frac{s_m}{L_m}, \quad \sum_{m < n} K_{\text{dir}}(n, m) = O(2^{-\kappa n})$$

for some $\kappa > 0$. This conclusion does not use the coarse estimate (1.1).

Proof. Apply the original pointing factor to the complete literal-axis sum, as in the preceding proof, and then use the first moment. An entry continues at $m = b + v = n - w + v < n$, with coefficient

$$\Gamma_{U,N}(b) = \frac{(b+w)!}{b! a(U)} D2^{\lambda b} \frac{L_{b+v}}{L_{b+w}}.$$

Define $K_{\text{dir}}(n, m)$ by summing these coefficients over all literal entries with this target degree, taking an entry to be zero when $n < w$. Distinct normals and actions remain separate summands, even when their target degrees agree. The divisor is still $a(U)$, and no additional quotient by target automorphisms or marker choices occurs.

Put $k = h - r > 0$. Since $n = m + 2k$, exact factorial cancellation gives

$$\Gamma_{U,N}(b) = \frac{D}{a(U)} \frac{m!}{b!} 2^{\lambda b} \Delta(n, m).$$

The fixed-width benchmark estimate above applies in either parity, and $m!/b! \leq m^v$. Retaining its quadratic width term gives the uniform estimate, valid for every $b \geq 0$,

$$\Gamma_{U,N}(b) \leq \phi^{-2} \frac{D}{a(U)} (n+1)^{(w+v)/2+1} 2^{-gb/8 - (w^2 - v^2)/16 + (w-v+2)/8}. \quad (6.6)$$

Indeed the exponent before simplification is $\lambda b - km/4 - k^2/4 + (k+1)/4$. Consequently, for an entry-dependent constant,

$$\Gamma_{U,N}(b) \leq C(n+1)^{v+k+1} 2^{\lambda b - km/4} \leq C(n+1)^{v+k+1} 2^{-gb/8}.$$

Absorb the fixed polynomial into the exponential and sum the fixed finite menu. This proves the row estimate without estimating any unknown subgroup count s_m . $\square$

The order markers $v = 2 \log_2 |U/N|$ and the character marker $v = 0$ fit this even-prefix argument whenever their strict width conditions hold. Any class-count input used to establish the character envelope remains necessary. The proposition supplies neither a physical cover by the menu nor uniform control when the widths, menu, or lift constants grow with n ; those wider regimes remain separate obligations.

Corollary 6.7 (Nonbinary pair actions). *Let U be a transitive action on $2s$ points with an invariant pair system whose faithful top $T \leq S_s$ is nonbinary, with even $s \geq 24$. For every $N \triangleleft U$, write K for the pair kernel and set*

$$A = KN/N \cong K/(K \cap N), \quad B = U/(KN).$$

This entry has the certificate of Theorem 6.5, with $g \geq s/48$. Hence every fixed finite menu of these entries admits (6.3).

Proof. The pair kernel is a submodule of $\mathbb{F}_2^s$; the action on its indicated section factors through B . Apply Theorem 1.2 and Proposition 6.3. With $c = \dim C$,

$$g = 2s - (s + 2c) - 4r(A/C) \geq s/48.$$

There was no assumption that $N \leq K$, nor that the common quotient or its subdirect cores are abelian. For labelled transport, fix one witness on each original-normalizer orbit of entries and fixed transporters before counting J . A normalizer-equivariant choice of an individual witness is not needed, and the witnesses are not counted as additional decorations. $\square$

Remark 6.8 (Scope of the positive enlargement). The original quotient, extension class, and restricted group transgression determine which lifts exist. The cocycle torsor bounds are applied after those data have been fixed. The capacity proof uses the distinct module annihilator in (5.4). The theorem does not grant bounds for arbitrary extra weights on the source groups, and a growing menu requires an additional uniform estimate for its size and the constants D . Only Φ , not the cocycle tail $2^{\lambda b}$, is raised to the moment power.

7 Complete quotient moments and uniform transfer

For finite X and a complete source J define

$$Z_J(X) = \sum_{N \triangleleft X} |\text{Epi}(J, X/N)|, \quad h(w) = w - (w \bmod 2).$$

The normal subgroups here are literal, including $N = X$. Distinct axes with isomorphic quotient groups remain distinct terms.

Lemma 7.1 (The complete quotient moment). *If R has a specified faithful permutation representation of degree v , then for every integer $q \geq 1$,*

$$\sum_{J \leq S_b} Z_J(R)^q \leq s_{b+qv}.$$

Proof. $Z_J(R)$ counts full subdirect subgroups of $R \times J$ by their intersection with R and their literal graph map. For an ordered q -tuple of such subgroups take their fibre product over the same J inside $R^q \times J$. Projection recovers J and every member of the tuple: each of the other subgroups can lift the same element of J . This is an injection on actual counting sets, including coincident maps and proper joint quotient images. $\square$

Lemma 7.2 (Uniform normalization). *Uniformly for $n = b + w$, $w \geq 1$,*

$$\log \Delta(n, b) \leq -\frac{h(w)b}{8} - \frac{w^2}{16} + O(w \log(n+2)). \quad (7.1)$$

Proof. Let $r = \lfloor b/2 \rfloor$ and $R = \lfloor n/2 \rfloor$. The recurrence gives $c_r/c_R \leq (2R)^{R-r}$, and $p_b \leq (1 + b/6)c_r$, whereas $p_n \geq c_R$. These terms cost $O(w \log(n+2))$, since $R - r \leq (w+1)/2$. The Gaussian bounds give $\log(G_r/G_R) = (r^2 - R^2)/4 + O(1)$. For even w this is $-wb/8 - w^2/16 + O(w)$. For odd w , the parity correction is at most $b/8 + O(w)$, giving (7.1). The smallest indices are absorbed in the same absolute constant. $\square$

Theorem 7.3 (A finite complete comparator). *Fix a finite family of actual transitive actions U of degrees w . Suppose each has a fixed faithfully represented comparator R_U of degree $0 < v < h(w)$ and constants $B > 0$, $C \geq 0$, $\eta < h(w)/8$ such that for all complete $J \leq S_b$,*

$$Z_J(U) \leq BZ_J(R_U) + C2^{\eta b}.$$

Then their complete pointed union has one quadratic-exponentially small scalar and one exponentially small complete forward row, with the original action weights. The comparator need not be a quotient of U .

Proof. Put $\alpha = (v+h)/16$, $e = (h-v)/16$ and $\chi = h/8 - \max(\alpha, \eta) > 0$. Split at $Z_J(R) > 2^{\alpha b}$. The cold summand is exactly bounded by

$$\frac{\Delta(n, b)}{a(U)} (B2^{\alpha b} + C2^{\eta b}).$$

Equation (7.1) gives exponential row decay after the finite sum. On the hot part put $d = \max(\eta - \alpha, 0)$, so the whole fibre is at most $(B+C)2^{db}Z_J(R)$. Use Lemma 7.1 at $q = \max(1, \lceil 8eb/v^2 \rceil)$, $q_* = 8eb/v^2$. The main normalized exponent is

$$\frac{(b+qv)^2}{16} - (q-1)\alpha b + db - \frac{(b+w)^2}{16} = -\frac{4e^2b^2}{v^2} - \chi b - \frac{(w-h)b}{8} - \frac{w^2}{16} + \frac{v^2}{16}(q-q_*)^2.$$

Here $0 \leq q - q_* \leq 1$ and all parameters are fixed. The coarse error in (1.1) and the original factorials cost $O(n^{3/2} + n \log(n+2))$, smaller than the quadratic loss. Take one any-hot union and its cold complement as in Theorem 6.5. $\square$

Theorem 7.4 (Growing-parameter quotient transfer). *Fix $w_0 \geq 5$, $0 < \theta < 1$, and $0 < \rho \leq 1/8$. For each accepted transitive action U of degree $w \geq w_0$, choose one certificate consisting of a quotient R , a faithful action of R on $v_0 \leq \theta h(w)$ points, and $\eta_U \geq 0$. Require $v_0 = 0$ only when $R = 1$, and*

$$\frac{h(w) - v_0}{8} - \eta_U \geq \rho w.$$

Suppose that, for every $J \leq S_b$,

$$Z_J(U) \leq 2^{\eta_U b + B_U(b)} Z_J(R), \quad B_U(b) \leq \beta(w) + Cw \log^2(w+b+1), \quad (7.2)$$

where C is fixed and $\beta(w) = o(w^2)$ uniformly over the certificates. Then the complete accepted-orbit family has a forward kernel and scalar with the uniform eventual bounds

$$K(n, n-w) \leq 2^{-\rho w n/8}, \quad \sum_b K(n, b) \leq 2 \cdot 2^{-\rho w_0 n/8}, \quad \eta(n) \leq 2^{-\rho^2 n^2/8}. \quad (7.3)$$

Proof. Pad the given representation by fixed points to $v = \max(v_0, \lfloor 4\rho h(w) \rfloor)$ and put

$$\delta = (h(w) - v)/8 - \eta_U, \quad a = v/8 + \delta/2.$$

Then $\delta \geq \rho w/2$, $v \leq (1 - 4\rho)w$, and $\eta_U + a = h(w)/8 - \delta/2$. If $R \neq 1$, also $v \geq \rho w$: use $v_0 \geq 2$ when $\rho w \leq 1$, and otherwise $\lfloor 4\rho h(w) \rfloor \geq 4\rho w - 4\rho - 1 > \rho w$. For $R = 1$ the hot event below is empty, so no lower bound on v is needed. This controlled padding preserves the original margin.

Split at $Z_J(R) > 2^{ab}$. The cold kernel is

$$K(n, b) = \Delta(n, b) \sum_{\substack{U \text{ accepted} \\ \deg U = n-b}} \frac{2^{(\eta_U + a)b + B_U(b)}}{a(U)}.$$

By (7.1) its main exponent is at most $-\rho w b/4 - w^2/16 \leq -\rho w n/4 - w^2/32$. The number of transitive action classes is $2^{o(w^2)}$ by the transitive-subgroup bound of [LMM00], in the labelled form quoted in the introduction of [RDT25]. Consequently, after increasing a finite constant, $\beta(w)$ plus this class entropy is at most $w^2/64 + C_1 w$ for every w . The remaining error is $O(w \log^2(n+2)) + C_1 w = o(w n)$ uniformly. This proves the first bound in (7.3). Sum the geometric tail over $w \geq w_0$ to obtain the second.

For a nonempty hot part use $q = \max(1, \lfloor 4\delta b/v^2 \rfloor)$, $q_* = 4\delta b/v^2$, and $M = b + qv$. Lemma 7.1 gives $\sum_{J \text{ hot}} Z_J(R) \leq 2^{-(q-1)ab} s_M$. The exact main exponent, including the lift factor, is

$$\begin{aligned} \frac{M^2}{16} - (q-1)ab + \eta_U b - \frac{n^2}{16} &= -\frac{\delta^2 b^2}{v^2} + (\eta_U + a - w/8)b - \frac{w^2}{16} + \frac{v^2}{16}(q - q_*)^2 \\ &\leq -\frac{\rho^2 b^2}{4} - \frac{\rho w b}{4} - (\rho/2 - \rho^2)w^2 \leq -\rho^2 n^2/4. \end{aligned}$$

The last comparison uses $\rho \leq 1/8$; the adverse odd correction $(h(w) - w)b/8$ is nonpositive. Moreover $M \leq (1 + 1/(2\rho))b + w$, so the coarse error is uniformly $O_\rho(n^{3/2})$. The original hot sum is

$$\sum_{w=w_0}^n \sum_{\deg U=w} \frac{2^{\eta_U b + B_U(b)}}{b! a(U) p_n G_{\lfloor n/2 \rfloor}} \sum_{J \text{ hot}} Z_J(R), \quad b = n - w.$$

Its remaining normalization costs $O(n \log(n+2))$. For any prescribed small positive coefficient, the uniform $o(w^2)$ errors and class entropy are bounded by that coefficient times w^2 plus $O(w)$. All errors, including the sum over widths and $Cw \log^2(n+1)$, are $o(n^2)$ uniformly. Retaining half the quadratic reserve proves the last bound in (7.3). No normalized-count boundedness has been assumed. $\square$

Lemma 7.5 (Central nonbinary pair sections). *For a faithful nonbinary transitive action of even degree $s \geq 4$, every trivial section of its binary permutation module has dimension $t \leq s/2 - 1$. Consequently a central pair-kernel quotient at original width $2s$ has an effective graph moment degree $s + 2t \leq 2s - 2$.*

Proof. If an odd prime p divides s , its Sylow subgroup has no fixed points. Exactness of invariants bounds a trivial section by the number of its orbits, at most $s/p \leq s/2 - 1$. If $s = 2^a \geq 8$, a Sylow 2-subgroup is transitive. The soluble permutation-module bound gives $t \leq B_a \leq 3s/8 \leq s/2 - 1$. At $s = 4$, the 3-cycle fixed-vector argument in Lemma 5.3 gives $t \leq 1$. For a central kernel A take $C = A$ in Proposition 6.3. Then the cocycle tail vanishes and the moment degree is $s + 2t$. $\square$

8 Faithful character tuples on a common source

All characters in this section are complex irreducible characters. A faithful tuple means that the intersection of its kernels is trivial. Slots may contain the trivial character; this permits harmless padding. Binary-primary linear means every power-of-two order, rather than only order two.

8.1 Whole-socle feasibility

Write the abelian part of $\text{Soc}(Q)$ as a direct sum of simple $\mathbb{F}_p Q$ -modules over its various primes. For a nontrivial simple type X let m_X be its multiplicity and $e_X = \dim_{\text{End}_Q} X$. For each prime let m_p be the dimension of the central elementary p -socle and let f_p be the dimension of its intersection with Q' . Put

$$T = \max(\{f_p\}_p, \{[m_X/e_X]\}_{X \neq 1}, \{1 : \text{Soc}(Q) \text{ has a nonabelian factor}\}), \quad M_o = \max_{p \text{ odd}} m_p,$$

with empty maxima zero.

Theorem 8.1 (Three-category feasibility). *A faithful tuple with r binary-primary linear slots, l unrestricted linear slots, and t general irreducible slots exists if and only if*

$$t \geq T, \quad l + t \geq M_o, \quad r + l + t \geq m_2. \quad (8.1)$$

The trivial group permits the empty tuple.

Proof. Every nontrivial normal subgroup intersects the socle nontrivially. It therefore suffices to detect the entire socle with the joint tuple. For the abelian socle, Clifford theory says that the restriction of an irreducible contains one orbit of linear characters. In a simple isotype X^{m_X} , a single orbit spans at most e_X independent multiplicity directions. To see this, identify the image algebra on X with $M_{e_X}(\text{End}_Q X)$; a vector in the isotype supplies at most e_X rows in its multiplicity space. Conversely t vectors generate that isotype whenever $m_X \leq te_X$, by choosing rows to span the whole multiplicity space. The dual module has the same dimensions and multiplicity, so this criterion applies equally to character orbits.

Linear characters kill Q' . All nontrivial abelian socle types lie in Q' , because their commutator with Q is the whole simple type. Thus they require the general slots. So does each derived-central direction, and a nonabelian socle factor requires at least one general slot. Each irreducible is scalar on a central elementary p -group, so a slot detects at most one independent central direction. A binary-primary linear character kills every odd central element. These facts prove necessity of (8.1).

For sufficiency, choose bases of each central p -socle adapted to its derived-central subspace. Assign its first f_p dual coordinates to the general slots, and as many remaining coordinates as those slots can hold. At odd p , put the remaining coordinates in the l linear slots; at $p = 2$, use the $r + l$ linear slots. The inequalities guarantee room. The functionals assigned to linear slots vanish on the entire intersection of the abelian socle with Q' . They extend first to its image in Q^{ab} and then to complex linear characters of Q^{ab} , since a character of a subgroup of a finite abelian group extends to that group. For a binary-primary slot, extend inside the binary component and take the trivial character on all odd components. Higher binary roots may be necessary and are allowed.

On each nontrivial simple isotype choose t dual vectors whose Q -orbits generate its full dual, using the matrix-algebra criterion above. Combine their assignments over all primes with the

chosen central functionals to obtain t characters of the abelian socle. If the nonabelian socle is nontrivial, choose a nontrivial irreducible on each of its simple direct factors and take their tensor product. It is faithful: the simple factors are centreless, and a tensor product can have scalar cancellation only from scalar images. Assign this character to one general slot and use the trivial character there in the other slots.

Each slot now specifies an irreducible of the whole socle. Induce it to Q and choose an irreducible constituent. By Clifford theory, its restriction contains every conjugate of the specified socle character; its kernel on the socle is precisely the corresponding normal core. The chosen orbit spans and central coordinates make the intersection of these cores, together with the extended linear kernels, trivial. The resulting tuple is faithful on Q . $\square$

For the source costs below, the cheapest feasible profile is

$$t = T, \quad l = \max(M_o - T, 0), \quad r = \max(m_2 - \max(M_o, T), 0). \quad (8.2)$$

Indeed replacing an unrestricted linear slot by a general one increases its coefficient from $(\log 3)/3$ to $(\log 5)/3$, and replacing a binary-primary linear slot by either type also increases its coefficient. For fixed t choose the least feasible l and then the least r ; increasing t cannot improve that cost.

8.2 Joint kernel counting

Proposition 8.2. *If Q has a specified faithful tuple with profile (r, l, t) , then for every actual $J \leq S_b$,*

$$|\text{Epi}(J, Q)| \leq |\text{Aut } Q| 2^{rb/2} 3^{lb/3} 5^{tb/3}. \quad (8.3)$$

Proof. For each onto kernel choose one epimorphism and inflate the fixed tuple. Its joint kernel is that original kernel, so distinct kernels give distinct ordered tuples of characters of the same J . There are at most $2^{b/2}$ binary-primary linear characters: a Sylow 2-subgroup of J maps onto the binary component of J^{ab} , and its abelianization has order at most $2^{b/2}$. The total number of linear characters is $|J^{\text{ab}}| \leq 3^{b/3}$ by [RDT25, Theorem 2.8] (with the perfect case immediate). The number of all irreducibles is $k(J) \leq 5^{b/3}$ by [GM15, Theorem 1.1], including $b \leq 3$ by direct inspection. Enlarge the joint tuple set to the product of these three lists. Finally each onto kernel has exactly $|\text{Aut } Q|$ maps. $\square$

Proposition 8.3 (Several anchors, one preimage). *In a specified faithful tuple of profile (r, l, t) choose an anchor subtuple whose joint kernel K is a p -group. Let u be the number of general slots among the anchors, $g = t - u$, and $d = [Q : O_p(Q)]$. Then*

$$|\text{Epi}(J, Q)| \leq |\text{Aut } Q| d^g 2^{rb/2} 3^{lb/3} 5^{ub/3} (38/25)^{gb}. \quad (8.4)$$

If Q itself is a p -group the empty anchor is allowed, with $u = 0, d = 1$.

Proof. The inflated anchor tuple determines $M = \varphi^{-1}(K)$. Since $K \leq O_p(Q)$ and $O_p(Q/K) = O_p(Q)/K$, it determines the single subgroup $F = \varphi^{-1}(O_p(Q))$ as the preimage of the characteristic p -core of J/M . Each remaining general character, restricted to F , has constituents with p -group image. All such constituents factor through the canonical maximal p -quotient of F . An actual Sylow p -subgroup $P \leq F$ maps onto that quotient, so its irreducible list has size at most $k(P) \leq (38/25)^b$ by the nilpotent class bound [Mar05, Theorem 1.5]. No faithful degree is assigned to the abstract quotient. Frobenius reciprocity bounds the number of irreducibles of J above a fixed constituent by $[J : F] = d$: their dimensions are at least the constituent's dimension, whereas its induction

has d times that dimension. The linear slots use the global lists of Proposition 8.2, and only the u general anchors use the general list. Enlarge the joint lists after F has been determined. Faithfulness recovers the onto kernel, and one factor $|\text{Aut } Q|$ counts the maps. For a p -group target $F = J$ is already known and no anchor is needed. $\square$

Corollary 8.4 (Complete finite character entries). *Fix a finite menu of actual actions U . For each literal $N \triangleleft U$ choose one tuple or anchor certificate whose coefficient of b in (8.3) or (8.4) is at most $h(w)/8 - \delta$, where $w = \deg U$ and $\delta > 0$ is common to the menu. The complete accepted-orbit union has zero scalar and a forward kernel with exponentially small row, in original weights.*

Proof. Sum the chosen bounds over the actual normal axes of each U , always on the same complete J . Their fixed target constants have a finite sum D_U , so $Z_J(U) \leq D_U 2^{(h(w)/8 - \delta)b}$. The pointed cold row is $\Delta(n, b) \sum_{\deg U = n-b} D_U 2^{(h(w)/8 - \delta)b} / a(U)$. By Lemma 7.2 its logarithm is at most $-\delta b + O_W(\log(n+2))$, and $b \geq n - W$ for a fixed width cap W . The complete finite sum is exponentially small. $\square$

9 Relative heads of actual transitive actions

For $N \triangleleft A$ and a prime p , write

$$h_{p,A}(N) = \dim_{\mathbb{F}_p} N / (N^p[N, A]), \quad \rho_A(N) = h_{2,A}(N), \quad \delta_A(N) = h_{3,A}(N).$$

The distinction between a relative head and the abelianization of an abstract isomorphic group is essential. For $B \leq C$ normal in A , the quotient word maps onto the corresponding quotient word downstairs; its kernel is an image of the head of B . Consequently

$$h_{p,A}(C) \leq h_{p,A}(B) + h_{p,A/B}(C/B). \quad (9.1)$$

If $G \leq \prod_{i=1}^t A_i$ is full on its actual orbit actions, filter G by the kernels of projection onto coordinates $i+1, \dots, t$. The image of the i th factor is a normal subgroup $N_i \triangleleft A_i$. Conjugation on that factor is exactly the action of A_i , since the coordinate projection of G is onto. Thus

$$d_p(G) \leq \sum_i h_{p,A_i}(N_i). \quad (9.2)$$

This assertion does not add the ranks of the full coordinate images.

Theorem 9.1 (Binary first head). *For a faithful transitive $U \leq S_w$, $w \geq 2$, and $N \triangleleft U$,*

$$\rho_U(N) \leq 3w/8,$$

except for the whole-action pairs (U, U) of the four critical actions in Table 1. Their heads have dimensions 1, 2, 2, 4.

Proof. We use strong induction on w . Tracey's induced-module theorem [Tra18, Theorem 1.6] gives, for a submodule of an induced binary module with fibre dimension a and index $q \geq 2$, at most $aE(q, 2) \leq aq/2$ module generators. Indeed, if q has odd part greater than one, the largest-prime-power branch is at most $q/3$; if $q = 2^k$, the other branch is $\lfloor \sqrt{2/\pi} q / \sqrt{k} \rfloor$, which is at most $q/2$ for $k \geq 3$ and equals 1, 2 for $k = 1, 2$.

Suppose first that N is intransitive, with $q \geq 2$ orbits of size m . Let K be the kernel of U on these orbits and let H be its transitive action on one of them. The faithful embedding of K in H^q and its coordinate-kernel filtration give

$$\rho_U(N) \leq \rho_K(N) \leq \sum_{i=1}^q \rho_H(E_i), \quad E_i \triangleleft H.$$

If H is not critical, induction suffices. In each critical action every proper normal subgroup has relative head at most $3m/8$: for C_2, V_4 this is immediate; in D_8 , squares and commutators kill its central involution in each proper noncentral normal subgroup; in E_8 , every noncentral normal subgroup has commutator with E_8 equal to its centre and has image of dimension at most three modulo that centre. These observations also suffice if the common image of N on an orbit is proper in H .

The remaining case is a subdirect $N \leq H^q$ with H critical. The block stabilizer induces the actual local automorphisms on every characteristic elementary section of H ; the sum of its conjugate sections is its induced module of index q . For C_2, V_4 the preceding module bound gives $q/2, q$. For D_8, E_8 , use both $N \cap Z(H)^q$ and its image in $(H/Z(H))^q$. Their fibre dimensions are respectively $(1, 2)$ and $(1, 4)$, so (9.1) gives $3q/2, 5q/2$, respectively. All four bounds are at most $3mq/8$. In particular the central line of a dihedral block has not been omitted.

If N is transitive, then

$$\rho_U(N) \leq d(N) \leq \left\lfloor \frac{\sqrt{3}w}{2\sqrt{\log_2 w}} \right\rfloor$$

by [Tra21, Theorem 1.1]. This is at most $3w/8$ for $w \geq 38$, and also for

$$w \in \{16, 19, 24, 27, 30, 32, 33, 35, 36\}.$$

The remaining finite input checks the whole heads in the complete transitive catalogues of degrees

$$2, 3, \dots, 23, 25, 26, 28, 29, 31, 34, 37.$$

Its only failures are the four critical actions. The catalogue classification is that of [HR20, Hul23]; this slice has 7,259 action classes. A separate enumeration of all 233 transitive normal pairs in degrees 2, 4, 8 shows that a critical transitive N has a head above $3w/8$ only when $U = N$ is the corresponding critical action. Thus the whole-head test is applied to the actual action of N , and the separate pair test handles its possible overgroups. The finite calculation constructs $N/(N^2[N, U])$ and verifies its elementary binary invariants; its precise bounded contract is recorded in Appendix A. Together with the preceding induction this proves the assertion in all degrees. $\square$

We next give the relative ternary bounds needed for the one-triple case. Let $\mathcal{E}_9$ consist of the four transitive actions $9T2, 9T6, 9T7, 9T17$, of orders 9, 27, 27, 81.

Lemma 9.2 (Ordered-transversal head bound). *Let P be a finite p -group, $H \leq P$, $[P : H] = p^t$, and let V be an a -dimensional representation of the original H over a field k . For every actual submodule $M \leq \text{Ind}_H^P V$,*

$$\dim_k \text{Hom}_P(M, k) \leq a \cdot \text{width}(\{0, \dots, p-1\}^t).$$

The order defining width is componentwise. For $p = 3$ its width is at most $W(t)$, where

$$W(0) = W(1) = 1, \quad W(2) = 3, \quad W(t) = 7 \cdot 3^{t-3} \quad (t \geq 3).$$

Proof. First choose an actual chain $H = H_0 \triangleleft H_1 \triangleleft \cdots \triangleleft H_t = P$ with each index p . For existence, a proper subgroup $L < P$ acts on P/L . The number of fixed cosets is positive and divisible by p , since $[P : L]$ is and all nontrivial orbit sizes are. A fixed coset gL satisfies $g^{-1}Lg \leq L$, hence equality by finiteness. Thus L is properly contained in its normalizer. An element of order p in $N_P(L)/L$ supplies the next subgroup of index p . Iteration and index multiplication give exactly t steps. Only consecutive normality is required; H need not be normal in P .

Choose $x_i \in H_i$ generating H_i/H_{i-1} , and set $T(b) = x_1^{b_1} \cdots x_t^{b_t}$ for $0 \leq b_i < p$. Successively extracting the highest digit proves that these are unique representatives for the original right cosets of H . Order the labels by $L(b) = \sum_i b_i p^{i-1}$, so the largest differing coordinate decides the comparison. If $d < b$ in this order and $b \leq c$ componentwise, then the coset of $T(d)T(b)^{-1}T(c)$ has label strictly below c . Indeed, at the largest differing coordinate i , the equal higher tails of d, b cancel. With $A, B, C \in H_{i-1}$ the remaining expression is

$$Ax_i^{d_i-b_i} B^{-1} C x_i^{c_i} S_c = D x_i^{d_i-b_i+c_i} S_c, \quad D \in H_{i-1}.$$

This uses only $H_{i-1} \triangleleft H_i$. The exponent satisfies $0 \leq d_i - b_i + c_i < c_i < p$, so no carry changes a higher coordinate. The lower coordinates absorb every commutator and stabilizer term.

Use the coinduced model

$$\{f : P \rightarrow V : f(hx) = \rho(h)f(x)\}, \quad (g \cdot f)(x) = f(xg).$$

It is the original induced representation: $g \otimes v$ corresponds to the function supported on Hg^{-1} with value v at g^{-1} ; the tensor relation gives precisely the displayed $\rho(h)$ twist. Values at the $T(b)$ give one copy of the original V per coset. For $q = T(b)^{-1}T(c)$, act by q^{-1} . Then

$$(q^{-1} \cdot f)(T(c)) = f(T(c)q^{-1}) = f(T(b)).$$

Thus the entire leading fibre vector moves unchanged from row b to row c . Every lower row moves to a row below c by the preceding calculation. Its original fibre twist is retained; it cannot create a higher row. In particular this argument does not require an invariant basis or a split fibre representation.

Choose an ordered basis of V and order full labels (b, j) by $aL(b) + j$, with coset position first. Work in the intrinsic quotient

$$Q = M / \langle g \cdot u - u : g \in P, u \in M \rangle_k.$$

Among lifts in M of all bases of Q , minimize the sum of their leading full-label heights. Such lifts exist, and the sum is a natural number. If two different lifts have leading labels $(b, j), (c, j)$ with $b \leq c$ componentwise, subtract from the second the scalar multiple of q^{-1} applied to the first which cancels its leading coefficient. The new lift has strictly smaller height: the leading fibre row is unchanged by transport, and every other term stays below (c, j) . In Q its class is the old second class minus that multiple of the first, so the classes still form a basis and the new lift is nonzero. This contradicts minimality. The case $b = c$ uses $q = 1$ and also excludes duplicate labels. Thus the leading labels inject into a disjoint copies of the grid, with an antichain in each copy. Invariant forms on M are exactly Q^* , proving the dimension bound. The quotient here is intrinsic to M , not its image in ambient coinvariants.

For three ternary coordinates the following seven chains partition all 27 labels:

$$\begin{aligned} &000, 100, 200, 210, 220, 221, 222; & 001, 101, 201, 211, 212; \\ &002, 102, 202; & 010, 110, 120, 121, 122; & 011, 111, 112; \\ &012; & 020, 021, 022. \end{aligned}$$

Fixing the other $t - 3$ coordinates gives 3^{t-3} slices, each met at most seven times by an antichain. For $t = 2$ use the chains 00, 10, 20, 21, 22; 01, 11, 12; and 02. The cases $t = 0, 1$ are immediate. $\square$

For a positive integer s , write $s_3 = 3^t$ and define the integer ternary envelope

$$B(s) = \min\{C(s), s/\text{lpp}(s/s_3)\}, \quad C(s) = \begin{cases} s, & t = 0, \\ s/3, & t = 1, 2, \\ 7s/27, & t \geq 3, \end{cases} \quad (9.3)$$

with $\text{lpp}(1) = 1$. We use this envelope directly in the relative-head recurrence.

Lemma 9.3 (Primitive inputs and block filtration). *For a primitive action R of degree r , let $a_3(R)$ count its composition factors of order three, with multiplicity. Then*

$$a_3(R) \leq r/3,$$

with equality only for the natural C_3, S_3 actions and the affine degree-nine actions of orders 216, 432. Outside these four actions, $a_3(R) < 3r/10$. Every primitive normal pair has $\delta_R(N) < 3r/20$, apart from the whole natural C_3, A_4 pairs. For a transitive A with a minimal block system of s blocks of size r , primitive block component R , top T , and block kernel K ,

$$\delta_A(N) \leq a_3(R)B(s) + \delta_T(NK/K). \quad (9.4)$$

Proof. The primitive generator bound [HRD13, Theorem 1.1] gives $\delta_R(N) \leq \lfloor \log_2 r \rfloor$; its exceptional natural S_3 case has zero relative ternary head. This proves the strict $3r/20$ bound for $r \geq 34$: check $34 \leq r \leq 63$, and then use $20j < 3 \cdot 2^j$ for $j \geq 6$. The composition-length bound [GPRV18, Theorem 1.3] gives $a_3(R) \leq (8/3) \log_2 r - 4/3 < 3r/10$ for $r \geq 45$. At 45 this follows from $45^{16} < 2^{89}$, and the difference is increasing thereafter. The bounded inputs enumerate all primitive normal pairs through degree 33 and all primitive composition series through degree 44. These are respectively 945 normal pairs in 253 actions and 336 actions, with exactly the exceptions stated above. The weaker $r/3$ assertion follows immediately, including those exceptions. Classification completeness uses [RD05, HRDR23]; the finite computations count order-three composition factors, rather than the ternary valuation of group order.

Here is the common ternary head bound used in the block argument. Write $s_3 = 3^t$. Restrict to a Sylow 3-subgroup and decompose the original induced module by its actual double cosets, retaining the twisted fibre actions. Its orbit indices are 3^j with $j \geq t$, and their sum is s . Filter the given submodule by the kernels of these actual coordinate maps. Head subadditivity and Lemma 9.2 bound each resulting head by $aW(j)$. For $t = 1, 2$, all $W(j) \leq 3^j/3$; for $t \geq 3$, all $W(j) = 7 \cdot 3^j/27$. Summation therefore gives $aC(s)$, including mixed orbit sizes. For $t = 0$ the ambient dimension as suffices. Restriction can only increase the invariant character head.

For a prime $q \neq 3$, restrict instead to a Sylow q -subgroup. Maschke's theorem makes every submodule of each induced component a direct summand. Its maps to the trivial line therefore extend to the component; Frobenius reciprocity bounds their dimension by a . Each actual orbit has size at least s_q , so there are at most s/s_q components. Maximizing s_q gives $as/\text{lpp}(s/s_3)$; at $s/s_3 = 1$ use the ambient dimension as instead. Together these give $aB(s)$. All local coefficients and their sums are integers; no floor is moved through the original fibre dimension.

For the block assertion, choose a chief series R_i of R and filter $N \cap K$ by the inverse images of R_i under all original conjugate block evaluations. These maps separate the block kernel. Each elementary layer is a submodule of the actual module induced from a block stabilizer. A ternary

layer of fibre dimension a_i therefore contributes at most $a_i B(s)$; a layer at another prime has zero ternary head.

For a nonabelian chief factor Q , each coordinate image of the corresponding layer is normal in the original quotient of R , hence either trivial or all of Q . Minimal normality makes Q perfect. Choose a maximal proper normal subgroup $M \triangleleft Q$. The ambient conjugates of the maps $Q \rightarrow Q/M$ are jointly injective, since their common kernel is ambient-normal and proper in Q . They give surjections onto simple perfect groups. Thus the original layer, including any proper subdirect correlations, is a subdirect product of simple perfect groups and is perfect: induct on the number of coordinates, whose successive kernels are either trivial or a full simple factor. Its ternary head is zero.

Sum this single chain using (9.1). The relative second isomorphism theorem identifies the final top section with NK/K under the original ambient conjugation. The ternary fibre dimensions sum to $a_3(R)$, proving (9.4). $\square$

Theorem 9.4 (Ternary relative stability). *Except for the whole natural C_3, A_4 pairs and the four whole $\mathcal{E}_9$ pairs, every transitive normal pair of degree w obeys*

$$\delta_A(N) \leq \lfloor 5w/27 \rfloor.$$

Consequently the bounds $w/4$ and $2w/9$ hold outside, respectively, the whole natural C_3 pair and the two whole natural pairs. Positive equality in the former occurs only at natural A_4 ; equality in the latter occurs precisely at the four whole $\mathcal{E}_9$ pairs. For an arbitrary $G \leq S_m$ with c actual regular C_3 orbits and a actual natural A_4 orbits,

$$9d_3(G) \leq 2m + 3c + a. \tag{9.5}$$

Proof. The finite primitive input in Lemma 9.3 already implies the assertion in the primitive case. The finite transitive normal-pair input in degree nine says that the only two-ninth equalities are the four whole $\mathcal{E}_9$ pairs and every other pair has head at most one. This is a complete finite catalogue, not an assumption on possible block systems.

From (9.3), $B(s) \leq s/3$ for $s \geq 3$ and

$$B(2), B(3), B(4), B(6), B(9), B(18) = 1, 1, 1, 2, 3, 6.$$

For $3 \nmid s$ and $s \geq 3$, the prime-to-three denominator is at least four; the other cases follow from $C(s)$. Use induction in (9.4). For an ordinary top with $s \geq 3$ and $r \geq 3$,

$$\delta_A(N) \leq rs/9 + 5s/27 \leq 5rs/27.$$

For $r = 2$ there is no ternary layer. At $s = 2$ the top head vanishes and $\delta_A(N) \leq a_3(R) \leq r/3 \leq 10r/27$. For a whole $\mathcal{E}_9$ top the bound is $3a_3(R) + 2 \leq r + 2 \leq 5r/3$ for $r \geq 3$; $r = 2$ is immediate. A natural A_4 top gives $a_3(R) + 1$, which is at most $20r/27$ for $r \geq 3$; again $r = 2$ is immediate. A natural C_3 top also gives $a_3(R) + 1$. Use the degree-nine input at $r = 3$, $a_3(R) \leq 1$ at $r = 4$, and $r/3 + 1 \leq 5r/9$ for $r \geq 5$; the binary-block case is immediate. These cases exhaust the induction. Integrality gives the stated floors.

The coarser conclusions now follow by comparing $5/27 < 2/9 < 1/4$ and inspecting the six stated exceptions. Apply the two-ninth bound to (9.2); a natural C_3 layer costs three and a natural A_4 layer one above twice its degree. This proves (9.5). In particular a no- C_3 , no- A_4 residual has $d_3(G) \leq 2m/9$; neither exclusion can be omitted. $\square$

Theorem 9.5 (Necessary high ternary actions). *If a transitive normal pair (A, N) of degree w satisfies $20\delta_A(N) > 3w$, its actual action is one of the following:*

1. natural C_3 or natural A_4 ;
2. $6T1, 6T4, 6T5, 6T6$;
3. $12T20, 12T85, 12T164, 12T194, 12T228, 12T229, 12T265$;
4. a transitive 3-group of degree 9 or 27.

This is a necessary list of actions, not a claim that all their normal pairs have high relative head.

Proof. Primitive actions are covered by Lemma 9.3. We retain the complete normal-pair catalogues in degrees 6, 12, 18: 1,300 actions and 20,410 pairs. Their strict high-action list is exactly items two and three, with no degree-eighteen hit. In particular every degree-eighteen pair has head at most two. These remain bounded classification inputs for the actual actions and all their normal subgroups.

Use a minimal block system and put $v = a_3(R)$ in (9.4). For a nonexceptional top with $s \geq 3$, $B(s) \leq s/3$. If $r \geq 5$, then

$$\frac{vs/3 + 5s/27}{rs} \leq \frac{1}{9} + \frac{5}{27r} \leq \frac{4}{27} < \frac{3}{20};$$

at $r = 4$, $v \leq 1$ gives $14s/27 < 3(4s)/20$. At $s = 2$ the top head vanishes and the stronger primitive density bound gives $v \leq 3r/10$ for $r \geq 4$, except possibly $r = 9$; the degree-eighteen input removes that exception.

For a natural C_3 top the bound is $v + 1$: the only $r \geq 4$ case not below $9r/20$ is $r = 4$, giving degree twelve. A natural A_4 top gives $v + 1 \leq 3r/5$ for $r \geq 4$. A whole $\mathcal{E}_9$ top gives $3v + 2 \leq 27r/20$ for $r \geq 6$, and $v \leq 1$ deals with $r = 4, 5$. For binary blocks $v = 0$, only the natural C_3 top can fail, giving degree six.

It remains to treat ternary blocks, $r = 3$. For a nonexceptional top,

$$B(s) + \lfloor 5s/27 \rfloor \leq \lfloor 9s/20 \rfloor \quad (s \notin \{2, 6, 18\}).$$

To check all indices, write $s = 3^t m$ with $3 \nmid m$. If $m \notin \{1, 2\}$, use $1/4 + 5/27 < 9/20$. If $m \in \{1, 2\}$ and $t \geq 3$, use $7/27 + 5/27 = 4/9 < 9/20$. For $t \leq 2$ the only remaining indices $s \geq 2$ are 2, 3, 6, 9, 18; the sums at $s = 3, 9$ are 1, 4 and satisfy the displayed floors. At $s = 18$, use the actual top bound two from the finite input, giving $B(18) + 2 = 8 < 81/10$. No separate top-degree 27, 54, 81, 162 refinement is needed.

The only child degrees still to inspect are 6, 9, 12, 18, 27. Degrees 6, 12, 18 use the retained finite catalogues. Degree nine is structural. Its minimal block system is 3×3 , and equality in the head recurrence forces both the primitive-fibre contribution and the original normal top head to equal one. A transitive three-point top is C_3 or S_3 . In the second case every permutation is conjugate to its inverse, so invariance sends each ternary character value to its negative; the relative ternary head is zero. The rank-one top is therefore C_3 . For the literal block kernel, the simultaneous fibre-sign map now gives the same dichotomy used below: a nontrivial sign annihilates its relative ternary head and would give $\delta_A(N) \leq 1$, while trivial signs make the kernel a 3-group. Thus the whole degree-nine action is a 3-group, without a finite order or action-list premise.

At degree 27 the same recurrence removes an apparent choice of block orientation. If r is the primitive block size, s the number of blocks, $d = \delta_A(N)$ and t the relative head of the original normal top image, then the high inequality and the stability envelope give

$$r = 3, \quad s = 9, \quad d = 5, \quad t = 2.$$

Indeed $rs = 27$ leaves $(r, s) = (3, 9)$ or $(9, 3)$. In the second case $v \leq 3$, $B(3) = 1$ and $t \leq 1$, so $d \leq 4$, a contradiction. In the first case $v \leq 1$, $B(9) = 3$ and $t \leq 2$; the high inequality forces equality throughout. The structural degree-nine argument just proved therefore makes the actual top a 3-group.

Let K be the literal block kernel and let $\epsilon : K \rightarrow C_2^9$ be the simultaneous vector of its nine fibre signs. Put $K_0 = \ker \epsilon$. Restriction to the original fibres injects K_0 into C_3^9 ; this retains every diagonal or proper subdirect correlation. If ϵ is nontrivial, top transitivity makes each coordinate sign nontrivial. Conjugation through that sign acts by inversion on the corresponding ternary coordinate. Consequently every original normal subgroup of K_0 , including every proper nonabelian subdirect core, has zero relative ternary head. The quotient K/K_0 is binary and has zero ternary head as well. Restriction to the literal kernel followed by the original top map now gives $\delta_A(N) \leq t = 2$, contradicting $20d > 81$.

If ϵ is trivial, $K = K_0$ is a 3-group. Since the actual top is also a 3-group, the original extension A is a 3-group, with no splitting assumption. Thus degrees nine and twenty seven enter the same existing 3-group owner.

The two smallest degrees are structural as well. At degree three, a nonzero relative character forces a quotient of order three; transitivity and the embedding in S_3 leave A_3 or S_3 . The latter conjugates every element to its inverse and hence annihilates every invariant ternary character, so only the natural $C_3 = A_3$ action remains. At degree four, the same quotient makes three divide the group order, while transitivity makes four divide it. A subgroup of S_4 therefore has order twelve or twenty four. The order-twelve subgroup has index two and is A_4 ; the order-twenty-four subgroup is S_4 , where conjugacy to inverses again annihilates the relative ternary head. Thus degree four is exactly the natural A_4 owner. The only genuinely residual bounded degrees are now 6 and 12. This exhausts the final case and proves the theorem. $\square$

For the separate bounded counting argument below, we retain the finite semiregular witnesses for all 91 nonsoluble degree-eighteen actions: each contains an actual fixed-point-free permutation of order three. Those witnesses are used for the stronger module-generator capacities in Theorem 15.9; they are not needed for the relative-head classification just proved.

10 Complete chief-layer fibres on one source

Throughout this section $Z_J(G) = \sum_{N \triangleleft G} |\text{Epi}(J, G/N)|$ retains every literal normal subgroup of the target. Put $h_b = b \log(b+1)$, with $h_0 = 0$.

Lemma 10.1 (Semisimple full fibre). *Let $E = S_1 \times \cdots \times S_t$ be a normal subgroup of an arbitrary finite group G , with each S_i nonabelian simple. Then, for every actual $J \leq S_b$,*

$$Z_J(G) \leq Z_J(G/E) \prod_{i=1}^t \left(1 + \frac{|\text{Aut } S_i| h_b}{\log |S_i|} \right).$$

The empty product is one. No permutation representation of G/E or of a simple section is required.

Proof. For an original $N \triangleleft G$, retain $I = N \cap E$ and $D = NE/E$. The subgroup I is an invariant subproduct. In G/I , the group N/I centralizes E/I and intersects it trivially. Projection from $C_{G/I}(E/I)$ to G/E is injective, since E/I is centreless. Thus I, D determine N whenever it exists.

Fix an onto map $\theta : J \rightarrow (G/E)/D$ and its actual kernel $F \leq S_b$. An onto lift restricts to an onto map $F \rightarrow E/I$. After that restriction is fixed, a lift is unique: two lifts have pointwise difference in E/I centralizing the onto image of F , and hence have trivial difference.

Distinct kernels of maps F onto one nonabelian simple S give a quotient S^k . Indeed, inductively a proper subdirect subgroup of $S^k \times S$ identifies the last coordinate with one of the preceding coordinates, and would give a repeated kernel. Therefore $k \log |S| \leq \log |F| \leq h_b$. Each kernel supports exactly $|\text{Aut } S|$ epimorphisms. Bound the maps to E/I by the product of these coordinate bounds, and then sum over the invariant subproducts I , enlarging to all subproducts. This gives the displayed product. All D and θ remain in the single $Z_J(G/E)$; no independent normal-menu multiplier is added. $\square$

Define the composition invariant

$$\kappa(G) = \sum_{C_p \text{ a composition factor of } G} \frac{\log p}{p},$$

with multiplicity. Nonabelian composition factors have weight zero.

Theorem 10.2 (Fixed-target composition envelope). *For every fixed finite G there are $C_G \geq 1$ and an integer $D_G \geq 0$ such that*

$$Z_J(G) \leq C_G(1 + h_b)^{D_G} 2^{\kappa(G)b} \quad (J \leq S_b).$$

One may take D_G to be the number of nonabelian composition factors. These constants are uniform on any fixed finite target menu.

Proof. Induct along a chief series, starting with $C_1 = 1, D_1 = 0$. For an elementary abelian minimal normal subgroup $E = \mathbb{F}_p^a$ put $Q = G/E$ and $q = \lfloor \log |Q| \rfloor$. For each original $N \triangleleft G$, put $I = N \cap E$, $D = NE/E$. A lift of an onto $\theta : J \rightarrow Q/D$ restricts on $F = \ker \theta$ to a map $F \rightarrow E/I$. There are at most $p^{ad_p(F)} \leq p^{ab/p}$ such restrictions. A fixed generating tuple of Q/D has length at most q ; choose its preimages in J . Their images, together with the restriction to F , determine the lift and have at most $|E|^q$ choices. Incompatible assignments contribute zero, so nonsplit extensions require no extra assertion. If $\nu(G)$ is the number of literal normal subgroups of G , then

$$Z_J(G) \leq \nu(G) |E|^q p^{ab/p} Z_J(Q).$$

Thus take $C_G = \nu(G) |E|^q C_Q$ and $D_G = D_Q$. For a semisimple minimal normal $E = \prod_i S_i$, use Lemma 10.1, taking $C_G = C_Q \prod_i (1 + |\text{Aut } S_i|)$ and $D_G = D_Q + t$. The abelian exponents add to $\kappa(G)$ by Jordan–Hölder. All permutation rank bounds were applied to the actual $F \leq S_b$. $\square$

Proposition 10.3 (Irreducible affine last layer). *Let $U = V \rtimes R$ be a fixed primitive affine action of degree $w = p^d$, with $R \neq 1$ faithful and irreducible on V . Write $\text{End}_R(V) = \mathbb{F}_{p^f}$ and $A_U = |Z^1(R, V)|$. Then*

$$Z_J(U) \leq (1 + A_U) 2^{fb \log p / (dp)} Z_J(R).$$

If R is nonabelian, then $f/d \leq 1/2$. In particular this proposition and Theorem 10.2 give a direct contractive row for every fixed finite menu satisfying

$$\kappa(R) + \frac{f \log p}{dp} < \frac{w - (w \bmod 2)}{8}.$$

Proof. Since $C_U(V) = V$, a normal subgroup disjoint from V is trivial; irreducibility makes every other normal subgroup contain V . Thus $Z_J(U) = Z_J(R) + |\text{Epi}(J, U)|$ exactly. Fix an onto $\theta : J \rightarrow R$ and $F = \ker \theta$. If V has multiplicity m in the semisimple head of $H_1(F, \mathbb{F}_p)$, then $dm \leq d_p(F) \leq b/p$ and the space of equivariant restrictions to V has size p^{fm} . Restriction of cocycles has kernel precisely the inflated $Z^1(R, V)$; each nonempty fibre therefore has size A_U . This proves the claimed bound, including every obstruction to a lift being onto. The integer f divides d . If $f = d$, the action is scalar over its commuting field, so R is abelian. Otherwise $f \leq d/2$. The last assertion follows from original-weight pointing and Lemma 7.2; all target constants are fixed. $\square$

Theorem 10.4 (Entire affine block kernel). *Let $U \leq S_w$ be transitive with $s \geq 2$ blocks of size $r = p^a \geq 2$, exact primitive affine component L , and faithful block action $T \leq S_s$. Put $w = rs$ and $g(w) = \lceil w/\sqrt{\log w} \rceil$. Choose a chief series $1 = N_0 < \dots < N_e = L$ and write $K_i = U \cap N_i^s$. For each abelian local chief factor $\mathbb{F}_{p_i}^{a_i}$, let H_i bound the number of module generators of every submodule of its actual induced ambient module and its dual. Then*

$$Z_J(U) \leq 2^{\eta_U b + \beta_U} \Psi_U(b) Z_J(T), \quad (10.1)$$

where

$$\begin{aligned} \eta_U &= \sum_i H_i \log(p_i)/p_i, \\ \beta_U &= \sum_i \log(p_i) [a_i s H_i + H_i w/p_i + a_i s (g(w) + 1)], \end{aligned}$$

and Ψ_U is the product in Lemma 10.1 over the simple factors of all actual nonabelian intersection layers. With $H_i \leq \lceil 4a_i s / \sqrt{\log s} \rceil$ these satisfy, uniformly as $rs \rightarrow \infty$,

$$\eta_U = o(w), \quad \beta_U = o(w^2), \quad \log \Psi_U(b) \leq Cw \log^2(w + b + 1).$$

All proper, zero and full block kernels, and all original normal quotients, occur in this inequality.

Proof. The endpoint of the local-chief filtration is the block kernel $K = U \cap L^s$, and the final quotient is T . Put $G_i = U/K_{i-1}$, $E_i = K_i/K_{i-1}$, $R_i = U/K_i$. For an abelian local factor, the ambient module $V_i = (\mathbb{F}_{p_i}^{a_i})^s$ factors through R_i : K_i centralizes the whole local abelian layer. It is induced from one block stabilizer of index s . The same is true of its dual. The local-chief construction is that of [Tra18, Lemma 5.8], with this explicit endpoint. For a nonabelian local factor, coordinate projections of E_i are normal in the exact L and hence are trivial or the full local chief factor. Transitivity makes the choice uniform. The subdirect description of products of nonabelian simple groups makes E_i semisimple, possibly trivial.

For one abelian layer abbreviate G, E, R, p, a, H accordingly. Duality gives, for any R -module X and every subquotient M of V_i ,

$$|\text{Hom}_R(X, M)| = |\text{Hom}_R(M^*, X^*)| \leq p^{H \dim X}.$$

Indeed a subquotient is a quotient of a submodule, so its dual requires at most H module generators. For $N \triangleleft G$ retain $I = N \cap E$ and $D = NE/E$. At most p^{asH} invariant submodules I occur, by padded generating tuples in V_i . For fixed I, D , put $M = E/I$. If a normal complement N/I exists, all others are its graphs of equivariant homomorphisms $D/[D, D]D^p \rightarrow M$. Here D acts trivially on M : its normal lift centralizes M . Thus this is an exact torsor when nonempty. The actual preimage $\tilde{D} \leq U \leq S_w$ gives $d_p(D) \leq d_p(\tilde{D}) \leq w/p$, so the graph count is at most $p^{Hw/p}$.

Fix now an onto $\theta : J \rightarrow R/D$ and its actual kernel $F \leq S_b$. Restrictions of compatible lifts to F have at most $p^{Hb/p}$ choices. Once a restriction is fixed, differences descend to cocycles of R/D on M . Every quotient of U has at most $g(w)$ generators, by [Tra18, Theorem 1.1(1)]; hence the lift torsor has size at most $|M|^{g(w)} \leq p^{asg(w)}$. This argument uses no splitting or cohomology vanishing. Summing I , the normal graphs, and the lifts leaves all D and θ in one $Z_J(R)$. It gives the stated abelian factor, even with the harmless extra $as \log p$ in β_U . Apply Lemma 10.1 at the other layers and iterate, always on the same literal J , to prove (10.1).

For completeness the universal induced-module bound [Tra18, Corollary 4.27(iii)] gives $H_i = \lceil 4a_i s / \sqrt{\log s} \rceil$ for $s \geq 1261$; below that threshold the dimension bound suffices since $\sqrt{\log s} < 4$. Put $L_0 = \log |L|$ and $q_s = \sqrt{\log s}$. The chief-order budget yields

$$\begin{aligned}\eta_U &\leq 2sL_0/q_s + L_0/2, \\ \beta_U &\leq 4s^2L_0^2/q_s + 2swL_0/q_s + swL_0/\sqrt{\log w} + 3sL_0 + wL_0/2.\end{aligned}$$

Since L is affine, $L_0 \leq \log r + (\log r)^2$. Divide the displayed bounds by rs and r^2s^2 , respectively. For large r their terms tend uniformly to zero; for bounded r they tend uniformly to zero as s increases. This proves both two-variable little-oh estimates, including the ceiling costs.

Every simple factor S in these layers has $\log |\text{Aut } S| \leq (\log |S|)^2$, since a generating tuple of length at most $\log |S|$ determines an automorphism. The total of $\log |S|$ is at most sL_0 , and each such term is at most L_0 . Therefore

$$\log \Psi_U(b) \leq sL_0^2 + 2sL_0 \log(b+1) \leq Cw \log^2(w+b+1).$$

The last bound follows since L_0/r and L_0^2/r are bounded for all $r \geq 2$. No faithful action was assigned to a simple section in deriving this estimate. $\square$

11 Uniform capacity of affine local components

The estimates here apply to the exact primitive local component of an imprimitive action, with arbitrary faithful transitive top.

Lemma 11.1 (Refined dual capacities). *For a module induced from a block stabilizer of index $s \geq 2$ with local dimension a in characteristic p , each of the following available integers bounds the number of generators of every submodule of the ambient module and its dual:*

$$\begin{aligned}\lfloor as/2 \rfloor, & \quad \left\lfloor \frac{as}{\text{lpp}(s/s_p)} \right\rfloor \quad (s/s_p > 1), \\ \left\lfloor as \sqrt{\frac{2}{3(p-1)e}} \right\rfloor & \quad (s_p = p^e, \ e > 0).\end{aligned}$$

If the block action contains a soluble transitive subgroup, two more bounds are $\lfloor a\tilde{\omega}(s) \rfloor$ and as_p , where

$$k(s) = \sum_{q|s} \nu_q(s)(q-1), \quad \tilde{\omega}(s) = \frac{s}{2^{k(s)}} \binom{k(s)}{\lfloor k(s)/2 \rfloor}.$$

The minimum of these bounds and $\lceil 4as/\sqrt{\log s} \rceil$ may be used in Theorem 10.4.

Proof. The prime-to- p and square-root bounds follow from [Tra18, Theorem 1.6, Corollary 4.27], using $\pi > 3$; the two conditional bounds are its soluble-transitive alternative. The dual is induced from the dual local module. If s is not a pure p -power, the prime-to- p denominator is at least two. If s is a p -power, a Sylow p -subgroup containing a Sylow subgroup of a point stabilizer is transitive, by orbit-stabilizer. Thus the conditional estimate applies. Since $k(s) \geq 1$ and $2^{-k} \binom{k}{\lfloor k/2 \rfloor} \leq 1/2$, it gives the same half bound. Integrality gives its floor. Finally the older uniform bound is valid by Theorem 10.4. Taking a minimum preserves every bound and its uniform error estimates. $\square$

Theorem 11.2 (All but seven affine local degrees). *In Theorem 10.4, suppose*

$$r \notin \{2, 3, 4, 5, 8, 9, 16\}.$$

Take the refined capacities of Lemma 11.1. With $h(w) = w - (w \bmod 2)$, they satisfy

$$\frac{h(w) - s}{8} - \eta_U \geq \frac{w}{1024}.$$

Consequently all these actions form one complete transfer family with an exponentially small ordinary kernel row and a quadratically small scalar, uniformly in both the local degree and block count.

Proof. Put $\gamma = (\log 3)/3 < 17/32$, the strict inequality following from $3^{32} < 2^{51}$. The half bound gives

$$\eta_U \leq \frac{s}{2} \kappa(L).$$

For every prime p , $\log(p)/p \leq \gamma$. The composition-length theorem [GPRV18, Theorem 1.3] gives

$$\kappa(L) \leq \gamma c(L), \quad c(L) \leq \frac{8}{3} \log r - \frac{4}{3}.$$

For $25 \leq r \leq 31$, the latter upper bound is below 12, so $c(L) \leq 11$ and $\kappa(L) < 187/32$. The margin is therefore greater than $5s/64 - (rs \bmod 2)/8$. If rs is even, this exceeds $rs/1024$. If it is odd, then $s \geq 3$ is odd, and the difference from $rs/1024$ exceeds

$$\frac{(80 - r)s - 128}{1024} \geq \frac{49 \cdot 3 - 128}{1024} > 0.$$

This retains the parity cost at the smallest odd endpoint.

For $r \geq 32$, use $s \geq 2$ to bound the margin below by

$$\frac{6r + 8 - 34 \log r}{48} s.$$

Subtract $rs/1024$. The resulting coefficient at $r = 32$ is $19/32$, and its derivative is positive for $r \geq 32$, since $\log_e 2 > 1/2$ gives

$$\frac{6 - 34/(r \log_e 2)}{48} - \frac{1}{1024} > \frac{6 - 68/r}{48} - \frac{1}{1024} > 0.$$

This proves the entire unbounded range analytically.

The remaining accepted prime powers are 7, 11, 13, 17, 19, 23. For prime local degree p , the stabilizer is cyclic of order dividing $p - 1$, all chief factors have dimension one, and

$$\eta_U \leq \lfloor s/2 \rfloor \kappa(L), \quad \kappa(L) \leq \log(p)/p + \kappa(C_{p-1}).$$

At $p = 7$, $\kappa(L) < 3/7 + 1/2 + 17/32 = 327/224$. For even s , the margin exceeds $9s/448 > 7s/1024$; for odd s , the retained floor gives $(9s + 271)/448$ instead. For a cyclic group, $\kappa(C_m) \leq \log(m)/2$. At $p = 11$, use $10^3 < 2^{10}$ to get $\kappa(L) < 13/6$; the margin is then greater than $s/6 - 1/8 \geq 5s/48$. At $p = 13$, $\kappa(L) < 5/2$ gives margin $s/4 - 1/8 \geq 3s/16$. Both exceed $ps/1024$. For $p \geq 17$, $\log(p-1) \leq (p-1)/4$ and $\log(p)/p < 1/2$, so the margin exceeds

$$s(p-5)/16 - 1/8 \geq s(p-6)/16 > ps/1024.$$

The last inequality holds already at $p = 17$. This exhausts the prime powers outside the seven stated degrees.

The actual quotient T acts on $s \leq h(w)/2$ points and $w \geq 14$. Choose one block and chief-series certificate per action class. Equation (10.1) retains $\beta_U = o(w^2)$ and $\log \Psi_U(b) \leq Cw \log^2(w+b+1)$. Theorem 7.4 applies with $\theta = 1/2$, $\rho = 1/1024$, $w_0 = 14$. In particular its complete kernel cells are eventually at most $2^{-wn/8192}$ and its scalar at most $2^{-n^2/8388608}$. The original action weight and all quotient and extension data are those already present in (10.1). $\square$

12 Explicit affine block counts

Write $s_p = p^{v_p(s)}$, let $\text{lpp}(t)$ denote the largest prime-power divisor of $t > 1$, and put

$$k(s) = \sum_q v_q(s)(q-1), \quad \omega(s) = s \binom{k(s)}{\lfloor k(s)/2 \rfloor} 2^{-k(s)}.$$

Use Lemma 11.1 for the induced module and its dual. In the notation of that lemma, write

$$H(s, p, a; T) = \min \left\{ \bar{H}, \left\lceil \frac{4as}{\sqrt{\log s}} \right\rceil \right\}, \quad (12.1)$$

where $\bar{H}$ is the minimum of all its available integer bounds. The soluble-transitive entries are always available at prime-power block counts.

Always use (12.1) in Theorem 10.4. Its upper cutoff retains the uniform $\beta_U = o(w^2)$ and the semisimple error estimate. Thus a uniform positive margin

$$\frac{h(w) - s}{8} - \eta_U \geq \rho w \quad (12.2)$$

is precisely an application of Theorem 7.4, with the actual top T as quotient. In the following arithmetic $\gamma = \log(3)/3 < 17/32$, using $3^{32} < 2^{51}$.

Theorem 12.1 (Regular prime and symmetric-three components). *For an actual nonbinary action with exact local C_p on p points, (12.2) holds with $\rho = 1/4096$ except possibly at the following block counts:*

$$\begin{array}{l|l} p \geq 5 & \emptyset \\ p = 3 & 2, 6 \\ p = 2 & \{2^a : 1 \leq a \leq 8\} \cup \{3 \cdot 2^a : 0 \leq a \leq 10\}. \end{array}$$

For exact natural S_3 , it holds with $\rho = 1/1024$ outside $\{2, 3, 4, 6, 8, 12, 18\}$.

Proof. For C_p the coefficient is $H \log(p)/p$. If $p \geq 5$, use $H \leq s/2$ and $\log(p)/p \leq 17/32$. Subtracting the desired larger margin $w/64$ leaves at least $((7p-25)s-8)/64 > 0$.

For $p = 3$, if s has a prime divisor at least five, use $H \leq s/5$; the margin exceeds $23s/160 - 1/8$. If $4 \mid s$, use $H \leq s/4$; the margin exceeds $15s/128$. These cases leave only $s = 3^b$ and $s = 2 \cdot 3^b$. At $s = 3$, $H \leq 1$ gives margin $> 3/32$. For $s = 3^b, b \geq 2$, the soluble bound gives $H \leq 3s/8$, hence margin $> 13s/256 - 1/8$. For $s = 2 \cdot 3^b, b \geq 2$, use $H \leq s/\sqrt{6} < 7s/16$, hence margin $> 9s/512$. All exceed $3s/4096$. Only two and six remain.

For $p = 2$, the margin is $s/8 - H/2$. If the odd part has a prime-power divisor at least five, $H \leq s/5$ leaves $s/40$. For $s = 2^a, a \geq 9$, the central-binomial ratio is at most $63/256$, leaving $s/512$. For $s = 3 \cdot 2^a, a \geq 11$, use $H \leq s\sqrt{2/33} < 127s/512$, leaving more than $s/1024$. This exhausts all counts outside the displayed binary list.

For S_3 , write $H_p = H(s, p, 1; T)$. Its coefficient is $H_2/2 + \gamma H_3$ and its reserve is $s/4 - (s \bmod 2)/8$. If s has a prime divisor at least five, $H_2, H_3 \leq s/5$ leave more than $7s/160 - 1/8 \geq 3s/160$. For $s = 2^a, a \geq 4$, use $H_2 \leq 3s/8, H_3 \leq 1$; the gap is greater than $s/16 - 17/32 > 3s/1024$. At $s = 9$, the integer bounds $H_2 \leq 1, H_3 \leq 3$ leave more than $1/32 > 27/1024$. For $s = 3^b, b \geq 3$, use $H_3 \leq 5s/16, H_2 \leq 1$; the gap exceeds $43s/512 - 5/8 > 3s/1024$. Finally let $s = 2^a 3^b, a, b \geq 1$. For $a \geq 3, \eta/s < 1/6 + 17/256$; for $a = 2, b \geq 2, \eta/s < 1/18 + 17/128$; for $a = 1, b \geq 3, H_2 \leq 2, H_3 \leq s/3$ give $\eta/s < 1/54 + 17/96$. Each is below $1/4 - 3/1024$. The untreated mixed cases are 6, 12, 18, proving the exact stated sufficient list. $\square$

Theorem 12.2 (Four-point component capacities). *For exact natural A_4 , (12.2) holds with $\rho = 1/4096$ outside $\{2, 3, 4, 6, 8, 12, 16, 24\}$. For exact natural S_4 it holds with the same ρ outside*

$$\{2^a : 1 \leq a \leq 8\} \cup \{3 \cdot 2^a : 0 \leq a \leq 10\} \cup \{5, 10, 15, 20, 30, 60\}.$$

These statements allow all proper, zero and nonsplit block kernels.

Proof. Put $H_{pa} = H(s, p, a; T)$. The respective coefficients are $\eta_A = H_{22}/2 + \gamma H_{31}$ and $\eta_S = \eta_A + H_{21}/2$, and the reserve is $3s/8$.

For A_4 , a prime divisor at least five gives $\eta_A/s < (1 + 17/32)/5 = 49/160$. Otherwise $s = 2^a 3^b$. For pure binary powers $a \geq 5, H_{22} \leq 5s/8, H_{31} \leq 1$ leave more than $s/16 - 17/32$. For pure ternary powers $b \geq 2, H_{22} \leq 2, H_{31} \leq 3s/8$ leave more than $45s/256 - 1$. For $a, b \geq 1$ with $a, b \geq 2$ use $\eta_A/s < 1/9 + 17/128$; for $a = 1, b \geq 2$ use $H_{31} \leq s/\sqrt{6} < 5s/12$, giving $\eta_A/s < 1/9 + 85/384$. Finally $b = 1, a \geq 4$ gives $\eta_A/s < 1/3 + 17/512$, whose gap is $13/1536$. All displayed gaps are at least $s/1024$, and their complement is exactly the asserted eight counts.

For S_4 , a prime divisor at least seven gives $\eta_S/s < 65/224$; a factor 25 gives the still smaller bound $65/800$. Write therefore $s = 2^a 3^b 5^c, c \in \{0, 1\}$. When $c = 1$, the prime-to-characteristic estimates give

$$\eta_S/s \leq \frac{3/2}{\max(3^b, 5)} + \frac{\gamma}{\max(2^a, 5)}.$$

For $a \geq 3$ this is less than $3/10 + 17/256$, leaving $11/1280$; for $b \geq 2$ it is less than $1/6 + 17/160$. The six complementary counts are 5, 10, 15, 20, 30, 60. When $c = 0$, pure binary powers $a \geq 9$ give $\eta_S \leq 189s/512 + \gamma$, leaving more than $3s/512 - 17/32 \geq s/1024$. Pure ternary powers $b \geq 2$ leave more than $45s/256 - 3/2 \geq s/1024$. For mixed counts, $b \geq 3$ gives $\eta_S/s < 1/18 + 17/64$, and $a, b \geq 2$ gives $\eta_S/s < 1/6 + 17/128$. At the remaining $s = 18$, the integer bounds $(H_{22}, H_{21}, H_{31}) \leq (4, 2, 7)$ leave more than $1/32 > 18/1024$. Finally $b = 1, a \geq 11$ gives

$$H_{22} \leq 2s\sqrt{2/33} < 79s/160, \quad H_{21} < 79s/320, \quad H_{31} \leq 3;$$

the gap exceeds $3s/640 - 51/32 \geq s/1024$ for $s \geq 6144$. The stated exceptional set is exactly the complement of these cases. $\square$

Theorem 12.3 (The four remaining affine local degrees). *For $r \in \{5, 8, 9, 16\}$, every action with exact primitive affine local degree r satisfies (12.2) with $\rho = 1/4096$, except possibly at*

r	s
5	2, 3, 4
8	2, 4, 6
9	2, 3, 4, 6, 12
16	2, 4.

Proof. Let $H(s, p, a)$ be the minimum in Lemma 11.1, using the soluble entries only for prime-power s and omitting the logarithmic cutoff. It has the form $\lfloor ax(s, p) \rfloor$, so $\sum_j H(s, p, a_j) \leq H(s, p, \sum_j a_j)$. Floors are taken after multiplying by the dimension. Put $t_2 = 1/2, t_3 = 17/32, t_5 = 7/15, t_7 = 3/7$, which upper-bound $\log p/p$. For brevity write $H_{pa} = H(s, p, a)$. The following upper bounds retain the true translation dimension:

$$\begin{aligned}
E_5 &= t_5 H_{51} + \frac{1}{2} H_{22}, \\
E_8 &= \frac{1}{2} H_{23} + t_3 H_{31} + t_7 H_{71}, \\
E_9 &= \frac{1}{2} H_{24} + t_3 (H_{32} + H_{31}), \\
E_{16} &= \frac{1}{2} H_{24} + \max_{\mathbf{n} \in V} \sum_{p=2,3,5,7} t_p H_{pn_p},
\end{aligned}$$

where $V = \{0 \leq \mathbf{n} \leq (6, 2, 1, 1) : \sum n_p \leq 5\}$. Indeed the stabilizers at degrees five and nine divide C_4 and $\mathrm{GL}_2(3)$, respectively. At degree sixteen the order $|\mathrm{GL}_4(2)| = 2^6 3^2 \cdot 5 \cdot 7$ gives the coordinate bounds, and [GPRV18, Theorem 1.3] gives at most five stabilizer composition factors after subtracting the four translation factors.

For degree eight, a proper subgroup of the simple group $\mathrm{GL}_3(2)$ has index at least seven, since an index below seven would embed the group in S_6 . Its order divides 168 and is at most 24, and therefore has at most two prime divisors and is soluble. An irreducible soluble stabilizer has trivial normal 2-subgroup. A minimal normal subgroup is then C_3 or C_7 ; C_3 has a nonzero proper fixed space on $\mathbb{F}_2^3$, contradicting irreducibility. A normal C_7 puts the stabilizer between C_7 and its order-21 Singer normalizer. Thus the only possibilities are $C_7, C_7 : C_3, \mathrm{GL}_3(2)$, proving the envelope E_8 . Nonabelian local factors in these arguments stay in Ψ_U .

Here is an exhaustive arithmetic verification for all s , without a group census. Independent upper bounds on weighted dimensions and a binary exponent cutoff are

r	A_2	A_3	A_5	A_7	a_0
5	1	0	7/15	0	5
8	3/2	17/32	0	3/7	3
9	2	51/32	0	0	4
16	9/2	17/16	7/15	3/7	5.

The reserve per s is at least $B_r = (r - 1)/8 - 1/16$ for odd r , and $B_r = (r - 1)/8$ for even r . If $5 \mid s$, the capacity per dimension at $p = 5$ is less than $5s/12$, and at the other primes at most $s/5$. If $7 \mid s$, use $s/3$ at seven and $s/7$ elsewhere. A prime divisor at least eleven gives $s/11$ everywhere. If $9 \mid s$, use $5s/12$ at three and $s/9$ elsewhere. If $2^{a_0} \mid s$, use $3s/8$ at two for $r = 5, 16$, $s/2$ for $r = 8$, and $5s/12$ for $r = 9$; other primes get $s/2^{a_0}$. Substitution gives these lower gaps per s ,

respectively:

r	$5 \mid s$	$7 \mid s$	$q \geq 11$	$9 \mid s$	$2^{a_0} \mid s$
5	31/720	383/1680	73/240	593/2160	23/480
8	429/1120	99/224	1605/2464	1181/2688	9/1792
9	7/32	95/224	215/352	59/1152	7/1536
16	2431/5040	209/240	23801/18480	50369/60480	6791/53760

Every entry exceeds $r/4096$. The complement is exactly $s = 2^a 3^b$, $0 \leq a < a_0$, $b \in \{0, 1\}$, $s \geq 2$. The accepted cells there, with exact gaps $(h(rs) - s)/8 - E_r$, are

5	$(6, 1/15), (8, 8/15), (12, 3/5), (16, 23/15), (24, 13/5), (48, 33/5)$
8	$(3, 37/224), (12, 363/224)$
9	$(8, 13/32), (24, 103/32)$
16	$(3, 17/32), (6, 17/16), (8, 3/2), (12, 9/2), (16, 3), (24, 9), (48, 18)$.

Direct substitution into the displayed integer formula proves each entry; each exceeds $rs/4096$. The thirteen remaining cells are exactly the exceptions in the theorem. Taking the logarithmic cutoff again restores the uniform error estimates and cannot worsen any margin. $\square$

Theorem 12.4 (Zero ternary kernel at local degree four). *Let U have exact local A_4 or S_4 on $s \geq 2$ blocks, put $M = (\mathbb{F}_2^2)^s$, $E = U \cap M$, and view $R = U/E$ inside $R_0 \wr T$, where $R_0 = C_3$ or S_3 . If $R \cap C_3^s = 1$, then $R = T$, counts two and three are impossible, and every count outside $\{4, 8, 16\}$ satisfies (12.2) with $\rho = 1/96$.*

Proof. For local A_4 , the linear block kernel is already $R \cap C_3^s$. For local S_4 , it embeds in C_2^s . Each of its coordinate images is a normal 2-subgroup of the exact S_3 , and is trivial. Thus $R = T$ in both cases. At counts two and three its point stabilizer cannot surject onto the required local C_3 or S_3 .

Only the translation intersection layer remains. With $H = H(s, 2, 2; T)$ its entire fibre is bounded by

$$Z_J(U) \leq 2^{Hb/2 + 4sH + 2s(g(4s) + 1)} Z_J(T).$$

If s is not a power of two, $H \leq 2s/3$ leaves gap $3s/8 - H/2 \geq s/24 = w/96$. If $s = 2^a$, $a \geq 5$, $H \leq 2\binom{a}{\lfloor a/2 \rfloor} \leq 5s/8$ leaves $s/16$. The other powers are two, four, eight and sixteen, of which two is impossible. The cutoff in (12.1) supplies the uniform quadratic error, so Theorem 7.4 applies. $\square$

13 Nonaffine primitive components and compressed quotients

Lemma 13.1 (Core-free subgroups of a simple power). *Let S be nonabelian simple, let ℓ be its least proper subgroup index, and let $H \leq S^a$ be core-free. Then $[S^a : H] \geq \ell^a$.*

Proof. We use $\ell^2 < |S|$, proved in [BBERJS22, Theorem A(4)]. Induct on a . Let $P \leq S^{a-1}$ and $B \leq S$ be the projections of H , with common Goursat quotient Q . If $B < S$, then $|Q| < |S|$. A full coordinate factor contained in P must map trivially to Q , and would therefore be contained in H , contrary to core-freeness. Thus P is core-free and the induction hypothesis gives $[S^a : H] \geq [S^{a-1} : P][S : B] \geq \ell^a$.

If $B = S$, then $Q = S$: otherwise the last factor is contained in H . Every full coordinate factor contained in P maps onto Q , so there is at most one, since images of distinct factors commute. If there is none, the index is $|S|[S^{a-1} : P] \geq \ell^a$. If there is one, write $P = S_i \times P'$; the map to Q is an isomorphism on S_i and trivial on P' , and P' is core-free in S^{a-2} . Hence $[S^a : H] = |S|[S^{a-2} : P'] > \ell^a$. This proves the induction. $\square$

Theorem 13.2 (Primitive compression). *If $L \leq S_r$ is nonaffine primitive, its socle E is semisimple and L/E has a specified faithful action of positive degree u with $2u \leq h(r)$. This assertion uses the finite primitive classification only in actual degrees $5 \leq r \leq 29$.*

Proof. Every nontrivial normal subgroup of a primitive group is transitive. An abelian minimal normal subgroup is regular and gives the affine case. Distinct minimal normal subgroups commute; since the centralizer of a transitive group is semiregular, two such subgroups must be regular mutual centralizers. They are isomorphic, and a third is impossible. Consequently $E = S^a$ for a nonabelian simple S . Also $C_L(E) = 1$: a nontrivial centralizer would contain a minimal normal subgroup of L , whereas its intersection with the centreless E is trivial. Thus

$$L/E \leq \text{Out}(S) \wr S_a.$$

Put $q = |\text{Out}(S)|$. Its regular action gives the wreath product a faithful action of degree aq , including $q = 1$.

Theorem A(4) of the cited paper gives $q \leq 3 \log \ell$, where $\ell \geq 5$ is the least proper subgroup index of S . Lemma 13.1 gives $r \geq \ell^a$. For $a \geq 2$, $2aq \leq \ell^a$: at $a = 2, \ell = 5$, integrality gives $q \leq 6$ and $4q \leq 24 < 25$; for $\ell \geq 6$ use $12 \log \ell \leq \ell^2$. Increasing a multiplies the right side by at least five and the left side by at most $3/2$. For $a = 1, \ell \geq 30$, use $6 \log \ell \leq \ell$ instead.

In the remaining almost simple case, failure of the regular action of the actual quotient to have $2|L/E| \leq r$ would imply $r < 2|L/E| \leq 6 \log \ell < 30$. The finite primitive certificate therefore checks precisely the required bounded range. Its complete degree-5–29 primitive alphabet has 116 simple nonabelian socles; each carries the literal socle and an actual regular quotient map with $2|L/E| \leq r$. Completeness is relative to the stated primitive classification [RD05, HRDR23]; the four nonsimple nonaffine socles in this range are covered by the preceding argument. Thus $2u \leq r$, and parity gives $2u \leq h(r)$. $\square$

Theorem 13.3 (Complete semisimple compression). *Suppose an actual transitive $U \leq S_w$, $w \geq 5$, has a normal semisimple subgroup E , possibly trivial, such that U/E has a faithful action of degree $v_0 \leq h(w)/2$. The complete family of these actions satisfies Theorem 7.4, with $\eta_U = 0$, $\theta = 1/2$ and $\rho = 1/20$. Every action with a nonaffine primitive exact local component has such a certificate.*

Proof. Write $E = \prod_i S_i$. Each S_i is a literal subgroup of S_w . A nontrivial orbit, followed by a maximal overgroup of its point stabilizer, supplies a faithful primitive action of S_i of degree at most w . Hence $d(S_i) \leq \log w$ by [HRD13, Theorem 1.1], and $|\text{Aut}(S_i)| \leq |S_i|^{\log w}$. Since $\sum_i \log |S_i| \leq \log(w!) \leq w \log w$, the product in Lemma 10.1 has logarithm at most $Cw \log^2(w+b+1)$ for an absolute C . The reserve is $(h(w) - v_0)/8 \geq h(w)/16 \geq w/20$ for $w \geq 5$. Theorem 7.4 applies with $\beta = 0$.

For a block system of size r and count s , let L be its exact primitive component and put $E_0 = \text{Soc}(L)$, $E = U \cap E_0^s$. Each coordinate projection of E is normal in the exact L , and hence is a normal subproduct of E_0 . The subdirect-product description of products of nonabelian simple groups makes E semisimple, allowing proper diagonal and zero intersections. Moreover U/E embeds into $(L/E_0) \wr T$, whose faithful action has degree $v_0 = us$. Theorem 13.2 gives $2v_0 \leq h(r)s \leq h(rs)$. This proves the last assertion, including primitive actions by taking $s = 1$. $\square$

14 Complete primitive-action consumers

Throughout this section $Z_J(U) = \sum_{N \triangleleft U} |\text{Epi}(J, U/N)|$. Every quotient is literal, and J is the same complete source throughout.

Lemma 14.1 (One derived source for soluble affine targets). *Let $U = V \rtimes R$, where $V = \mathbb{F}_p^d$ and $R \neq 1$ is soluble, faithful and irreducible. Put $\text{End}_R(V) = \mathbb{F}_{p^f}$, $c_R = |N_{\text{GL}_d(p)}(R)|/|C_{\text{GL}_d(p)}(R)|$, and $\eta = f \log(p)/(pd)$. For every $J \leq S_b$,*

$$Z_J(U) = Z_J(R) + |\text{Epi}(J, U)|, \quad |\text{Epi}(J, U)| \leq |V|c_R 2^{\eta b}.$$

More precisely, if t is the derived length of R and a_X are the multiplicities in the head of $H_1(J^{(t)}, \mathbb{F}_p)$ of the distinct pulled-back natural R -modules, then

$$|\text{Epi}(J, U)| = |V|c_R \sum_X (p^{f a_X} - 1). \quad (14.1)$$

Proof. Put $F = J^{(t)}$, $Q = J/F$. The nontrivial last derived group $R^{(t-1)}$ is abelian and has order prime to p , because $O_p(R) = 1$ for a faithful irreducible linear group. It has no fixed vector in V . For an onto $\theta : Q \rightarrow R$, the prime-to- p part of the abelian group $Q^{(t-1)}$ maps onto $R^{(t-1)}$. Averaging and inflation–restriction therefore give $H^i(Q, V_\theta) = 0$ for all $i \geq 0$. Restriction identifies $H^1(J, V_\theta)$ with $\text{Hom}_Q(H_1(F, \mathbb{F}_p), V_\theta)$, and each restriction has exactly $|V|$ literal cocycles. A nonzero restriction is onto V by irreducibility, and is exactly the condition that the lift to U is onto: $U^{(t)} = V$.

Two top maps give isomorphic modules precisely when they differ by an automorphism of R induced by $N_{\text{GL}_d(p)}(R)$. Each such orbit has c_R members, proving (14.1). All types lie in one head, so $d \sum_X a_X \leq d_p(F) \leq b/p$ and $\sum_X (p^{f a_X} - 1) \leq p^f \sum_X a_X - 1$. Finally, an original normal subgroup of U either contains V or intersects it trivially. In the latter case it centralizes V ; since $C_U(V) = V$, it is trivial. This proves the exact full-menu identity and the bound. $\square$

Theorem 14.2 (All sufficiently small-order actual actions). *All transitive nonbinary actions of degree $w \geq 64$ with $\log |U| \leq w/8$ have one complete ordinary forward kernel*

$$K_{\text{so}}(n, b) = \Delta(n, b) \sum_{\substack{[U]: \deg U = n-b \geq 64 \\ \log |U| \leq (n-b)/8}} \frac{2^{(\log |U|)^2 + (\log |U|)(b/2+1)}}{a(U)},$$

with row at most $2 \cdot 2^{-n}$ eventually and zero scalar. In particular this covers every primitive affine action of degree at least 1024.

Proof. The ordinary permutation-generator bound [HRD13, Proposition 2.1] gives $d(J) \leq b/2 + 1$. For $\ell = \log |U|$, the number of target normal subgroups is at most 2^{ℓ^2} , by padding a generating tuple of each subgroup to length $\lfloor \ell \rfloor$. The images of source generators then prove the displayed fibre envelope. Original pointing gives this kernel directly.

The uniform coefficient/Gaussian comparison gives

$$\log \Delta(n, b) \leq -\frac{(w - (w \bmod 2))b}{8} - \frac{w^2}{16} + O(w \log(n+2)).$$

The envelope costs at most $w^2/64 + wb/16 + w/8$. The action-class entropy is $o(w^2)$ by the transitive enumeration theorem. For $w \geq 64$, the negative terms are at most $-wn/32 - w^2/64$; uniformly absorb class entropy into $w^2/128 + O(w)$ and the remaining terms into $wn/64$. Hence $K_{\text{so}}(n, n-w) \leq 2^{-wn/64}$. Sum over all $w \geq 64$. For affine degree $w = p^d$, $\log |U| < (d+1) \log w \leq (\log w)^2 + \log w \leq w/8$ when $w \geq 1024$. The last inequality follows at $\log w = 10$ and thereafter by differentiating twice. A primitive affine group of that degree cannot be binary: in characteristic two an irreducible binary linear group has only the trivial simple module. $\square$

Theorem 14.3 (The soluble primitive affine range). *All soluble primitive affine actions of degrees $5 \leq w < 1024$, except the natural action $C_5 \rtimes C_4$ of degree five, have a complete ordinary forward row $O(2^{-an})$ and a complete scalar $O(2^{-a'n^2})$, for fixed $a, a' > 0$.*

Proof. For each actual $U = \mathbb{F}_p^d \rtimes R$, use Lemma 14.1. A regular prime action has $Z_J(C_p) = |\text{Hom}(J, C_p)| \leq p^{b/p}$; for $p \geq 5$, $\log(p)/p < (p-1)/8$. Suppose $R \neq 1$. If $p = 2$, $w \geq 8$, use the faithful action of R on the $w-1$ nonzero vectors. It has degree $v = w-1 < w$ and $\eta \leq 1/2 < w/8$, so Theorem 7.3 applies with comparator R .

For odd p and abelian R , irreducibility makes R cyclic in the multiplicative group of a field. Thus $Z_J(R) = |\text{Hom}(J, R)| \leq |J/J'| \leq 3^{b/3}$. At $w \geq 7$, its exponent and η are smaller than $(w-1)/8$. At degree five the complements are $1, C_2, C_4$. For C_2 use its degree-two comparator; the C_4 case is the specified exception.

At degree nine let $Z = R \cap \{\pm I\}$ and $P = R/Z$, acting faithfully on the four projective points. For each original normal subgroup of R , a fixed top map has at most $|\text{Hom}(J, C_2)|$ lifts through its central kernel. If $h_2(J) = |\text{Hom}(J, C_2)|$ and $e = d_2(P)$, then

$$h_2(J)Z_J(P) \leq 2^e Z_J(P \times C_2).$$

Indeed, for one quotient P/N with $c = |\text{Hom}(P/N, C_2)|$, the two product kernels contribute $[1 + h_2(J) - c]|\text{Epi}(J, P/N)|$, and $h_2(J) \leq 2^e[1 + h_2(J) - c]$ whenever the epimorphism count is nonzero. These kernels are distinct for each N . Consequently $Z_J(R) \leq \nu(R)2^{d_2(P)}Z_J(P \times C_2)$. The comparator has faithful degree six, smaller than $h(9) = 8$; the affine exponent is at most $\log(3)/3 < 1$.

At degree twenty-five the scalar subgroup of $R \leq \text{GL}_2(5)$ has order dividing four. Its projective image is a soluble subgroup of $\text{PGL}_2(5) \cong S_5$, of order at most twenty-four. An intransitive subgroup has this bound immediately; a soluble transitive subgroup of prime degree five has a regular normal C_5 and lies in its order-twenty normalizer. Each central quotient lift costs at most $|\text{Hom}(J, C_4)| \leq 2^{b/2}$. Counting all fixed target normals and using $d(J) \leq b/2 + 1$ gives $Z_J(R) \leq D_R 2^{(\log 48)b/2}$. Its exponent is strictly below $h(25)/8 = 3$.

At degree twenty-seven every soluble irreducible $R \leq \text{GL}_3(3)$ has order at most seventy-eight. To see this, its Fitting subgroup is a nontrivial nilpotent $3'$ -group. A nonhomogeneous restriction of the natural module has three one-dimensional homogeneous components, giving a monomial group of order at most forty-eight. In the homogeneous case the constituent has dimension three: a one-dimensional constituent would make the Fitting subgroup scalar and central, contradicting its self-centralizing property and irreducibility. Over the algebraic closure, a nilpotent $3'$ -group has no absolutely irreducible character of degree three, since its irreducible degrees are products of powers of its Sylow primes. Thus the faithful constituent splits into three Galois-conjugate linear characters. The Fitting subgroup is abelian and its field algebra is $\mathbb{F}_{27}$. Its normalizer lies in $\Gamma L_1(27)$, of order seventy-eight. Hence $Z_J(R) \leq D_R 2^{(\log 78)b/2}$, with $(\log 78)/2 < 26/8$.

In every other odd prime-power degree, the base-size theorem [HM16, Theorem 1.1] gives $|R| \leq w^2$ for $p \geq 5$ and $|R| \leq w^3$ for $p = 3$. Its complete-reducibility and p -solubility hypotheses follow from irreducibility and solubility here. The remaining ranges start at $w = 49$ and 81 . The inequalities $49^8 < 2^{48}$ and $81^{12} < 2^{80}$, and monotonicity of $w-1-8\log w$ and $w-1-12\log w$, give $\log |R|/2 < (w-1)/8$ in both ranges. The fixed-target ordinary generator envelope applies as before.

These cases exhaust the finite set of action classes. Every pointwise exponent has a strict physical gap; every comparator has $0 < v < h(w)$ and $\eta < h(w)/8$. Their original kernels are $\Delta(n, n-w)$ times their respective fibre envelopes divided by $a(U)$. Apply Theorem 7.3 and take

finite positive minima of these gaps. The pointwise classes have scalar zero. This proves complete row and scalar rates without assuming any normalized count is bounded. $\square$

Proposition 14.4 (Natural symmetric-four and Frobenius-five actions). *The natural S_4 and $C_5 \rtimes C_4$ actions have complete ordinary forward rows $O(2^{-an})$ and complete scalars $O(2^{-a'n^2})$, for positive constants.*

Proof. For $S_4 = \mathbb{F}_2^2 \rtimes S_3$, Lemma 14.1 gives $Z_J(S_4) \leq Z_J(S_3) + 24 \cdot 2^{b/4}$. The comparator S_3 has faithful degree three, less than the original width four, and $1/4 < 4/8$. The finite comparator theorem applies, using the original normalizer 24.

For the natural degree-five action put $z(J) = Z_J(C_4) = |\text{Hom}(J, C_4)|$ and $\eta = \log(5)/5$. The same lemma gives $Z_J(U) \leq z(J) + 5 \cdot 2^{\eta b}$. Declare $z(J) \leq 2^{3b/8}$ cold. Its exact kernel is

$$\frac{\Delta(n, n-5)}{20} \left(2^{3(n-5)/8} + 5 \cdot 2^{\eta(n-5)} \right),$$

whose gap is positive since $\eta < 15/32 < 1/2$. On the hot event, the whole original fibre is at most $6 \cdot 2^{(\eta-3/8)b} z(J)$. Use Theorem 26.1 with $r = \max(1, \lceil b/8 \rceil)$. Its weighted first moment, including the hot cutoff, has quadratic exponent

$$b^2/16 + br/4 + r^2/2 - (r-1)3b/8 + (\eta-3/8)b - n^2/16 = -b^2/128 + (r-b/8)^2/2 + (\eta-5/8)b - 25/16.$$

The actual scalar includes exactly the original factor $1/(20b!p_n G_{\lfloor n/2 \rfloor})$ on the hot source sum. Its logarithmic normalization costs $O(n \log(n+2))$; the moment error is $o(n^2)$. Thus it is at most $2^{-n^2/512}$ eventually. The hot event is empty at $b = 0$. $\square$

Theorem 14.5 (The nonsoluble primitive affine range). *All primitive affine actions $U = \mathbb{F}_p^d \rtimes R$ of degrees $5 \leq w = p^d \leq 1024$ with nonsoluble R have one complete pointwise ordinary forward row at most $2^{-n/8}$ eventually and scalar zero.*

Proof. By Theorem 10.2 and Proposition 10.3, the complete fibre has the bound

$$Z_J(U) \leq (1 + A_U) C_R (1 + b \log(b+1))^{D_R} 2^{(\kappa(R) + \eta_U)b},$$

where $A_U = |Z^1(R, \mathbb{F}_p^d)|$, $\eta_U = f \log(p)/(dp)$, and $\mathbb{F}_{p^f} = \text{End}_R(\mathbb{F}_p^d)$. Because R is nonabelian, f is a proper divisor of d ; otherwise the faithful image would be scalar over its commuting field. Thus $d \geq 2$ and $\eta_U \leq \log(p)/(2p)$. Write $\gamma = \log(3)/3 < 17/32$.

The primitive composition bound [GPRV18, Theorem 1.3] gives $c(U) \leq (8/3) \log w - 4/3$. Since $c(U) = d + c(R)$ and R has a nonabelian composition factor, the number of abelian composition factors of R is at most $(8/3) \log w - 7/3 - d$. Hence

$$\kappa(R) + \eta_U \leq \gamma((8/3) \log w - 11/6 - d) \leq \frac{17}{32}((8/3) \log w - 23/6).$$

For $x \geq 64$, the difference between $(x-1)/8$ and the last expression is increasing: using $\log_e 2 > 1/2$, its derivative exceeds $1/8 - 17/(6x) > 0$. At $x = 64$ it is $271/192 > 1/4$. Thus $h(w)/8 - \kappa(R) - \eta_U > 1/4$ throughout that range.

At $w = 32$, the integer composition bound gives at most six abelian factors in R , so the exponent is at most $6(17/32) + 1/4 = 55/16$, leaving gap $9/16$. At $w = 49$, use $7^{16} < 2^{45}$ to obtain at most ten abelian factors, giving exponent at most $357/64$ and gap $27/64$. At $w = 25$, quotient the scalar subgroup of order dividing four. The projective image is a nonsoluble subgroup

of $\mathrm{PGL}_2(5) \cong S_5$, hence A_5 or S_5 . For completeness, an intransitive subgroup of S_5 is soluble; a transitive subgroup has one or six Sylow-five subgroups. The former case lies in the order-twenty normalizer. In the latter, its order is 30, 60, or 120; order thirty would give an impossible faithful degree-four action of S_5 by cosets. The other two orders give A_5, S_5 . Therefore R has at most three abelian factors, all C_2 , so its exponent is at most $3/2 + 17/64 = 113/64 < 3 - 1/4$.

The only remaining prime-power degrees below sixty-four are 8, 16, 27: prime-degree stabilizers are cyclic, and $\mathrm{GL}_2(2), \mathrm{GL}_2(3)$ are soluble. The complete primitive action catalogue at these three degrees gives the following finite composition-factor data for the literal stabilizers:

degree	primitive indices	abelian factors of R	bound on $\kappa(R)$
8	3	none	0
16	11–20	at most one C_2 and one C_3	33/32
27	10,11	at most one C_2	1/2

The classification and representative conventions are those of [HRDR23, RD05]. The displayed composition series supply respectively exponent bounds $1/4, 41/32, 49/64$, all with physical gap greater than $1/4$. The catalogue completeness is used only at these three degrees; the other ranges were covered analytically.

Finally use the original pointed kernel

$$\Delta(n, b) \sum_{\substack{[U] \text{ in this family} \\ w=n-b}} \frac{(1 + A_U)C_R(1 + b \log(b+1))^{D_R(\kappa(R)+\eta_U)b}}{a(U)}.$$

The family is fixed and finite, so all target constants and polynomial exponents are fixed. Its complete row is $O(n^D)2^{-n/4} \leq 2^{-n/8}$ eventually. The full cocycle factor A_U remains in this argument; it is not set equal to $|\mathbb{F}_p^d|$ for nonsoluble stabilizers. $\square$

15 The bounded ternary block counts

Put $\gamma = (\log_2 3)/3 < 17/32$. All fibres in this section use the complete source $J \leq S_b$, without any restriction on its orbits.

Lemma 15.1 (Finite multiplicative comparison). *Fix finitely many actual actions U of widths w , and fixed finite comparators Q_U with faithful actions of degrees $0 < v_U < h(w)$. Suppose*

$$Z_J(U) \leq C_U 2^{\eta_U b} Z_J(Q_U), \quad \delta_U = (h(w) - v_U)/8 - \eta_U > 0.$$

Their complete pointed union has an exponentially small forward row and a quadratically small scalar, with original normalizers $a(U)$. The comparators need not be quotients of U .

Proof. Set $\alpha = v/8 + \delta/2$, and call J hot when $Z_J(Q) > 2^{\alpha b}$. The cold coefficient is exactly $\Delta(n, b)C_U 2^{(\eta_U + \alpha)b}/a(U)$. Its exponent is $-\delta b/2 - w^2/16 + O_w(\log(n+2))$. For the full hot fibre use Lemma 7.1 with $q = \max(1, \lceil 4\delta b/v^2 \rceil)$ and $q_* = 4\delta b/v^2$. The normalized leading exponent is

$$\frac{(b+qv)^2}{16} - (q-1)\alpha b + \eta b - \frac{(b+w)^2}{16} = -\frac{\delta^2 b^2}{v^2} - \frac{\delta b}{2} - \frac{(w-h(w))b}{8} - \frac{w^2}{16} + \frac{v^2(q-q_*)^2}{16}.$$

The rounding term is bounded and the coarse counting error is $O(n^{3/2} + n \log(n+2))$. The original hot weight is $(b!a(U)p_n G_{\lfloor n/2 \rfloor})^{-1}$ times the complete hot sum of $Z_J(U)$. A finite union therefore preserves both strict losses. Choose certificates before the any-hot/cold partition. $\square$

Lemma 15.2 (Central binary comparison). *If X has central subgroup C_2 and quotient T , then*

$$Z_J(X) \leq \nu(X) |\operatorname{Hom}(J, C_2)| Z_J(T), \quad |\operatorname{Hom}(J, C_2)| Z_J(T) \leq 2^{d_2(T)} Z_J(T \times C_2),$$

where $\nu(X)$ is its number of literal normal subgroups.

Proof. For each $N \triangleleft X$, X/N is a central extension of a quotient of T by a group of order at most two. A fixed map to that quotient has no lifts or at most $|\operatorname{Hom}(J, C_2)|$ lifts. Summing over all N proves the first inequality, including repetitions of the same normal subgroup of T . For the second put $h = |\operatorname{Hom}(J, C_2)|$. For each $S \triangleleft T$ put $Q = T/S$, $a = |\operatorname{Hom}(Q, C_2)|$. An onto map $J \rightarrow Q$ and a binary character are jointly onto $Q \times C_2$ precisely when that character does not factor through the map. Thus the distinct product normals $S \times 1, S \times C_2$ contribute $(h - a + 1) |\operatorname{Epi}(J, Q)|$. Whenever the latter epimorphism set is nonempty, $h \geq a$ and $h \leq a(h - a + 1) \leq 2^{d_2(T)}(h - a + 1)$. All remaining product normals add nonnegative terms. $\square$

Proposition 15.3 (Regular ternary exceptions). *Every actual transitive action with exact local regular C_3 and two or six blocks has a complete exponentially small forward row and quadratically small scalar.*

Proof. Write $U \leq C_3 \wr T$, $E = U \cap C_3^s$. At $s = 6$, every transitive $T \leq S_6$ contains a semiregular C_3 : a Sylow three-subgroup of order three has trivial point stabilizers, while one of order nine is a Sylow subgroup of S_6 and contains two disjoint three-cycles. The ambient module and its dual restrict to two regular $\mathbb{F}_3[C_3]$ modules. Since $\mathbb{F}_3[C_3] \cong \mathbb{F}_3[t]/(t^3)$, every submodule has at most two generators: its number of Jordan blocks is its socle dimension, at most two. Quotients retain this bound. The full-chief proof of Theorem 10.4, with this capacity, gives $Z_J(U) \leq C_U 2^{2\gamma b} Z_J(T)$. Its physical width is eighteen, its comparator degree six, and $3/2 - 2\gamma > 0$, so Lemma 15.1 applies.

At $s = 2$, exact local C_3 forces E to project onto each coordinate. Its only C_2 -invariant possibilities in $\mathbb{F}_3^2$ are the diagonal, the antidiagonal, and the full plane. Coprimality splits the extension, giving $C_6, S_3, C_3 \times S_3$, respectively, all in their original degree-six actions. The first has full fibre $|\operatorname{Hom}(J, C_6)| \leq 3^{b/3}$. For the other two use the common source $F = J'J^2$, $B = J/F$. The semisimple module $H_1(F, \mathbb{F}_3)$ has trivial multiplicity $d = d_3(J)$ and nontrivial sign multiplicities a_θ , with $d + \sum a_\theta \leq b/3$. Coprime inflation–restriction gives

$$|\operatorname{Epi}(J, S_3)| = 3 \sum_{\theta \neq 0} (3^{a_\theta} - 1), \quad |\operatorname{Epi}(J, C_3 \times S_3)| = (3^d - 1) 3 \sum_{\theta \neq 0} (3^{a_\theta} - 1).$$

The second joint map is onto because C_3, S_3 have no common nontrivial quotient. Each expression is at most $3^{b/3+1}$. The complete normal quotients of $C_3 \times S_3$ are $1, C_2, C_3, C_6, S_3, C_3 \times S_3$; the cyclic terms have the same abelianization bound. Thus $Z_J(U) \leq C_U 2^{\gamma b}$. Its slope is less than $h(6)/8 = 3/4$, so direct normalization already gives an exponentially small row and zero scalar. $\square$

15.1 Signed modules on a bounded number of triples

For exact natural S_3 on s blocks write

$$E = U \cap C_3^s, \quad R = U/E \leq C_2 \wr T, \quad D = \ker(R \longrightarrow T) \leq C_2^s.$$

These are the actual kernel and actual top of the chosen block system. If $E = 1$, the whole block kernel is trivial: a nontrivial normal local image is C_3 or S_3 , and in the latter case its derived

subgroup projects onto C_3 . Hence $U \cong T$ has faithful degree s and satisfies Theorem 7.3. If an ambient signed ternary module and its dual have capacity H_3 , the full-chief argument gives

$$Z_J(U) \leq C_U 2^{\gamma H_3 b} Z_J(R). \quad (15.1)$$

Adding a binary capacity H_2 gives the same bound with $2^{(\gamma H_3 + H_2/2)b} Z_J(T)$. These bounds include all proper kernels, normal graphs and nonsplit extensions.

Lemma 15.4 (Signed isotypes). *Assume $D \neq 1$. Its coordinate characters are all nonzero and have q distinct values, each repeated $a = s/q$ times. They span D^* ; thus $q \mid s$, $d = \dim D \leq q \leq 2^d - 1$, and $q = 1, 2$ forces $d = 1, 2$, respectively. If a is a power of two or $a = 3$, every submodule of the signed ambient module and of its dual is cyclic.*

Proof. Transitivity makes the character multiplicities equal, and their common kernel is zero because D is its actual coordinate image. The idempotents of $\mathbb{F}_3[D]$ decompose a submodule into isotypes. The stabilizer of one isotype acts transitively on its a coordinates by signed permutations. If a is a power of two, a transitive Sylow two-subgroup of its coordinate image lifts to a two-group of signed permutations. The coordinate module is cyclic and semisimple over this group; an equivariant projection of a cyclic generator generates any submodule. If $a = 3$, a Sylow three-subgroup maps isomorphically onto a coordinate three-cycle. Its signs have product one, so it untwists to the regular $\mathbb{F}_3[C_3]$ module, whose ideals are cyclic. Transport a generator of the chosen isotype by R to generate all isotypes. Signed permutation matrices identify the dual with the original module. Quotients of cyclic modules are cyclic as well. $\square$

Lemma 15.5 (Two common-source onto bounds). *(i) Let V be elementary abelian of order 3^e , and R a two-group acting faithfully with cyclic dual V^* . Then*

$$|\text{Epi}(J, V \rtimes R)| \leq 3^e |\text{GL}_e(3)| 3^{b/3}.$$

The same conclusion holds for $R = C_6$ acting faithfully on a single unipotent ternary block, with its involution acting as -1 .

(ii) If X has a normal Sylow p -subgroup P with cyclic center and $C_X(P) = Z(P)$, then

$$|\text{Epi}(J, X)| \leq |Z(P)| |\text{Aut}(P)| p^{b/(p-1)}.$$

Proof. For (i), fix $F = O^2(J)$ in the two-group case and $F = J'$ in the C_6 case. Formation residuals commute with onto maps: the image of the source residual has the required quotient type, and the preimage of the target residual has that type, proving both inclusions. Consequently every map under consideration sends this same F onto V . Put $B = J/F$, $A = H_1(F, \mathbb{F}_3)$. The restriction is an onto B -map $A \rightarrow V_\theta$ and dualizes to an injection $\iota : V_\theta^* \rightarrow A^*$. Fix a cyclic vector $v_0 \in V^*$. Its image f determines the whole submodule $\langle Bf \rangle$. For each f , at most $|\text{GL}_e(3)|$ identifications with V^* are possible, and faithfulness determines the quotient map θ uniquely from each identification. Thus all quotient maps and onto restrictions together have at most $|\text{GL}_e(3)| 3^{\dim A}$ choices. For fixed restriction the lift fibre is empty or a cocycle torsor on B . Averaging gives $H^1(B, V_\theta) = 0$ when B is a two-group. In the second case B is abelian and its two-primary Hall subgroup has zero invariants because it maps onto the scalar involution; averaging and inflation-restriction give the same conclusion. The torsor has size at most $|V|$. Finally $\dim A = d_3(F) \leq b/3$.

For (ii), fix $F = O^{p'}(J)$ and its largest p -group quotient $Q = F/O^p(F)$. Every onto map to X sends F onto P and factors there through Q . Since $F \leq S_b$, $|Q| \leq p^{v_p(b!)} \leq p^{b/(p-1)}$. A faithful linear character of $Z(P)$ induces an irreducible constituent faithful on P : every nontrivial normal

subgroup of a p -group meets its center. Fix its faithful character χ . Pullback along onto maps $Q \rightarrow P$ gives irreducible characters of Q ; one pulled-back character determines the kernel and therefore at most $|\text{Aut}(P)|$ maps. The degree-square identity bounds their number by $|\text{Aut}(P)||Q|$. For a fixed onto restriction, two extensions to J differ in $C_X(P) = Z(P)$, so their quotient maps coincide. Their differences form a cocycle torsor on the p' -group J/F with coefficients $Z(P)$. Coprime averaging bounds its size by $|Z(P)|$. $\square$

Proposition 15.6 (Two symmetric triples). *All exact-local- S_3 actions on two blocks satisfy a complete finite forward estimate.*

Proof. The block stabilizer is its kernel, so its binary image is either C_2^2 or diagonal C_2 . In the first case distinct signs force $E = \mathbb{F}_3^2$ and $U = S_3 \wr C_2$. The ternary kernel is faithful irreducible for $R = D_8$, and Lemma 14.1 gives $Z_J(U) \leq Z_J(D_8) + 216 \cdot 3^{b/6}$. The comparator degree is four. In the diagonal case R is C_4 or V_4 . Coprimality splits the extension. For C_4 , its square is $-I$, so $E = \mathbb{F}_3^2$ is irreducible and the derived-source bound gives $Z_J(U) \leq Z_J(C_4) + 18 \cdot 3^{b/3} \leq 19 \cdot 3^{b/3}$. For V_4 , the ambient module has two distinct sign lines; choosing one or both gives $S_3 \times C_2$ or S_3^2 . The former has a faithful degree-five action.

For the latter use the same $F = J'J^2$ and sign multiplicities a_θ as above. An ordered pair of onto maps to S_3 is onto S_3^2 precisely when its binary characters differ: a proper subdirect has common quotient C_2 or S_3 , forcing equality. Hence

$$|\text{Epi}(J, S_3^2)| = 9 \sum_{\theta \neq \psi} (3^{a_\theta} - 1)(3^{a_\psi} - 1) \leq 18 \cdot 3^{b/3}.$$

The last inequality expands $\prod_\theta (1 + (3^{a_\theta} - 1))$ and uses $\sum a_\theta \leq b/3$. Its complete normal menu is

$$Z_J(S_3^2) = |\text{Epi}(J, S_3^2)| + 2|\text{Epi}(J, S_3 \times C_2)| + 2|\text{Epi}(J, S_3)| + |\text{Epi}(J, V_4)| + 3|\text{Epi}(J, C_2)| + 1.$$

Indeed intersection with C_3^2 is zero, one of its two sign lines, or the full plane; self-centralization resolves the zero case, and the other cases give exactly these quotients. Thus $Z_J(S_3^2) \leq 2Z_J(S_3 \times C_2) + 18 \cdot 3^{b/3}$. Every slope is below $h(6)/8$ and every comparator degree is below six, so Theorem 7.3 finishes all cases. $\square$

Proposition 15.7 (Four blocks with two signs). *All exact-local- S_3 four-block actions whose binary kernel has two distinct coordinate characters have a complete finite forward estimate.*

Proof. Their equal character classes have size two, $D = C_2^2$, and the top preserves the two-class partition. Thus $T \leq D_8$ has order four or eight, and R is a two-group of order sixteen or thirty-two. Coprimality splits $U = E \rtimes R$. The two isotypes have equal dimension in every submodule, so nonzero E has dimension two or four. Every two-dimensional section has monomial image exactly $D_8 \leq \text{GL}_2(3)$: its diagonal signs are independent and its two lines are interchanged. Its action kernel K in R has order two or four. Such a normal kernel has a central series of at most two order-two factors, first by meeting $Z(R)$ and then because a normal subgroup of order two is central.

Put $P = \mathbb{F}_3^2 \rtimes D_8 = S_3 \wr C_2$ in its faithful degree-six action and $Q_* = P \times C_2^2$ in degree ten. The central series, Lemma 15.2, and the literal quotient-menu inclusion imply $Z_J(V \rtimes R) \leq A_V Z_J(Q_*)$ for every two-dimensional section V , with a fixed finite A_V . They also give $Z_J(R) \leq A_R Z_J(Q_*)$ whenever such a section exists. No splitting of the central extensions is required.

If E is the full four-dimensional module, it is faithful and its dual is cyclic: the coordinate module is cyclic and semisimple for the two-group R . Lemma 15.5(i) bounds its onto maps by

$C_0 3^{b/3}$, where $C_0 = 81|\mathrm{GL}_4(3)|$. A normal subgroup disjoint from E centralizes E , so is trivial. All other proper intersections are two-dimensional sections. Consequently

$$Z_J(U) \leq |\mathrm{Epi}(J, U)| + Z_J(R) + \sum_{\substack{I < E \\ \dim I = 2, I \text{ } R\text{-stable}}} Z_J(U/I).$$

If E is reducible, the preceding comparisons bound this by $B_U Z_J(Q_*) + C_0 3^{b/3}$. If it is irreducible, the sum is empty and use R in faithful degree eight instead. If $\dim E = 2$, the central comparison already bounds the entire fibre. These are all normal strata. Comparator degrees at most ten and slope γ are strictly below the required width-twelve thresholds. $\square$

Proposition 15.8 (Three blocks with one sign). *All exact-local- S_3 three-block actions with diagonal binary kernel have a complete finite forward estimate.*

Proof. The top is C_3 or S_3 . A lifted coordinate three-cycle of order three untwists its signs. In the first case $R = C_2 \times C_3$. In the second the determinant is nontrivial on the scalar binary kernel and splits $R \rightarrow S_3$; a transposition conjugating the three-cycle to its inverse has constant signs. Thus after diagonal rescaling $R = C_2 \times S_3$ acts by scalar signs and ordinary coordinate permutations. Its nonzero invariant ternary submodules are exactly the three ideals in $\mathbb{F}_3[t]/(t^3)$, of lengths $e = 1, 2, 3$. All are stable under S_3 as well. The scalar involution has zero invariants on the module and on every quotient. Averaging and inflation–restriction give $H^1(R, \mathbb{F}_3^3/E) = 0$; a translation therefore removes the cocycle of the actual embedded subgroup. Hence $U_e = E_e \rtimes R$.

For top C_3 , $U_1 = S_3 \times C_3 =: Q$, faithfully of degree six. At $e = 2, 3$, the action is faithful, $U'_e = E_e$, and the dual is cyclic. Lemma 15.5(i) gives $|\mathrm{Epi}(J, U_e)| \leq C_e 3^{b/3}$. Normal intersections with E_e form its uniserial chain. At surviving length at least two, faithfulness forces a normal subgroup disjoint from the ternary kernel to be trivial. At length one the quotient is Q , whose normals disjoint from its first ternary line are 1 and its central C_3 . There are no mixed normal graphs, because the involution inverts one line and fixes the other. Thus, exactly,

$$\begin{aligned} Z_J(U_1) &= Z_J(Q), \\ Z_J(U_2) &= Z_J(Q) + |\mathrm{Epi}(J, U_2)|, \\ Z_J(U_3) &= Z_J(Q) + |\mathrm{Epi}(J, U_2)| + |\mathrm{Epi}(J, U_3)|. \end{aligned}$$

For top S_3 , $U_1 = S_3^2 =: Q_*$, again faithfully of degree six. At lengths two and three the action of R is faithful. The normal Sylow three-subgroup $P_e = E_e \rtimes C_3$ has center the first ideal C , of order three, and $C_{U_e}(P_e) = C$. This follows since centralizing E_e forces an element into E_e , where centralizing the top three-cycle forces it into C . Lemma 15.5(ii) gives $|\mathrm{Epi}(J, U_e)| \leq 3|\mathrm{Aut}(P_e)|3^{b/2}$. Every nontrivial normal subgroup contains C , by the uniserial chain and faithfulness. Moreover $U_2/C \cong Q_*$ and $U_3/C \cong U_2$. To see the necessary twist, the length-one quotient is the sign representation of S_3 , while $(\mathbb{F}_3^3/C)^*$ is its augmentation module. The dual of a two-dimensional module is its determinant-inverse twist; here the determinant is the sign of the S_3 factor. Multiplying the scalar C_2 coordinate of R by this sign is an automorphism of R and supplies both asserted abstract identifications. Consequently the same three displayed identities hold with Q_* in place of Q and these new onto tails. Both comparators have degree six, physical $h(9) = 8$, and the largest tail slope $(\log_2 3)/2$ is less than one. Apply Theorem 7.3. $\square$

Theorem 15.9 (All bounded symmetric-three counts). *Every actual action with exact local natural S_3 and $s \in \{2, 3, 4, 6, 8, 12, 18\}$ blocks has a complete exponentially small forward row and a quadratically small scalar. Together with Theorem 12.1, this covers every imprimitive exact-local- S_3 action.*

Proof. The $E = 1$ case was disposed of above. When $D = 1$, use (15.1) with comparator T and the respective capacities

s	3	4	6	8	12	18
H_3	1	1	2	1	3	7.

These follow from Lemma 11.1; at six use the semiregular three-cycle argument of Proposition 15.3, lifting through a binary kernel by a Sylow three-subgroup. Every margin $(h(3s) - s)/8 - \gamma H_3$ is positive. At $s = 2$, $D = 1$ contradicts the exact local S_3 image; Proposition 15.6 covers that count completely.

At $s = 8$, irrespective of D , the prime-to-three bound is $H_3 = 1$; retain R in degree sixteen, with margin $1 - \gamma > 0$. At $s = 18$, a soluble transitive subgroup of T gives $H_3 \leq 5$ and $H_2 \leq 2$, so the degree-eighteen comparator T has margin $7/2 - 5\gamma > 0$. Otherwise use the actual semiregular C_3 witness for every nonsoluble degree-eighteen action recorded in the finite input to Theorem 9.5. Its lift through D has order three. The signed ternary module and its dual untwist to six regular modules over $\mathbb{F}_3[C_3]$, so $H_3 \leq 6$. The same comparator has margin $7/2 - 6\gamma > 0$.

Now assume $D \neq 1$ and use the actual isotype count q . At $s = 12$, if $q \leq 2$, then $d = q \leq 2$ and $H_3 \leq 3$. For each literal normal quotient of R , a fixed map to its top quotient has at most $2^{db/2}$ restrictions into its binary kernel, times a fixed cocycle constant. Thus $Z_J(R) \leq C_R 2^{db/2} Z_J(T)$, including all normal graphs. The margin is at least $2 - 3\gamma > 0$. If $q \geq 3$, its isotype size is 4, 3, 2 or one; Lemma 15.4 gives $H_3 = 1$ and the degree-twenty-four comparator R has margin $3/2 - \gamma > 0$. At $s = 6$, $q \geq 2$ similarly gives isotype sizes 3, 2, 1 and margin $3/4 - \gamma > 0$ with degree-twelve comparator R .

At $s = 4, 6$, the remaining $q = 1$ means that D is central of order two in R . Combine Lemma 15.2 with the capacities $H_3 = 1, 2$. The comparator $T \times C_2$ has degrees six and eight; its margins are respectively $3/4 - \gamma > 0$ and $5/4 - 2\gamma > 0$. The $s = 4, q = 2$ branch is Proposition 15.7. For $s = 4, q = 4$ or $s = 3, q = 3$, every coordinate sign space is distinct. Transitivity makes E the whole ambient module, irreducible and faithful for R . Since R is soluble, the common-derived estimate Lemma 14.1 gives $Z_J(U) \leq Z_J(R) + C_U 2^{\gamma b}$; the comparator degrees $2s < h(3s)$ and tail slopes meet Theorem 7.3. The only remaining case is $s = 3, q = 1$, covered by Proposition 15.8.

The listed cases exhaust all divisors $q \mid s$ and the zero kernels. Each multiplicative bound has strictly positive margin, so Lemma 15.1 applies. There are finitely many actual actions in these widths. Choose one block system and the first applicable certificate for each action, then one any-hot/cold partition on the complete accepted pointing union. The cold coefficients are summed with their original $\Delta(n, n-w)/a(U)$, and the hot sums with their original $((n-w)!a(U)p_n G_{\lfloor n/2 \rfloor})^{-1}$. The finite sum preserves exponential and quadratic decay. There is no additional action weight for isotypes, quotient representations, or alternative block systems. $\square$

16 The bounded affine exceptions

We retain the complete quotient killing translations as a comparator. Only the other normal quotients need character covers. This distinction allows the following arguments to include arbitrary nonsplit extensions.

Lemma 16.1 (Homogeneous sections of a block module). *Suppose G acts linearly on $M = \bigoplus_{i=1}^s V_i$, with $\dim_F V_i = d$ and transitive permutation of the $s \geq 2$ summands. If X^m is a homogeneous semisimple section, then either $\dim_F X \geq 2$ and $m \leq ds/2$, or $\dim_F X = 1$ and $m \leq d \lfloor s/2 \rfloor$. In particular a transitive monomial module on s lines has homogeneous multiplicity at most $\lfloor s/2 \rfloor$.*

Proof. The first assertion is dimension. For the second choose a derangement in the transitive block action: the average fixed-point count is one, whereas the identity fixes $s > 1$ points. A lift g has at most $\lfloor s/2 \rfloor$ block cycles. One basis of a starting block generates its whole cycle over $F[x]$, with x acting as g and the return twist retained. Thus M has at most $d\lfloor s/2 \rfloor$ generators over the PID $F[x]$. Every subquotient has at most that many: take its preimage in a free module of that rank, then use the submodule theorem over a PID. On X^m , the element g is a single scalar, so its polynomial-module generator number is m . This proof does not require semisimplicity of the ambient module. $\square$

Lemma 16.2 (One orbit of constituent types). *Let R act faithfully on $M = \bigoplus_i V_i$ and transitively on the summands, with each V_i irreducible under its stabilizer. Let $D \triangleleft R$ fix every summand, with nontrivial coordinate images D_i . Suppose each D_i is normal in the exact local stabilizer. Then $M \downarrow_D$ is semisimple, has no trivial constituent, and its simple types form one R -orbit. Every nonzero R -section of M is faithful on D .*

Proof. A simple D_i -submodule of V_i and its stabilizer translates span V_i by irreducibility; their sum is semisimple. The isotypic summands form one orbit, since the sum over a proper orbit would be invariant under the stabilizer. The fixed space is invariant under that stabilizer, and is zero by faithfulness and $D_i \neq 1$. Transitivity on blocks joins the local orbits of types into one R -orbit. A nonzero R -section has nonempty invariant type support, so it contains every type in that orbit. The intersection of their kernels is the kernel of D on the faithful M , which is trivial. $\square$

We repeatedly use the following elementary section observation. An elementary abelian p -section of an actual $H \leq S_d$ has rank at most $\lfloor d/p \rfloor$: lift it to a subgroup of H and apply the abelian p -quotient bound there. This does not assign degree d to an arbitrary quotient of H .

Theorem 16.3 (Binary affine local degrees eight and sixteen). *Let U have exact primitive affine local degree $r = 2^d$ on s blocks, where (r, s) is one of*

$$(8, 2), (8, 4), (8, 6), (16, 2), (16, 4).$$

Put $E = U \cap (\mathbb{F}_2^d)^s$ and $R = U/E$. For every literal $N \triangleleft U$ with $E \not\leq N$, the quotient U/N has a faithful tuple of at most $t = s$ irreducible complex characters when $r = 8$, and at most $t = 2s$ when $r = 16$. Consequently all these actions have complete contractive forward rows and negligible scalars.

Proof. Write the exact linear stabilizer as $R_0 \leq \mathrm{GL}_d(2)$, and let $K = R \cap R_0^s$. Its coordinate image K_i is normal in R_0 . If it were trivial, the onto local map from a block stabilizer would factor through $T_i \leq S_{s-1}$. For $d = 3$, the three possibilities $C_7, C_7 : C_3, \mathrm{GL}_3(2)$ proved in Theorem 12.3 all have order divisible by seven, impossible for $s \leq 6$. For $d = 4, s \leq 4$, it would make R_0 a quotient of a subgroup of S_3 . The groups of order dividing six, namely $1, C_2, C_3, C_6, S_3$, have no irreducible binary module of dimension four. For the last assertion, a normal 2-subgroup acts trivially on a simple module. On a nontrivial simple binary S_3 -module, restriction to its normal C_3 supplies scalar multiplication by $\mathbb{F}_4$, and the inverting involution acts semilinearly. Choose a nonzero vector fixed by that involution, which exists in characteristic two. Its $\mathbb{F}_4$ -line is invariant under both generators, so irreducibility forces binary dimension two. Thus $K_i \neq 1$ in all cases.

Every normal 2-subgroup of a faithful irreducible binary linear group is trivial, since its fixed space is nonzero and invariant. Therefore $O_2(K_i) = 1$ and $O_2(K) = 1$. By Lemma 16.2, the physical module is semisimple on K , with no fixed constituent, and each nonzero translation section is faithful on K . Its simple K -constituents have dimension at least two. At degree eight

they have dimension three: a nontrivial normal subgroup of $C_7 : C_3$ contains C_7 , and the other two local possibilities are C_7 and the simple $\mathrm{GL}_3(2)$.

Fix a retained quotient $Q = U/N$, write $\bar{E}$ for its nonzero translation image, and let B be the preimage of K in U . Faithfulness gives $N \cap B = N \cap E$, so $\bar{B}/\bar{E} = K$. A binary minimal normal subgroup outside $\bar{E}$ meets it trivially; its intersection with $\bar{B}$ maps to a normal 2-subgroup of K and is trivial. Such a subgroup therefore centralizes $\bar{B}$. Every nontrivial-core binary type in the whole socle lies in $\bar{E}$, and its multiplicity is at most s for $d = 3$ and $2s$ for $d = 4$.

Keep the whole sum A_2^0 of the core-trivial binary socle types. It meets $\bar{E}$ trivially, because $\bar{E}^{\bar{B}} = 0$; its intersection with $\bar{B}$ again maps to $O_2(K) = 1$. Thus it embeds in a quotient of the actual top T , and has total rank at most $\lfloor s/2 \rfloor$. These are different simple types from those in $\bar{E}$, so the bounds take a maximum.

For an odd prime p , the whole p -socle intersects $\bar{B}$ in a normal p -subgroup. That subgroup centralizes $\bar{E}$, and faithfulness of K on $\bar{E}$ forces its image in K to be trivial. Hence the whole odd socle injects a quotient of T and has rank at most $\lfloor s/p \rfloor$. The whole-socle criterion of Theorem 8.1 therefore supplies the stated common tuple length, including any nonabelian socle in the same slots.

Normals containing E contribute exactly $Z_J(R)$. Fixing tuples on all remaining target quotients before inspecting J gives

$$Z_J(U) \leq Z_J(R) + C_U 5^{tb/3}, \quad C_U = \sum_{N \triangleleft U, E \not\leq N} |\mathrm{Aut}(U/N)|.$$

The action on the nonzero vectors of each local space gives R a concrete faithful degree $(r-1)s < rs$. The exponents satisfy $s \log 5/3 < s$ and $2s \log 5/3 < 2s$, respectively. Theorem 7.3 now applies to this finite set of physical degrees. If $E = 1$, its retained sum is empty and the same comparison is exact. $\square$

Theorem 16.4 (All exceptional affine-nine counts). *Every action with exact primitive affine local degree nine and $s \in \{2, 3, 4, 6, 12\}$ has a complete contractive forward row and negligible scalar, on arbitrary complete sources.*

Proof. Write $M = (\mathbb{F}_3^2)^s$, $E = U \cap M$, $R = U/E \leq R_0 \wr T$, $B_0 = O_2(R_0)$ and $D = R \cap B_0^s$. The complete local matrix classification gives

R_0	B_0	R_0/B_0
$C_4, C_8, D_8, Q_8, QD_{16}$	R_0	1
$\mathrm{SL}_2(3)$	Q_8	C_3
$\mathrm{GL}_2(3)$	Q_8	S_3 .

This is the seven-local input of the finite affine-nine structural certificate: adjoining all elements to the retained sixteen subgroup classes of the literal $\mathrm{GL}_2(3)$ proves subgroup coverage; irreducibility is checked against its four lines. The displayed normal 2-cores and quotients are the indicated elementary matrix groups, not a classification of any imprimitive degree.

Suppose first $D \neq 1$. Its coordinate images are nontrivial normal subgroups of the exact R_0 . The physical module is semisimple on D , and Lemma 16.2 shows every nonzero translation section is faithful on D . In a retained quotient $Q = U/N$, let $\bar{B}$ be the image of the preimage of D . Then $\bar{B}/\bar{E} = D$ and $O_2(\bar{B}) = 1$: a normal 2-subgroup centralizes the normal ternary $\bar{E}$ and its image in D would act trivially.

The entire binary socle therefore injects into a quotient of $R/D \leq S_3 \wr T$. Removing the normal ternary base embeds it further in a quotient of $C_2 \wr T \leq S_{2s}$, so its rank is at most s . Ternary simple

types nontrivial on $\bar{B}$ lie in $\bar{E}$; Lemma 16.1, with local dimension two, bounds their multiplicities by s . The whole $\bar{B}$ -trivial ternary type sum meets $\bar{E}$ and then $\bar{B}$ trivially, and embeds in a quotient of $R/D \leq S_{3s}$, again giving rank at most s . The two sets of types are distinct. For $p \geq 5$, the whole p -socle injects a quotient of T , since both local kernels have order divisible only by two and three. The whole-socle criterion gives a faithful tuple of at most s characters for every retained quotient. Thus

$$Z_J(U) \leq Z_J(R) + C_U 5^{sb/3}.$$

The actual action on nonzero local vectors represents R faithfully in degree $8s$. For each of the five counts, $8s < h(9s)$ and $s \log 5/3 < h(9s)/8$. The finite comparator theorem applies, including the odd physical width 27 with $h(27) = 26$.

If $D = 1$, the linear block kernel $K = R \cap R_0^s$ is trivial. For binary R_0 this is immediate. Otherwise K embeds in S_3^s and every binary subgroup of K , and hence every binary subgroup of its quotients, is elementary. A normal subgroup of $\mathrm{SL}_2(3)$ or $\mathrm{GL}_2(3)$ outside B_0 contains Q_8 and cannot have this property. Every coordinate image of K therefore lies in B_0 , giving $K = D = 1$. So $R = T$. For $s \leq 4$, a point stabilizer of T cannot surject onto any displayed R_0 , whose orders are 4, 8, 16, 24, 48; these counts are impossible in this branch.

For $s = 6, 12$, any quotient has binary and other nonternary socles embedding in a quotient of T . Filter its whole ternary socle by its intersection with $\bar{E}$. The intersection is an actual section of M with type multiplicity at most s , and the outer elementary section of T has rank at most $\lfloor s/3 \rfloor$. The semisimple socle filtration gives a common cover of length $s + \lfloor s/3 \rfloor = 8, 16$. Its exponent is strictly less than $9s/8$, since $5^{64(s+\lfloor s/3 \rfloor)} < 2^{216s-3}$. Corollary 8.4 consumes these entire normal menus. Every argument retained the original source and action weight. $\square$

Theorem 16.5 (Nonzero ternary core at local degree four). *Let U have exact natural A_4 or S_4 local action on s blocks, $2 \leq s \leq 3072$, and put $E = U \cap (\mathbb{F}_2^2)^s$, $R = U/E \leq S_3 \wr T$, $D = R \cap C_3^s$. If $D \neq 1$, the complete action is accepted by Theorem 7.3.*

Proof. Every coordinate image of D is C_3 . Its binary simple types in $M = (\mathbb{F}_2^2)^s$ are the inverse pairs of the nonzero coordinate characters. They form one R -orbit, allowing semilinear inversion. Each nonzero R -section of M is therefore faithful on D and has no fixed vector, as in Lemma 16.2. For a retained quotient, write $\bar{E}, \bar{B}$ for translations and the preimage of D . Then $\bar{B}/\bar{E} = D$ and $O_3(\bar{B}) = 1$.

Consider a homogeneous binary R -section X^m of M and put $e = \dim_{\mathrm{End}_R(X)} X$. Its binary dimension is even. If it is at least four, then $m \leq \lfloor s/2 \rfloor$. In dimension two with endomorphism field $\mathbb{F}_2$, $e = 2$ and $\lceil m/e \rceil \leq \lceil s/2 \rceil$. In dimension two with endomorphism field $\mathbb{F}_4$, the image on X is C_3 . Faithfulness on D makes $D = C_3$ central in R . Changing the field convention on each coordinate by Frobenius aligns its generator with the global scalar ωI . Every R -map then commutes with this scalar and is $\mathbb{F}_4$ -linear; the frame remains monomial. Binary submodules and maps are now genuinely $\mathbb{F}_4$ -linear. Lemma 16.1 over $\mathbb{F}_4$ gives $m \leq \lfloor s/2 \rfloor$. This use explicitly fixes the field and makes no assertion about arbitrary restriction of scalars.

At $s = 3$ improve the remaining endomorphism- $\mathbb{F}_2$ case. The image on X is S_3 and faithful D is its C_3 . Choose a 3-power lift z of a physical three-cycle. It centralizes D . Multiply it by $d \in D$ cancelling its image on X ; the resulting $y = zd$ still has odd order and induces the physical three-cycle. Its fixed space on M has dimension at most two, since a fixed tuple is determined by one coordinate. Odd-order invariants are exact in characteristic two. All of X^m is fixed by y , so $2m \leq 2$. Thus the translation-type Schur capacity is

$$k(s) = 1 \quad (s = 2, 3), \quad k(s) = \lceil s/2 \rceil \quad (s \geq 4).$$

Every binary socle type nontrivial on $\bar{B}$ lies in $\bar{E}$. For the whole trivial-type sum A_2^0 , put $L = NE/E \triangleleft R$. Faithfulness gives $L \cap D = 1$, so $[L, D] = 1$. Its image in R/L is injective; the full preimage $H \leq R$ centralizes D because $[H, D] \leq L \cap D = 1$. It also meets D trivially. The actual frame gives $C_R(D) \cap S_3^s = D$: every coordinate centralizer of the projected C_3 is C_3 . Hence H embeds in the actual top T , and $A_2^0 = H/L$ has rank at most $\lfloor s/2 \rfloor$.

For odd p , the whole p -socle intersects $\bar{B}$ trivially: for $p = 3$ use $O_3(\bar{B}) = 1$, and otherwise use orders. It injects a quotient of $R/D \leq C_2 \wr T$, then a quotient of T , so has rank at most $\lfloor s/p \rfloor$. The whole-socle criterion supplies $k(s)$ characters on every retained quotient, including the nonabelian socle in the same slots.

Thus $Z_J(U) \leq Z_J(R) + C_U 5^{k(s)b/3}$, with all killed translation normals exactly in $Z_J(R)$. The comparator has its concrete faithful degree $3s < 4s$. The retained exponent is less than $4s/8$: for even s use $5^{32} < 2^{75}$, and for odd $s \geq 5$ use $75(s+1) \leq 96s - 3$; counts two and three are immediate. This is a fixed finite band of actual actions, so Theorem 7.3 applies once to their union. $\square$

17 The exceptional small pair actions

Put $\beta = \log_2(38/25)$ and $\gamma_5 = \frac{1}{3} \log_2 5$. Throughout this section, a pair action means an actual transitive group $U \leq C_2 \wr S_s$ in its specified action on $2s$ points. Write $E = U \cap C_2^s$, $T = U/E$, and $M_s = \mathbb{F}_2^s$ with its actual T action. Every normal subgroup mentioned below is a normal subgroup of this original U , unless another group is specified.

Lemma 17.1 (Relative socles and fixed prefixes). *Let $F \triangleleft U$ be a binary group and $S = U/F$. Suppose that, for every $N \triangleleft U$, the multiplicity m_X of each simple binary type X in $\text{Soc}(U/N) \cap FN/N$ satisfies $m_X \leq qe_X$, where $e_X = \dim_{\text{End}_U X} X$. Suppose every quotient of S has a faithful tuple of at most k irreducible characters. Then every $Q = U/N$ has a faithful tuple consisting of a k -character prefix with joint binary kernel, followed by q general characters. Consequently*

$$|\text{Epi}(J, Q)| \leq |\text{Aut } Q| [Q : O_2(Q)]^q 5^{kb/3} (38/25)^{qb} \quad (J \leq S_b).$$

If Q is binary, the whole tuple can instead be counted in the known binary-source mode. If a specified top quotient has a faithful tuple of r binary-primary linear characters, one may use those r positions and the same q relative general positions. For $F = E$, the relative hypothesis follows whenever every homogeneous section X^m of M_s satisfies $m \leq qe_X$.

Proof. Put $A = FN/N$. A minimal normal subgroup of Q contained in A is central in A and elementary abelian. Thus $V = \text{Soc}(Q) \cap A$ is a semisimple binary Q/A -module. The dual V^* has q module generators: for X^m the required number is $\lceil m/e_X \rceil$, and distinct simple types share positions. Induce each corresponding linear character of V to Q and choose an irreducible constituent. Its restriction to V consists of conjugates of that linear character; hence the intersection of these q kernels with V is trivial. Inflate a faithful tuple of Q/A . The joint kernel of the resulting tuple is normal in Q , lies in A , and meets $\text{Soc}(Q)$ trivially, so it is trivial. This does not extend a character to a nonabelian A or split the original extension. Apply Proposition 8.3 to the fixed prefix. When $F = E$, V is a section of the original M_s . All choices depend on Q , not on the later complete source J . $\square$

Lemma 17.2 (Two elementary module bounds). *A transitive monomial module of dimension $r > 1$ over a field has no homogeneous section X^m with $m > \lfloor r/2 \rfloor$. For the regular binary*

V_4 module $W = \mathbb{F}_2[x, y]/(x^2, y^2)$, every submodule has head dimension at most two; the unique submodule attaining two is the augmentation ideal $I = (x, y)$, of dimension three. These statements also hold after a field extension.

Proof. A transitive permutation group has a derangement: the average number of fixed points is one, and the identity fixes more than one point. Choose a monomial element whose permutation is a derangement. Its module over the principal ideal domain $k[t]$ is a sum of at most $\lfloor r/2 \rfloor$ cyclic modules, one per cycle. A submodule of a c -generated torsion module over a principal ideal domain is c -generated: lift it to a submodule of a free module of rank c . The same holds for a quotient. On the other hand X^m , restricted to this element, requires at least m generators, by any nonzero primary component and the invariant-factor theorem. For the second assertion a proper ideal lies in I , whose radical is $I^2 = \langle xy \rangle$. A one-dimensional image in I/I^2 generates a cyclic ideal, whereas a two-dimensional image forces the entire I . The whole ring has cyclic head. This proves the claim and its equality assertion over any field of characteristic two. $\square$

17.1 The finite inputs and what they certify

We use the following finite input on *top actions*, separately from the universal lifting arguments that follow. For a simple binary T -module X , write $\delta_X = \dim_{\mathbb{F}_2} X$. Define

$$\tau(T) = \max_{B \leq M_s, X} \left\lceil \frac{\dim_{\mathbb{F}_2} \text{Hom}_T(B, X)}{\delta_X} \right\rceil, \quad \kappa(T) = \max_{L \triangleleft T} c(T/L),$$

where c denotes the least size of a faithful irreducible character tuple. The formula for τ is exactly the largest Schur demand of a homogeneous section: the multiplicity of X in the head of B is $\dim \text{Hom}_T(B, X) / \dim \text{End}_T(X)$.

Lemma 17.3 (Certified top data). *The top-action certificates have the following properties.*

(i) *Among the 301 transitive degree-twelve actions, 283 have (τ, κ) equal to*

$$(1, 1), (1, 2), (1, 3), (2, 1), (2, 2), (3, 1),$$

with respective multiplicities 71, 102, 19, 33, 56, 2. The remaining eighteen consist of eight binary-core actions with quotient C_3 or S_3 as in Lemma 17.4; eight actions with an elementary normal ternary group on four triples as in Lemma 17.5; and two paired actions with six-point tops as in Lemma 17.6.

(ii) *For the two six-point groups*

$$S_{36} = \langle (124), (24)(56), (13)(25)(46) \rangle, \quad S_{72} = \langle (124), (24), (13)(25)(46) \rangle,$$

all homogeneous sections of M_6 have Schur demand at most one. Their only simple types have dimensions one and four, with commuting field $\mathbb{F}_2$. Every nontrivial quotient has a faithful irreducible character except C_2^2 , which has two faithful binary linear positions.

(iii) *For all 24 nonbinary transitive degree-eight groups S , every nontrivial homogeneous type in M_8 has Schur demand at most one, every trivial section has dimension at most two, and $\kappa(S) \leq 2$. If an invariant submodule has trivial head dimension two, it is a unique five-dimensional submodule B . Every invariant $D \geq B$ satisfies $\dim D / (B + [D, S]) \leq 1$.*

(iv) *The 29 zero-ternary minimal-four degree-sixteen tops have $\tau \leq 4$, $\kappa \leq 2$. The 15 degree-sixteen tops with two primitive affine eight-point blocks have $\kappa \leq 1$ and binary permutation composition Schur demand at most four. The 20 primitive affine degree-sixteen tops have the respective bounds two and four. The two inequivalent nontrivial binary composition factors of the projective eight-point $\mathrm{PSL}_2(7)$ module have dimension three, commuting field $\mathbb{F}_2$, and each occurs once; the trivial factor occurs twice.*

Proof. These are the finite calculations recorded by the pair-top certificate contract in Appendix A. We specify their mathematical coverage, since a list of tested module examples would not suffice. Starting with every vector, form its cyclic submodule and close the resulting list under sums. Every invariant subspace is the sum of cyclic submodules of its vectors, so this constructs the complete invariant-subspace lattice. Maximal intervals give the simple modules; intertwining linear equations on the same generators identify their isomorphism types, their endomorphism fields, and the displayed Hom spaces. A composition chain gives the separate composition bounds. Thus every numerator and every homogeneous quotient is covered. For normal subgroups start with kernels of all irreducible characters and close under intersections. Every normal subgroup occurs, since the regular representation of its quotient is faithful and decomposes into irreducibles. Literal kernels, not quotient isomorphism types, are retained. Searching these kernel intersections proves the asserted covers for all quotients.

Completeness of the degree-twelve and primitive action lists is the published classification input [Hul05, HR20, RD05, Hul23, HRDR23]. The imprimitive degree-eight and two-affine-block constructions use all normal quotients of the local group, all graph isomorphisms between these quotients, and all block-swap cosets in their actual normalizers; Goursat's lemma and the order-two coset condition prove completeness. The four-pair constructions use the full invariant-kernel list and all lifts of fixed top generators. The zero-ternary list similarly uses all degree-four local subgroups, both induced binary linear frames, all invariant kernels, and every solution of the generator-relator cocycle equations. The computed cocycle-space dimension, modulo the coboundary-space dimension, verifies that every complement class occurs. Transporters identify actions in the stated lists. These procedures certify only the finite top data. In particular they make no assertion about unlisted normal subgroups of a subsequent pair lift. $\square$

17.2 The twelve-pair actions

Lemma 17.4 (The binary-core twelve-point tops). *Let T be one of the eight binary-core actions in Lemma 17.3(i). Put $P = O_2(T)$. For every pair lift U and every $N \triangleleft U$, put $Q = U/N$. If $T/P = C_3$, Q admits one binary-primary linear and four general faithful positions. If Q is nonbinary, the linear position can be made an ordinary linear anchor with binary kernel. If $T/P = S_3$, every nonbinary Q admits four general faithful positions, one of which has binary kernel. Every binary Q admits one binary-primary linear and four general faithful positions.*

Proof. First suppose $T/P = C_3$ and choose h of order three. There are three P -orbits of size four, and h cycles them. Identify them by h . Then $P \leq L^3$ has full local projections, with $L = C_4, V_4$, or D_8 . The simple binary types are 1 and the natural C_3 type X of binary dimension two and Schur dimension one. If $L = C_4$ or D_8 , restrict each coordinate module to an actual regular C_4 . Its module is $\mathbb{F}_2[t]/(t^4)$, with cyclic submodules. A three-coordinate filtration bounds every P -head by three; hence the homogeneous-section bounds are three for 1 and one for X . If $L = V_4$, Lemma 17.2 bounds a P -head by six. Equality forces dimension three in each coordinate filtration and rules out a full coordinate projection, since that would contribute only one generator. Thus

the numerator is the sum of the three augmentation ideals. Its head, with h cycling coordinates, is two regular C_3 modules, namely $1^2 \oplus X^2$. It cannot contain X^3 . Thus the X bound is two.

In this last case a trivial T -section of rank four is also impossible. Odd-order averaging $e = 1 + h + h^2$ would give, for its numerator B , $B^h = M_{12}^h$ and $([B, P])^h = 0$. Hence $e(p-1)e = 0$ for every $p = (p_1, p_2, p_3) \in P$, or on the regular four-point coordinate

$$R_{p_1} + R_{p_2} + R_{p_3} = I.$$

The four regular permutation matrices are linearly independent. Therefore $p = 0$ or p is a rotation of $(x, x, 0)$ for a nonzero $x \in V_4$. Rotation invariance and addition show that two independent values of x would produce a tuple with three nonzero entries, a contradiction. The coordinate projections could not then be all V_4 . The trivial section bound is three.

Coprime centralizer lifting maps $C_P(h)$ onto the fixed subgroup in every quotient of P by a T -normal subgroup. Evaluation on one coordinate embeds $C_P(h)$ in L . Consequently every quotient of T has central binary rank at most two. Moreover $T' = [P, h]P'$ and the fixed part of T'/P' is trivial, so $C_{T'}(h) = C_{P'}(h)$; evaluation embeds this in L' , of order at most two. Every quotient of T therefore has derived central binary rank at most one. Its nontrivial binary-type multiplicity is at most two for $L = V_4$, and at most three otherwise: filter any elementary section of $P \leq L^3$ through the coordinates, each of rank at most two.

For the whole Q , filter its binary socle by its intersection with EN/N and its image in $Q/(EN/N)$. Both are modules for the same original T ; same-type multiplicities add. The resulting central rank is at most $3 + 2 = 5$, the derived central rank at most $3 + 1 = 4$, and the nontrivial multiplicity at most $2 + 2 = 4$ or $1 + 3 = 4$. The odd socle is central of rank at most one and has zero intersection with Q' . There is no nonabelian socle. Theorem 8.1 gives the simultaneous profile $(r, l, t) = (1, 0, 4)$. If Q is nonbinary, multiply the designated binary-primary linear character by a faithful cubic character of $Q/O_2(Q)$. Its kernel shrinks to its intersection with $O_2(Q)$, since a binary-power root and a nontrivial cube root cannot have product one. This gives the claimed ordinary linear anchor.

Now suppose $T/P = S_3$. The certified actions have either six P -orbits of size two, with regular S_3 action on them, or three of size four, with natural S_3 action. For h of order three, $T_0 = P\langle h \rangle$ has respectively two six-point orbits or one twelve-point orbit. The simple binary types are 1 and the natural S_3 type Y , whose Schur dimension is two; restriction to $\langle h \rangle$ preserves its multiplicity. A nonbinary Q has binary kernel N , since the odd part of $|U|$ is three. Its top quotient is T/L for $L \leq P$. Any elementary section of the actual $P \leq S_{12}$ has rank at most six by [RDT25, Theorem 2.8]; its Y multiplicity is therefore at most three.

In the size-two case each local module is a cyclic regular C_2 module. Thus the pair-module Y multiplicity is at most three. Frattini's argument gives $T = PN_T(\langle h \rangle)$; take a binary element in this normalizer mapping to a reflection. It swaps the two triples of P -orbits and therefore acts without fixed points on the four h -orbits. It has at most two cycles on these four labels. Exact h -invariants and the cyclic module argument bound every trivial T -section by two. In the size-four case the preceding C_3 proof bounds pair-module trivial multiplicity by three and natural multiplicity by two.

In both cases $C = C_P(h)$ acts faithfully on the four h -orbits: the kernel in the ambient centralizer is odd. Thus $C \leq S_4$ is binary, and every current top quotient has central binary rank at most two by coprime fixed lifting. If that rank is two, C is noncyclic and contains a V_4 . A faithful V_4 action on four points is regular or has two two-point orbits with distinct kernels; in either case its trivial module-section bound is two. Exact h -invariants therefore improve the pair bound to two in this same case. The whole central rank is at most four, and the whole natural

Schur demand is at most $\lceil (3+3)/2 \rceil = 3$. The odd socle rank is at most one. Theorem 8.1 gives four general positions. One has binary kernel: otherwise all four normal kernels contain a Sylow three subgroup, hence all its conjugates, contradicting their trivial intersection.

It remains to handle binary Q in the S_3 case, for which the last argument about the top quotient was not asserted. An actual Sylow three element of U is semiregular on the original twenty-four points, with eight three-cycles, and is killed by N . Frattini gives $U = NN_U(\langle h \rangle)$. In the ambient normalizer $C_3^8 : (C_2 \times S_8)$, the common inversion factor C_2 must be retained. The odd base of the actual normalizer is killed by N ; its image $H \leq C_2 \times S_8$ is binary and surjects onto Q . It has a faithful ten-point action, so all elementary sections have rank at most five by the same published abelianization bound. For a Sylow subgroup $P_8 = C_2^4 : D_8$ of S_8 , one has $|P_8'| = 2^4$: the base commutator has order eight and $|D_8'| = 2$. Hence $|Q'| \leq 16$. The central rank of Q is at most five and its derived central rank at most four. The simultaneous feasibility criterion again gives one binary-primary linear and four general positions, now in the known binary-source mode. $\square$

Lemma 17.5 (Four ternary orbits). *Suppose T has an elementary normal ternary subgroup C of rank at least two with four orbits of size three, T/C binary, and $|O_2(T)| \leq 2$. Every pair lift U has a quotient R with a specified faithful twenty-point action and a constant D_U such that*

$$Z_J(U) \leq Z_J(R) + D_U 5^b \quad (J \leq S_b).$$

Proof. Averaging over C gives the T -equivariant decomposition $M_{12} = M_0 \oplus W$, with $\dim M_0 = 4$, $\dim W = 8$. The map $(v, t) \mapsto (\text{avg}_C(v), t)$ on the ambient semidirect product has kernel W . Restricted to U , its kernel is $E_1 = E \cap W = [E, C]$; put $R = U/E_1$. The ambient $M_0 : T$ acts faithfully on the disjoint union of the twelve pair labels and the eight C -orbit labels on the original twenty-four points. The first action detects T and the second detects M_0 , proving faithfulness of this twenty-point representation after restriction to R .

Fix an original N not containing E_1 , and choose a minimal normal S in the nonzero image of E_1 in $Q = U/N$. On restriction to C , W consists of four nontrivial two-dimensional local types. Their kernels have trivial intersection, and they form one T -orbit. There are at least two distinct types, since C has rank at least two; each has multiplicity at most two. Every nonzero T -section of W therefore acts faithfully on C . In particular if L is the kernel of the action of T on S , then $L \cap C = 1$, so L is normal binary and $L \leq O_2(T)$.

Put $A = EN/N$. Since A is abelian, there is an exact sequence

$$1 \longrightarrow A \longrightarrow C_Q(S) \longrightarrow L/\pi(N) \longrightarrow 1.$$

Every minimal normal subgroup of Q commutes with S , including S itself. Thus the entire socle lies in this binary centralizer. Its image modulo A has dimension at most one and is trivial. The trivial part in A is a section of M_0 , the ordinary permutation module on four blocks, so has rank at most two by Lemma 17.2. Every nontrivial type in A lies in W and has multiplicity at most two by restriction to C . The whole socle therefore has Schur demand at most three, and Theorem 8.1 supplies three general positions. Its epimorphisms cost at most $|\text{Aut } Q|5^b$. Finally the interval $N \geq E_1$ is exactly the normal-subgroup lattice of R , with the same quotients and epimorphism sets. Summing that interval gives exactly $Z_J(R)$; summing the other normals gives the asserted constant $D_U = \sum_{N \not\geq E_1} |\text{Aut}(U/N)|$. $\square$

Lemma 17.6 (The paired six-point tops). *Suppose $U \rightarrow T \rightarrow S$ has degrees 24, 12, 6, with both successive systems consisting of pairs, and $S = S_{36}$ or S_{72} from Lemma 17.3. Every original*

quotient admits either one general anchor and three relative general positions, or two binary-primary linears and three relative general positions in the known binary-source mode. A binary quotient with no exceptional C_2^2 top needs at most four general positions.

Proof. Let F be the kernel of $U \rightarrow S$. On the six original four-point blocks, $F \leq D_8^6$, and F/E is a submodule of M_6 . Let $C_0 \leq M_{12}$ be the vectors constant on the six pairs of pair labels. Both C_0 and M_{12}/C_0 are M_6 . Fix N , put $A = FN/N$, and take a homogeneous component X^m of $\text{Soc}(U/N) \cap A$. Its preimage $D \leq F$ has the three successive factors obtained from

$$D \cap E \cap C_0, \quad (D \cap E + C_0)/C_0, \quad \pi(D) \leq F/E.$$

The denominators are the respective images of $D \cap N$; the last is $\pi(F \cap N)$, since D contains $F \cap N$. Thus all three factors are homogeneous sections of the same M_6 for the same original S type. Their multiplicities add, and Lemma 17.3(ii) gives $m \leq 3e_X$. Apply Lemma 17.1. Except for top C_2^2 , a nontrivial top quotient has a one-character prefix with binary joint kernel. For top C_2^2 , Q itself is binary and the prefix consists of two full-order binary linear characters. A trivial top needs only the three relative positions. No part of this proof assumes that F or A is abelian. $\square$

Theorem 17.7 (Complete twelve-pair estimate). *The family of all actual pair actions with twelve pairs and nonbinary top has a complete exponentially small forward row and a quadratic-exponentially small scalar, with its original action weights.*

Proof. For the 283 generic tops, Lemma 17.1 gives slope $\kappa\gamma_5 + \tau\beta$ and target-dependent finite constants. For each of the six pairs in Lemma 17.3(i),

$$5^{64\kappa} 38^{192\tau} < 2^{573} 25^{192\tau},$$

so the slope is less than $3 - 1/64$. The binary-core slopes are $\frac{1}{3} \log_2 3 + 4\beta$, $\frac{1}{2} + 4\beta$, and $\gamma_5 + 3\beta$, all less than $3 - 1/64$. The paired-six slopes are $\gamma_5 + 3\beta$, $1 + 3\beta$, and 4β , with the same strict margin. These inequalities follow, without rounding logarithms, by raising to the 192nd power; the corresponding numerators are $3^{64} 38^{768}$, $2^{96} 38^{768}$, $5^{64} 38^{576}$, $2^{192} 38^{576}$, and 38^{768} , against 2^{573} times the matching power of 25. Proposition 8.3 and Corollary 8.4 sum every literal normal in these finite original width-twenty-four actions. Constants such as $[Q : O_2(Q)]^q |\text{Aut } Q|$ are fixed before choosing J . For the other eight tops, Lemma 17.5 and Theorem 7.3 apply with $v = 20 < 24$ and $\eta = \log_2 5 < 3$. This last assertion includes the entire normal interval of the quotient, rather than only its faithful target. There are finitely many subgroups of S_{24} , so the complete union is covered by one fusion, retaining the factor $1/(b!a(U)p_n G_{\lfloor n/2 \rfloor})$ in each original hot sum. $\square$

17.3 The sixteen-pair actions

Lemma 17.8 (Paired eight-point tops). *Suppose $U \rightarrow T \rightarrow S$ has degrees 32, 16, 8, with both successive systems consisting of pairs and S nonbinary. For every original normal subgroup, the relative socle in the direct quartet kernel has Schur demand at most four. Every quotient of U therefore admits a two-character general prefix with binary joint kernel and four relative general positions.*

Proof. Write $F = \ker(U \rightarrow S) \leq D_8^8$. The three layers used in Lemma 17.6 are now sections of M_8 . A nontrivial simple type has demand at most three by Lemma 17.3(iii). We must improve the possible sum of three trivial heads from six to four using the original commutator denominator.

For an invariant binary module B put $d_S(B) = \dim B/[B, S]$. For invariant $K, W, L \leq M_8$ such that their coordinatewise product span KW lies in L , we claim

$$d_S(K) + d_S(W) + \dim L/([L, S] + KW) \leq 4. \quad (17.1)$$

A Sylow binary subgroup of the actual transitive $S \leq S_8$ is transitive. Hence every nonzero invariant submodule contains the constant vector $\mathbf{1}$: a binary group has a nonzero fixed vector on every nonzero binary module, and M_8^P consists of constants. Dually, every proper invariant submodule lies in the augmentation hyperplane I . If K or W is zero, the two remaining heads are at most two each. If both heads are at most one, the result is again immediate. If $d_S(K) = 2$, $d_S(W) = 1$, then $KW \supseteq K$ and the last term is at most one by the five-dimensional equality assertion in Lemma 17.3. If both heads are two, $K = W$ is that five-dimensional submodule. It cannot be self-orthogonal in $\mathbb{F}_2^8$, so some product has odd coordinate sum. Thus KW is an invariant submodule not in I , whence $KW = M_8$ and the last term vanishes. This proves (17.1).

Fix N and put $A = FN/N$. Lift the trivial part of $\text{Soc}(U/N) \cap A$ to $D \leq F$. Set $B = D \cap E$ and let $C \leq M_{16}$ be constant on the eight pairs of pair labels. The three binary coordinate modules are

$$K = \pi(D) \leq F/E \leq M_8, \quad W = (B + C)/C \leq M_8, \quad L = B \cap C \leq M_8.$$

Inside a single quartet the commutator of an internal pair flip and a pair interchange is their coordinate product. Therefore the original commutator relation is $[B, D] = KW \leq L \cap N$. Also $[L, S] \leq L \cap N$, since the lifted section is central in U/N . The exact three-step socle filtration consequently has total rank at most the left side of (17.1), hence at most four. This uses the literal intersection with N in the bottom layer; replacing it by an unrelated denominator would lose the improvement. The top cover is two by Lemma 17.3(iii), so Lemma 17.1 completes the proof, even when A is nonabelian. $\square$

Lemma 17.9 (Quotients of an actual degree-eight group). *Let $G \leq S_8$ be transitive. Every quotient which is nonbinary or nonabelian has faithful character cover at most two. An abelian binary quotient has at most four faithful binary linear positions.*

Proof. We use the following precise consequence of the excessive binary normal-pair classification [RDT25, Definition 5.1 and Proposition 5.3(i)]: if P is transitive binary of degree eight and $S \triangleleft P$, then

$$d_P(S) > 2 \implies [P : S] \leq 2, \quad d(P) \leq 4.$$

Here $d_P(S) = \dim S/(S^2[S, P])$ by [RDT25, Lemma 2.5(i)]. Every Sylow binary subgroup of an actual transitive group of binary degree is transitive, by the orbit-stabilizer inequality.

We first show that $\dim A/(A^2[A, G]) \leq 2$ whenever G/A is nonbinary. Otherwise, with $S = A \cap P$, the displayed result gives $[G : A]_2 \leq 2$. The Frattini normalizer $L = N_G(S)$ satisfies $G = AL$ and is transitive on the original eight points: S is transitive on each original A -orbit, and L is transitive on their labels. The group L/S is nonbinary with binary part of order at most two. It has a nontrivial normal odd subgroup. Indeed this is immediate for odd order; for order twice an odd number use the kernel of the sign of its regular permutation action. This subgroup survives in $L/O_2(L)$.

Here is the needed compression fact, with its original action specified. If a faithful transitive L of degree n has a nontrivial normal odd subgroup modulo $P_0 = O_2(L)$, take an odd complement K in its full preimage. Every L -trivial elementary section of P_0 is an image of a subgroup of $C_{P_0}(K)$, by coprime fixed lifting. This centralizer acts faithfully on the K -orbits: the kernel on each odd orbit centralizes a transitive odd group, hence is semiregular of odd order. If K moves

the P_0 -orbit labels, its orbits cover at least three such labels, giving at most $n/3$ orbits. Otherwise its local action on every P_0 -orbit is nontrivial, by transitivity and conjugacy of complements. On a binary orbit of size d , a prime-order odd element has at most $d/4$ fixed points. To verify this, its fixed points form one orbit of its binary centralizer, by complement conjugacy in the binary point stabilizer; thus their number is $2^c < d = 2^a$. If its prime order is q , then $q \mid 2^a - 2^c$, so $a - c \geq \text{ord}_q(2) \geq 2$. The element has at most $(d + (q - 1)d/4)/q \leq d/2$ cycles. In either case K has at most $n/2$ orbits. Applying the actual abelianization bound [RDT25, Theorem 2.8] to the faithful centralizer action bounds every such binary section by $n/4$. Apply this with $n = 8$ to the image of S in $A/(A^2[A, G])$, a trivial section of $O_2(L)$, to obtain the contradiction.

For a nonbinary $Q = G/N$, take the preimage of $\Omega_1(Z(Q))_2$. The marked bound just proved gives central binary rank at most two. Every elementary binary socle section has rank at most four by the actual Sylow-preimage form of Theorem 2.8; a nontrivial simple binary type has dimension at least two and therefore multiplicity at most two. All odd primary ranks are at most $\lfloor 8/p \rfloor \leq 2$, again using actual preimages, so Theorem 8.1 gives cover two. For a nonabelian binary Q , the actual P surjects onto Q . The preimage of $\Omega_1(Z(Q))$ has index at least four, since a group cyclic modulo its center is abelian. The excessive-index bound gives central rank at most two. Finally an abelian binary quotient needs $d(Q) \leq d(P) \leq 4$ full-order linear characters. $\square$

Lemma 17.10 (Four ternary blocks and their top quotients). *Let $R \leq S_3 \wr B$ on twelve points, where B is transitive of degree four, and $H = R \cap C_3^4 \neq 1$. Put $L = R \cap S_3^4$ and $A = L/H$. If $A = 1$, every quotient of R has cover at most two. In general every quotient retaining a nontrivial image of H has cover at most two. In the exact-local- S_3 case, R/H has a specified faithful transitive degree-eight sign action.*

Proof. The coordinates of H are all nonzero, so R is transitive on the original twelve points. Suppose $A \neq 1$. Its nontrivial coordinate inversion characters form one R -orbit and have trivial common kernel. Maschke's theorem on this binary A shows that every nonzero R -section of H is faithful on A . Thus for a quotient retaining $\bar{H} \neq 1$, $N \cap L \leq H$ and $C_{\bar{L}}(\bar{H}) = \bar{H}$. The whole binary socle, and every $p > 3$ socle, consequently injects into a quotient of the actual four-point B . Their ranks are at most two and $\lfloor 4/p \rfloor$. The total ternary socle rank is at most four from the original twelve-point Sylow preimage. A nonscalar ternary type therefore has Schur demand at most two. For a scalar type χ , if $\chi|_A \neq 1$ its top image is zero, since a normal ternary subgroup of the quotient by $\bar{H}$ centralizes the normal binary A . Its entire contribution is a homogeneous section of the transitive monomial $\mathbb{F}_3^4$, of dimension at most two by Lemma 17.2. If $\chi|_A = 1$, its H piece is zero by the inversion-character support, and it injects into a quotient of B , giving rank at most one.

If $A = 1$, all nonternary socles inject past $\bar{H}$ into a quotient of B , while nonscalar ternary types still have demand at most two. For a scalar ternary socle X^m , let D be its preimage in R . For the trivial scalar, $D^3[D, R] \leq N$ and the degree-twelve relative bound of Theorem 9.5 gives $m \leq 2$. For a nontrivial scalar put $K = \ker \chi$. Then $D, N, H \leq K$. The actual K action is transitive of degree twelve or has two six-point orbits. Filtering the same relative head by those orbit kernels gives respectively the bounds 2 or $1 + 1 = 2$ from that theorem. This includes every original normal subgroup, whether or not it retains H . The simultaneous socle criterion proves the claimed cover bounds. Finally the sign map on each local S_3 has kernel exactly H on R . The induced group in $C_2 \wr B$ is transitive on its eight points because each block stabilizer contains local inversion. This is the asserted faithful action. $\square$

Lemma 17.11 (Nonzero ternary four-point frames). *Suppose $T \leq L \wr B$ has four blocks of size four with exact primitive local $L = A_4$ or S_4 . Put*

$$E_0 = T \cap V_4^4, \quad R = T/E_0, \quad H = R \cap C_3^4 \neq 1.$$

Let M_0 be the full preimage of E_0 in any original pair lift U . For every original N , the relative socle in M_0N/N has Schur demand at most four. In the S_4 case, its H -nontrivial types have demand at most two. If moreover $E_0 = V_4^4$, the whole relative demand is at most two.

Proof. The preimage T_0 of H in T has local image A_4 : that image is normal in L and maps onto C_3 . Write a local binary permutation module as $W = \mathbb{F}_2[V_4]$, with flag $0 < C = I^2 < I < W$. The natural translation module V_4 is canonically I/C , via $v \mapsto v - 1$ modulo I^2 . Consequently, for each H -nontrivial simple type in the relative socle, its pair piece and its E_0 piece are sections of the *same* natural module $V = \mathbb{F}_4^4$, with the same R action and endomorphism field. This remains true for correlated E_0 and nonsplit extensions. Every nonzero R -section of V is faithful on H : on restriction to H , the local nontrivial types form one orbit with trivial intersection of their kernels.

In the A_4 case, a type of binary dimension at least four has multiplicity at most two in each copy of V ; a type of dimension two with Schur dimension at least two has total demand at most $\lceil 8/2 \rceil = 4$. In the remaining case, dimension two and Schur dimension one, faithfulness on H implies $H = C_3$ and $[R, H] = 1$. Orient the coordinates so that H is the same scalar of $\mathbb{F}_4^*$ on all four coordinates. The actual R action is now transitive monomial over $\mathbb{F}_4$. Lemma 17.2 bounds the multiplicity in each copy by two, again giving total Schur demand four. This orientation is applied to the same frame; it does not merge distinct binary types.

For $L = S_4$ no natural type has Schur dimension one: such an image would be scalar and abelian, whereas faithfulness on H would then kill $[R, H]$, contrary to local inversion. A type of dimension at least four consequently has total demand at most two. For dimension two, faithfulness again gives $H = C_3$. After orienting H globally, $R^+ = C_R(H)$ is an index-two transitive monomial group over $\mathbb{F}_4$. It is transitive because every coordinate stabilizer contains an inverting element outside R^+ . Restricting the homogeneous section to R^+ and applying Lemma 17.2 bounds its multiplicity in each copy by two; division by Schur dimension two gives total demand at most two.

An H -trivial type has no E_0 piece and lies entirely in the pair layer. It is trivial under T_0 . The local A_4 module has trivial homogeneous sections of dimension at most one: its invariant submodules are $0, C, I, W$, and only C and W/I are trivial. Filter by the four coordinate kernels, with denominator the intersection of the original denominator with each kernel. Every factor is such a local section, so the whole dimension is at most four. This proves the general relative bound.

If $E_0 = V_4^4$, let D/F be an H -trivial homogeneous section in the pair layer. The coordinate projections of D are uniformly $0, C, I$, or W , by transitivity and the same local flag. If they are W , independent translations imply $I^4 \leq [D, E_0] \leq F$; the section is then a section of the ordinary four-coordinate module. If they are I , independent translations give $C^4 \leq F$, and the remaining natural layer has no H -trivial constituent, so the section vanishes. If they are C , it is again a section of the ordinary four-coordinate module. Its homogeneous multiplicity is at most two by Lemma 17.2. Together with the natural-type bound this proves the final assertion. $\square$

Lemma 17.12 (The rank-four sign boundary). *In the exact-local- S_4 situation of Lemma 17.11, if the actual sign group $K = R/H \leq C_2 \wr B$ has an abelian binary quotient needing four generators, then $E_0 = V_4^4$.*

Proof. Let $A = K \cap \mathbb{F}_2^4$. The elementary abelianization sequence gives $d_2(K) \leq \dim A/[A, B] + d_2(B)$. The five transitive groups of degree four are C_4, V_4, D_8, A_4, S_4 . For C_4, D_8, S_4 , an actual

regular four-cycle makes every invariant base submodule have head at most one; their binary abelianizations have ranks 1, 2, 1. For A_4 the corresponding bounds are one and zero. Thus only $B = V_4$ can reach four. Lemma 17.2 then forces $A = I$, the regular augmentation ideal, and B has rank two. Modulo I , the actual K is the graph of a homomorphism $\phi : B \rightarrow C_2$. For a lift (v, b) the square has image

$$(1 + b)v = \phi(b)(b - 1) \pmod{I^2}.$$

If $\phi \neq 0$ this kills a nonzero additional direction of the base head in the abelianization, giving at most three generators. Hence $\phi = 0$ and $K = I : B$.

The four coordinate inversion characters of I are nontrivial and pairwise distinct. Over $\mathbb{F}_3$ their weight spaces are the four coordinate lines. The nonzero invariant $H \leq \mathbb{F}_3^4$, with full coordinate projections, must therefore be C_3^4 . Its four natural binary coordinate types on V_4^4 are distinct. Each projection of E_0 is the full V_4 , since the local derived subgroup of T_0 is V_4 . Semisimplicity under this H now forces $E_0 = V_4^4$. $\square$

Lemma 17.13 (The two primitive eight-point blocks). *If the sixteen-point top T has two minimal eight-point blocks, every original pair lift has a general prefix of at most two and relative tail of at most four.*

Proof. The primitive degree-eight list is

$$2^3 : C_7, \quad 2^3 : (C_7 : C_3), \quad \text{AGL}_3(2), \quad \text{PSL}_2(7), \quad \text{PGL}_2(7), \quad A_8, \quad S_8.$$

This is the actual action list supplied by the primitive classification cited in Lemma 17.3. The first three local cases are covered by the complete fifteen-action graph-and-swap construction there, whose composition bounds imply the required homogeneous section bounds and whose top quotient cover is at most one.

For the other four put $S = \text{Soc}(L)$ for the local group. Then S is simple, $L' = S$, and L/S has order at most two. The block kernel projects onto L on both blocks. Its derived group projects onto S on both, so $P = T \cap S^2$ is, by Goursat's lemma, either S^2 or the graph of an automorphism of S . It is perfect, centerless, and normal in T , with T/P embedded in D_8 . In any quotient of T , the image of P is a product of nonabelian simple groups. The abelian socle meets it trivially and injects into a quotient of a subgroup of D_8 , whose elementary ranks are at most two. The simultaneous criterion gives top cover at most two.

For $S = A_8$, the natural binary module has two trivial composition factors and one simple factor of dimension six. For $S = \text{PSL}_2(7)$, Lemma 17.3(iv) gives two trivial and two inequivalent simple three-dimensional factors, each once. Thus the two-block module restricted to P has at most four trivial factors and at most two copies of any nontrivial simple type, whether P is a product or a twisted graph. A simple T -module restricts semisimply to P : its P -socle is nonzero and T -invariant, and hence is the whole module. Its constituents form one orbit. A type with trivial P action factors through the binary T/P and is therefore trivial; every other type has multiplicity at most two by selecting any one of its nontrivial P -constituents. The original module composition multiplicities are therefore at most four, hence so are the Schur demands of all its homogeneous sections. Apply Lemma 17.1. $\square$

Lemma 17.14 (Natural alternating and symmetric tops). *If $T = A_s$ or S_s in its natural action, $s \geq 5$, then every quotient of an original pair lift has either one faithful general character or at most two faithful full-order linear characters.*

Proof. The invariant submodules of M_s are $0, C, I, M_s$, where C is constant and I is augmentation. Indeed from a nonconstant vector choose unequal coordinates i, j and equal coordinates k, l disjoint from them. Subtract its image under $(ij)(kl)$ to obtain $e_i + e_j$; these vectors span I . For odd s , $M_s = C \oplus I$ with I simple; for even s , $0 < C < I < M_s$ and I/C is simple. In either case the socle of every subquotient has multiplicity at most one in each type. If $\pi(N) = 1$, then $N \leq E$, and every abelian minimal normal of $Q = U/N$ lies in E/N , since its image in T would be abelian normal. Their multiplicities are at most one; the nonabelian socle shares the same one position. Theorem 8.1 gives one faithful character. If $\pi(N) \neq 1$, it contains A_s and $[E, A_s] \leq E \cap N$, using actual lifts from N in the commutator. From the displayed submodule list, $[E, A_s]$ has codimension at most one in E . Since $T/\pi(N)$ has order at most two, $|Q| \leq 4$. This is abelian binary and needs at most two full-order linear characters. The original extension was never assumed split. $\square$

Theorem 17.15 (Complete sixteen-pair estimate). *All actual pair actions with sixteen pairs and nonbinary top have a complete exponentially small forward row, with their original action weights and zero additional scalar.*

Proof. First choose any pair system for T , if one exists, and apply Lemma 17.8. Otherwise a minimal proper block of T has size four or eight. Its local action is primitive, since an internal nontrivial block system would give a smaller block of T . The size-eight case is Lemma 17.13. The size-four case has exact local A_4 or S_4 , and gives the actual transported wreath embedding in Lemma 17.11. If $H = 1$, Lemma 17.3(iv) and Lemma 17.1 give prefix two and tail four. If $H \neq 1$ and the local action is A_4 , Lemmas 17.10 and 17.11 give the same profile.

For local S_4 , put $A = M_0 N/N$ for each original N and $V = Q/A$. If V retains H , its prefix has size two by Lemma 17.10. If it kills H and is nonbinary or nonabelian, use its actual degree-eight sign action and Lemma 17.9. These give general prefix two and tail four; a binary Q can count all six positions cheaply. If V is abelian binary of rank at most three, use three binary-primary linear prefix positions and four relative positions. If its rank is four, Lemma 17.12 forces the full original translation base, and Lemma 17.11 improves the relative demand to two. Use four binary-primary linears and two relative positions. Both last cases have binary Q , since both A and V are binary.

If T is primitive, the classified list consists of the twenty affine degree-sixteen actions and natural A_{16}, S_{16} . The former use the complete original module-composition and literal normal data in Lemma 17.3(iv); the latter use Lemma 17.14. This exhausts the actual top, and all arguments apply to every original N in its pair lift.

At original width 32 the possible slopes are

$$2\gamma_5 + 4\beta, \quad 6\beta, \quad \frac{3}{2} + 4\beta, \quad 2 + 2\beta, \quad \gamma_5, \quad \frac{2}{3} \log_2 3.$$

Each is less than $4 - 1/64$. Exact integer versions, with common right-hand binary exponent 765, are

$$\begin{aligned} 5^{128} 38^{768} &< 2^{765} 25^{768}, & 38^{1152} &< 2^{765} 25^{1152}, \\ 2^{288} 38^{768} &< 2^{765} 25^{768}, & 2^{384} 38^{384} &< 2^{765} 25^{384}, \\ 5^{64} &< 2^{765}, & 3^{128} &< 2^{765}. \end{aligned}$$

Proposition 8.3 and Corollary 8.4 now sum all original normals and all actual width-thirty-two actions. Their constants are finite; every character tuple is fixed in its target before the same complete source J is chosen. Coordinate transports preserve $a(U) = |N_{S_{32}}(U)|$ and the original pointing weight. In particular no width-sixty-four estimate, quotient permutation degree, or six-ordinary-character estimate is used here. $\square$

18 Repeated natural heads and the smaller pair counts

For a finite group J , write $d_p(J) = \dim_{\mathbb{F}_p} J/(J'J^p)$. All head modules below come from an actual subgroup of the same complete $J \leq S_b$.

Proposition 18.1 (Repeated natural heads with central binary lifts). *Let*

$$R_d^A = \mathbb{F}_4^d : C_3, \quad R_d^S = (\mathbb{F}_2^2)^d : S_3,$$

where the same scalar, respectively natural, action is used on all d copies. Suppose Q has a specified central elementary binary subgroup C of rank c and $Q/C \cong R_d^A$ or R_d^S , with $d \geq 1$. For every complete $J \leq S_b$,

$$|\text{Epi}(J, Q)| \leq B_T(d) 2^{\max(c, d)b/2}, \quad B_A(d) = 2 \cdot 4^d, \quad B_S(d) = 6 \cdot 4^d.$$

No factor $|\text{Aut } Q|$ is required in this bound.

Proof. Put $r = d_2(J)$. In the A case let $F = J'$, and in the S case let $F = J''$. In either case set $B = J/F$ and $D = H_1(F, \mathbb{F}_2)$, with its honest B action. For every onto top map θ to C_3 or S_3 , let V_θ be the natural two-dimensional binary module. In the first case the odd part of the abelian B maps onto C_3 ; in the second case the odd part of the abelian B' does so. This is a normal odd subgroup with zero invariants on V_θ . Averaging and inflation–restriction therefore give

$$H^i(B, V_\theta) = 0 \quad (i \geq 0), \quad H^1(J, V_\theta) \cong \text{Hom}_B(D, V_\theta).$$

The vanishing in degree two is what retains all restrictions even when the source extension is nonsplit.

Let a_I be the multiplicities of represented natural simple types in the one semisimple head of D , and put $A = \sum_I a_I$. Their commuting fields are $\mathbb{F}_4$ in the A case and $\mathbb{F}_2$ in the S case. Each type corresponds to two, respectively six, concrete top maps. Onto restrictions onto d identical copies are ordered independent rows over that field. Every cohomology class has exactly 4^d literal cocycles, since the module has no fixed vectors. The target's first, respectively second, derived group is precisely its repeated binary module. Thus a map is onto exactly when this restriction is onto, giving the exact formulas

$$|\text{Epi}(J, R_d^A)| = 2 \cdot 4^d \sum_I \prod_{j=0}^{d-1} (4^{a_I} - 4^j), \quad (18.1)$$

$$|\text{Epi}(J, R_d^S)| = 6 \cdot 4^d \sum_I \prod_{j=0}^{d-1} (2^{a_I} - 2^j). \quad (18.2)$$

A term is zero when $a_I < d$. No independent copy of the source is assigned to a repeated constituent.

We need the simultaneous constraints

$$r + 2A \leq \lfloor b/2 \rfloor \quad (A \text{ case}), \quad r + A \leq \lfloor b/2 \rfloor \quad (S \text{ case}). \quad (18.3)$$

Choose bases of all the Hom spaces at once. Their evaluations give an onto map from D to $V_{\text{all}} = \bigoplus_I V_I^{a_I}$. Lift all its coordinates by the displayed cohomology isomorphism. The joint image is exactly $V_{\text{all}} : R$, where R is the joint image of the chosen top maps: the restriction of the same

F is onto V_{all} , so the full inverse image of R is obtained. In the A case $R \leq C_3^I$ is odd and acts without trivial constituents on the binary module. This target has no binary quotient. Pair its quotient map with the maximal elementary binary quotient C_2^r of J ; Goursat's lemma gives the full product. The preimage of $V_{\text{all}} \times C_2^r$ is an actual subgroup of S_b with an elementary binary quotient of rank $2A + r$. The actual subgroup abelianization bound [RDT25, Theorem 2.8] proves the first inequality.

In the S case put $N = R \cap C_3^I$ and $K \cong R/N \leq C_2^I$. The coordinate sign characters on N are nontrivial and have no trivial constituent; hence $[N, K] = N$. A complement to the normal odd Hall subgroup gives $R = N : K$ with elementary binary K . The maximal binary quotient of $V_{\text{all}} : R$ is exactly K . Pairing with C_2^r therefore gives $(V_{\text{all}} : R) \times C_2^{r-u}$, where $u = \dim K$; it is the full fibre product over all u binary characters. Pull back $(V_{\text{all}} : K) \times C_2^{r-u}$. Each natural type restricted to K is a nontrivial transvection module, so its coinvariant dimension is one. This actual subgroup has binary abelian quotient rank $A + u + r - u = A + r$, proving the second inequality.

For $q = 4$ or 2 ,

$$\sum_I \prod_{j < d} (q^{a_I} - q^j) \leq \sum_I (q^{a_I} - 1)^d \leq \left(\sum_I (q^{a_I} - 1) \right)^d \leq (q^A - 1)^d < q^{dA}$$

when the sum is nonempty, and it is zero otherwise. Above one fixed map to Q/C , homomorphic lifts to Q are absent or a torsor under $\text{Hom}(J, C)$, of size 2^{cr} . Onto lifts are only a subset. Consequently the two exponents are bounded by $cr + 2dA$ and $cr + dA$, respectively. Equations (18.3) bound both by $\max(c, d)\lfloor b/2 \rfloor$. This proves the assertion while retaining every lift obstruction before the positive enlargement. $\square$

Corollary 18.2 (Finite menus with repeated-head entries). *Fix a finite menu of original transitive actions U of width w . For every original $N \triangleleft U$, suppose its quotient has either a fixed mixed character tuple with slope less than $h(w)/8$, or a specified central repeated-head certificate as in Proposition 18.1 with $\max(c, d)/2 < h(w)/8$. Then the complete menu has one exponentially small forward row and zero scalar, with original action weights.*

Proof. Fix one choice at every original normal subgroup. The finite set has a common positive strict gap; sum the estimates of Propositions 8.2 and 18.1 on the same J , and apply Lemma 7.2 to $\Delta(n, b)Z_J(U)/a(U)$. Different original normals remain different summands. Repeated-head entries already count literal epimorphisms and do not receive an additional automorphism factor. $\square$

Lemma 18.3 (A binary rank tail on complete sources). *For each fixed $\epsilon > 0$,*

$$\#\{J \leq S_b : d_2(J) > (1/4 + \epsilon)b\} \leq 2^{b^2/16 - \epsilon^2 b^2 + O_\epsilon(b^{3/2})}.$$

The same estimate, with an additional $O(b \log(b+2))$ in the exponent, counts complete J having a normal subgroup F of index three with $d_2(F) > (1/4 + \epsilon)b$.

Proof. For any integer $t \geq 0$, the graphs of all maps $J \rightarrow C_2^t$, with the latter on t fixed disjoint pairs, are subgroups of S_{b+2t} . Projection recovers J and its graph. There are $2^{td_2(J)}$ maps. Apply the coarse bound (1.1) and take t within one of $2\epsilon b$. The exact main exponent is

$$\frac{(b+2t)^2}{16} - (1/4 + \epsilon)bt = \frac{b^2}{16} - \epsilon^2 b^2 + \frac{(t - 2\epsilon b)^2}{4}.$$

For the second assertion, every index-three overgroup of a fixed $F \leq S_b$ is generated by F and one permutation, so there are at most $b!$ possibilities. This is a positive union bound over all such actual kernels; no eligible kernel multiplicity is divided out. $\square$

Proposition 18.4 (All six-pair actions). *Every transitive action on twelve points with six pairs has a complete exponentially small forward row and a quadratic-exponentially small scalar, with its original action weight.*

Proof. The six-block certificates in Appendix A give 182 physical action classes and 2476 literal normal records. Here is the constructive coverage behind this finite premise. The 56 subgroup representatives in S_6 contain the identity and are closed, up to explicit conjugacy, under adjoining every double-coset representative. If $x = h_1 g h_2$, then $\langle H, x \rangle = \langle H, g \rangle$; induction on generators therefore proves completeness. Sixteen are transitive. The 2825 distinct subspaces of $\mathbb{F}_2^6$, the exact Gaussian sum, give all 112 invariant-kernel interfaces. On each quotient module, evaluating a presentation on corrected generator lifts gives the linear cocycle equations. The number of base-conjugacy classes is exactly $2^{\dim Z^1 - \dim M + \dim M^T}$, proving completeness of the listed complement graphs and their full preimages, including nonsplit original lifts. The 224 transitive block certificates are identified by actual degree-twelve conjugacy into the 182 physical actions, with their original symmetric normalizers.

Every original normal list is closed under adjoining the normal closure of each conjugacy-class representative and contains the identity. Since a normal subgroup is generated by its conjugacy classes, this proves exhaustive normal coverage. The certificate at each normal is either a faithful irreducible tuple consisting of at most one linear and one general character (two linears also allowed), or a literal onto map with exactly that kernel to one of the five fixed models

$$C_2^3, \quad B_{32}, \quad C_2^e \times R_2^A \quad (0 \leq e \leq 2).$$

On two four-point blocks, B_{32} is generated by V_4^2 and (23)(67), of order 32. Kernel intersections of the character tuples and the explicit model maps verify these alternatives; labels or group orders alone are not used.

Call the complete J cold if $r = d_2(J) \leq 13b/50$. Character entries cost at most $|\text{Aut } Q|15^{b/3}$. Every map to a binary target factors through the canonical maximal binary quotient of J . That quotient has generator rank r , so the two binary models have at most $2^{5r} \leq 2^{13b/10}$ maps. Proposition 18.1 bounds each repeated-head model by $32 \cdot 2^b$. As $15 < 16$, summing every original normal gives a constant C_U with

$$Z_J(U) \leq C_U 2^{4b/3} \quad (J \text{ cold}).$$

This has strict margin $3/2 - 4/3 = 1/6$ at original width twelve. For arbitrary J , its generator number is at most $\log_2(b!)$; the fixed targets have orders at most $46080 < 2^{16}$. Thus $Z_J(U) \leq \nu(U)(b!)^{16}$, where $\nu(U)$ is its normal count. Lemma 18.3 with $\epsilon = 1/100$ bounds the hot complete-source count by $2^{b^2/16 - b^2/10000 + O(b^{3/2})}$. The full hot fibre and original normalization are consequently at most

$$\sum_U \frac{\nu(U)(b!)^{16}}{b!a(U)p_n G_{\lfloor n/2 \rfloor}} 2^{b^2/16 - b^2/10000 + O(b^{3/2})} = 2^{-\Omega(n^2)}, \quad n = b + 12.$$

The factorials cost only $O(n \log(n+2))$. The cold sum is $\Delta(n, b) \sum_U C_U 2^{4b/3}/a(U)$ and is exponentially small by Lemma 7.2. Fuse the whole family once, using the existence of any hot accepted pointing and its cold complement. There is one complete terminal on the cold row. $\square$

Lemma 18.5 (A cyclic dual on a common derived source). *Let $X = V : R$, where V is a binary vector space of dimension d , R acts faithfully and V^* is cyclic. Suppose, for $\ell = 1$ or 2 , that $R^{(\ell)} = 1$ and the odd part of $R^{(\ell-1)}$ has zero invariants on V . Then*

$$|\text{Epi}(J, X)| \leq 2^d |\text{GL}_d(2)| 2^{b/2}.$$

Proof. The hypotheses give $X^{(\ell)} = V$. Every map therefore sends the same $F = J^{(\ell)}$ onto V . Put $B = J/F$ and $D = H_1(F, \mathbb{F}_2)$. The odd part of $B^{(\ell-1)}$ maps onto the stipulated odd group under every onto top map, so averaging gives $H^i(B, V_\theta) = 0$ for all i . Onto restrictions $D \rightarrow V_\theta$ dualize to injections $V_\theta^* \rightarrow D^*$. Fix one cyclic vector in V^* . Its image f determines its whole image module $\mathbb{F}_2[B]f$. For a fixed f , there are at most $|\mathrm{GL}_d(2)|$ linear identifications; faithfulness determines the top map from each identification. Thus every top map and onto restriction together has at most $|\mathrm{GL}_d(2)|2^{\dim D}$ possibilities. Each restriction has at most $|V|$ cocycle lifts, and the actual $F \leq S_b$ has $d_2(F) \leq b/2$. This proves the bound for reducible and nonsemisimple modules as well. $\square$

Proposition 18.6 (All four-pair actions). *Every nonbinary transitive action on eight points with four pairs has a complete exponentially small forward row and a quadratic-exponentially small scalar, with its original action weight.*

Proof. Its top is A_4 or S_4 . In $M = \mathbb{F}_2^4$, the invariant modules are exactly $0, C, I, M$, with C constant and I augmentation: weight-two vectors span I , weight-one or weight-three orbits span M , and weight four is constant. For each kernel, try all 256 pairs of binary corrections to two fixed top generators, and require that the resulting whole kernel be the selected one. Every projecting subgroup contains such lifts, so this is exhaustive. The four-block certificates retain the ten transitive classes, their original normalizers and all 72 original normals. Normal completeness follows from intersection closure of irreducible character kernels, as in Lemma 17.3. Five classes are

$$C_2 \times A_4, \quad \mathrm{SL}_2(3), \quad S_4 \text{ in degree eight}, \quad C_2 \times S_4, \quad \mathrm{GL}_2(3).$$

The other five, denoted here $Y_1, \dots, Y_5$, have binary kernels I, M, I, I, M , respectively, with A_4 top in the first two cases and S_4 top in the last three. The two augmentation lifts over S_4 are the ordinary lift and the graph whose base parity equals the top sign; the latter need not split over I .

For these five, C is the unique minimal normal subgroup. The only minimal submodule of their binary kernel is C , and any normal subgroup disjoint from that kernel centralizes it. The top action on I and M is faithful, so such a subgroup is trivial. A character nontrivial on C , induced and then restricted to an irreducible constituent, is therefore faithful on Y_i . Thus $|\mathrm{Epi}(J, Y_i)| \leq |\mathrm{Aut} Y_i|5^{b/3}$. The exact central quotients recorded by the literal maps are

$$Y_1/C = R_2^A, \quad Y_2/C = X_c, \quad Y_3/C = Y_4/C = X_g, \quad Y_5/C = X_s,$$

where, on $V = \mathbb{F}_4^2$, a is common multiplication by ω , t common Frobenius and s coordinate swap, and

$$X_c = V : \langle a, s \rangle, \quad X_g = V : \langle a, ts \rangle, \quad X_s = V : \langle a, t, s \rangle.$$

Because every nontrivial normal contains C , in each case $Z_J(Y_i) = |\mathrm{Epi}(J, Y_i)| + Z_J(Y_i/C)$ exactly.

For X_c, X_s , the module is cyclic over $\mathbb{F}_4[s]$ with unique proper nonzero submodule the diagonal line. The invariant trace form $\mathrm{Tr}(x_1 y_1^2 + x_2 y_2^2)$ identifies it with its binary dual. For X_g , the three invariant $\mathbb{F}_4$ lines are $y = cx$ with $c^3 = 1$; any two are complementary natural S_3 modules. Their commuting field is $\mathbb{F}_2$, and the matrix algebra $\mathrm{Mat}_2(\mathbb{F}_2)$ shows that two natural copies have a cyclic dual (take components forming a basis). Thus Lemma 18.5 applies with $\ell = 1$ for X_c and $\ell = 2$ for the other two, giving onto bounds $C_0 2^{b/2}$, $C_0 = 16|\mathrm{GL}_4(2)|$. Their exact full normal menus are

$$\begin{aligned} Z_J(X_c) &= |\mathrm{Epi}(J, X_c)| + Z_J(C_2 \times A_4), \\ Z_J(X_g) &= |\mathrm{Epi}(J, X_g)| + 3|\mathrm{Epi}(J, S_4)| + Z_J(S_3), \\ Z_J(X_s) &= |\mathrm{Epi}(J, X_s)| + Z_J(C_2 \times S_4). \end{aligned}$$

Indeed in the first and third case every nontrivial normal contains the diagonal line. In the second case the three line kernels give three S_4 quotients, and all other proper normals contain V . These yield faithful degree-six comparators $C_2 \times A_4, S_4, C_2 \times S_4$ (padding S_4), with multipliers 1, 3, 1. Since both $1/2$ and γ_5 are less than one, Theorem 7.3 handles $Y_2, \dots, Y_5$.

For $C_2 \times T$, $T = A_4, S_4$, normals not containing the natural V_4 lie in the central C_2 . Put $z = 2^{d_2(J)}$. The complete remaining normal sum is $z|\text{Epi}(J, A_4)|$ in the alternating case and $(z-1)|\text{Epi}(J, S_4)|$ in the symmetric case: in the latter, the central binary row must be independent of the existing sign. The joint facets in Proposition 18.1, with $d = 1$, bound these sums by $8 \cdot 2^{b/2}$ and $24 \cdot 2^{b/2}$ respectively. Their complete quotient intervals are exactly $Z_J(C_2 \times C_3)$ or $Z_J(C_2 \times S_3)$, with faithful five-point representations. This proves the finite comparator estimate for these two classes. For $\text{SL}_2(3)$ and $\text{GL}_2(3)$ the center C_2 is unique minimal normal, so the same faithful-character argument and exact normal interval give comparators A_4, S_4 of degree four, with slope $\gamma_5 < 1$. The nonnatural S_4 itself has that comparator with zero error. This handles all classes except Y_1 .

For $Y = Y_1$, the repeated quotient has five distinct $\mathbb{F}_4$ -line kernels and therefore

$$Z_J(Y) = |\text{Epi}(J, Y)| + |\text{Epi}(J, R_2^A)| + 5|\text{Epi}(J, A_4)| + Z_J(C_3). \quad (18.4)$$

The first-derived proof of Proposition 18.1 gives one common head with natural multiplicities a_I and $r = d_2(J)$. For each corresponding $\theta : J \rightarrow C_3$, put $H = \ker \theta$. Averaging on $J/H = C_3$ gives

$$H^1(J, \mathbb{F}_2) = H^1(H, \mathbb{F}_2)^{C_3}, \quad H^1(J, V_\theta) = \text{Hom}_{C_3}(H_1(H, \mathbb{F}_2), V_\theta).$$

Comparing the latter with the same common-head Hom space shows that $H_1(H, \mathbb{F}_2)$ has trivial multiplicity r and natural multiplicity a_I . Thus

$$d_2(\ker \theta) = r + 2a_I. \quad (18.5)$$

There are at most $b/4$ represented types, since $2\sum_I a_I \leq d_2(J') \leq b/2$.

Declare J hot when *some* normal subgroup of index three has rank $> 51b/200$; test all such subgroups, even those whose natural type is absent. On the cold set, (18.5) and (18.1) give

$$5|\text{Epi}(J, A_4)| \leq 10b2^{51b/200}, \quad |\text{Epi}(J, R_2^A)| \leq 8b2^{51b/100}.$$

Together with $Z_J(C_3) \leq 3^{b/3}$ and the faithful-character bound on Y , the exact full sum (18.4) has slope at most $\gamma_5 < 1$, up to a fixed linear factor in b . For all sources that full sum is at most $C(1+b)2^b$: use $a_I \leq b/4$ in the repeated-head formula. Lemma 18.3, now with $\epsilon = 1/200$ and its index-three clause, gives the complete hot count $2^{b^2/16 - b^2/40000 + O(b^{3/2} + b \log(b+2))}$. Multiplying by the whole fibre and by the original factor $1/(384b!p_n G_{\lfloor n/2 \rfloor})$, $n = b+8$, leaves a quadratic-exponentially small scalar. The cold row is exponentially small by Lemma 7.2. The divisor 384 is the original S_8 normalizer: Y is normal in $C_2 \wr S_4$, and its characteristic C forces every normalizer to preserve those four pairs. The finite union of all ten classes is fused once, with all original weights and complete cold terminals retained. $\square$

18.1 The three exceptional five-point block counts

Theorem 18.7 (Exact affine-five exceptions). *Every action with exact primitive affine local degree five and $s \in \{2, 3, 4\}$ has a complete exponentially small forward row and a quadratic-exponentially small scalar.*

Proof. Write $M = \mathbb{F}_5^s$, $E = U \cap M$, $R = U/E \leq C_4 \wr T$. Since $s \leq 4$, the latter group has order prime to five. Transitivity forces $E \neq 1$, since otherwise $5 \nmid |U|$. Schur–Zassenhaus first splits $U = E : R$, then conjugates its complement into the actual monomial complement in $M : R$. This does not split $R \rightarrow T$. The semisimple monomial modules M, M^* are cyclic: the orbit of one coordinate vector spans the module. Every section $W = E/I$ and its dual are therefore cyclic and can be realized as actual submodules of M .

Let $D = \ker(R \rightarrow T)$ and $K = \ker(R \rightarrow \mathrm{GL}(W))$ for $W \neq 0$. Every coordinate projection of this realized W is nonzero by transitivity. A diagonal element trivial on W is therefore one, so $K \cap D = 1$ and K embeds normally in T . Its coordinate orbits have one common size $q \mid s$; a fixed vector on a monomial orbit is determined by one coordinate, giving $\dim W \leq s/q$. Thus $K = 1$ if $\dim W \geq 2$ at $s = 2, 3$, or if $\dim W \geq 3$ at $s = 4$. The only further possibility is $s = 4, \dim W = 2, q = 2$. Then $K \leq C_2^2$ and $T \leq D_8$, so R is binary. The normal subgroup K has a series of at most two central C_2 factors in R : choose an involution in $K \cap Z(R)$ and then use centrality of a normal order-two group in the quotient. The same holds after quotienting any normal $K_0 \leq K$.

We need a joint-source onto estimate. If $X = W : S$, with S a faithful $5'$ -group on $W = \mathbb{F}_5^d$ and W^* cyclic, then

$$|\mathrm{Epi}(J, X)| \leq 5^d |\mathrm{GL}_d(5)| 5^{b/5} \quad (J \leq S_b). \quad (18.6)$$

Indeed fix the single characteristic source subgroup $F = O^{5'}(J)$ and $B = J/F$. Residuals commute with epimorphisms, so every map onto X sends F onto W and induces a map $B \rightarrow S$. Put $A = H_1(F, \mathbb{F}_5)$. A surjective equivariant restriction dualizes to an injection $W^* \hookrightarrow A^*$. The image of one fixed cyclic vector determines its whole cyclic B -submodule. For each image there are at most $|\mathrm{GL}_d(5)|$ identifications; faithfulness determines the top map from each identification. Thus this bounds all top maps and restrictions jointly by $|\mathrm{GL}_d(5)||A^*|$. The remaining lift fibre is empty or a torsor for $Z^1(B, W)$; coprime averaging gives $|Z^1(B, W)| \leq |W|$. Finally $d_5(F) \leq b/5$ in its actual permutation action, proving (18.6). A central C_2 lift has at most $|\mathrm{Hom}(J, C_2)| \leq 2^{b/2}$ choices over a fixed map, whether split or not. The two central steps above therefore cost at most 2^b in total.

Fix now $N \triangleleft U$ and $I = N \cap E$. If $I = E$, these terms sum exactly to $Z_J(R)$. If $\dim(E/I) = 1$, retain the whole comparator U/I . For this line let $\chi : R \rightarrow \mathbb{F}_5^*$ be its action. The map (χ, π_T) is injective since $K \cap D = 1$, so $U/I \hookrightarrow \mathrm{AGL}_1(5) \times T$ has a faithful action of degree $5 + s \leq 4s$. For $\dim(E/I) \geq 2$, the normal N/I is disjoint from W and centralizes it. It lies in $W \times K$ and projects injectively to the $5'$ -group K ; its graph into W is trivial. Hence $N/I = K_0 \triangleleft R$. Apply (18.6) if $K = 1$, or apply it after the at most two central binary steps otherwise.

This proves, with a fixed list $\mathcal{Q}_U$ consisting of R and all literal line quotients U/I , that

$$Z_J(U) \leq \sum_{Q \in \mathcal{Q}_U} Z_J(Q) + C_U 2^{\eta_s b}, \quad C_U = \nu(U) 5^s |\mathrm{GL}_s(5)|,$$

where $\eta_2 = \eta_3 = \log 5/5$ and $\eta_4 = 1 + \log 5/5$. Equal abstract quotients at different I remain different entries of this fixed list. Pad every comparator to faithful degree $v = 4s$.

For clarity the finite comparator argument extends directly to this fixed list. Let $m = |\mathcal{Q}_U|$ and $S_U(J) = m^{-1} \sum_{Q \in \mathcal{Q}_U} Z_J(Q)$. Jensen and Lemma 7.1 give, for every integer $q \geq 1$,

$$\sum_{J \leq S_b} S_U(J)^q \leq m^{-1} \sum_Q \sum_J Z_J(Q)^q \leq s_{b+qv}.$$

The proof of Theorem 7.3 uses precisely this moment inequality and positivity, so applies with statistic S_U and multiplier m . Here $h(5s) = 10, 14, 20$ and $v = 8, 12, 16$, respectively; $v < h(5s)$

and $\eta_s < h(5s)/8$, since $\log 5/5 < 7/15$. The finite union therefore has the claimed bounds with its original physical weights and one complete residual terminal. $\square$

18.2 Three pairs and the complete degree-sixteen slice

Proposition 18.8 (Three original pairs). *Every nonbinary transitive action on three pairs has a complete contractive forward row and negligible scalar.*

Proof. Its top is C_3 or S_3 , and its binary permutation module is $\mathbb{F}_2^3 = C \oplus A$, with constant line C and augmentation plane A . The normal C_3 acts trivially on C and irreducibly on A , so the only invariant kernels are $0, C, A, C \oplus A$. Above each kernel, subgroups are graphs of cocycles in the quotient module. Coprime averaging kills all C_3 cocycles. For S_3 , the odd normal subgroup has no fixed vector on A , so $H^1(S_3, A) = 0$, while $H^1(S_3, C) = \text{Hom}(S_3, C_2)$ is generated by sign. This supplies the complete graph list, including nonsplit original lifts.

The transitive actual groups are $C_6, A_4, C_2 \times A_4$ over C_3 ; and regular $S_3, C_2 \times S_3$, two distinct degree-six actions of S_4 , and $C_2 \times S_4$ over S_3 . At zero kernel only the sign graph is transitive; at nonzero kernel the coordinate projections are full. The two S_4 graphs remain separate actual actions throughout the pointing.

The targets $C_6, A_4, S_3, C_2 \times S_3, S_4$ have specified faithful representations of degrees at most five, less than six. Use the target itself in Theorem 7.3, with $C = 0$. For $U = C_2 \times L$, $L = A_4$ or S_4 , let V be the standard V_4 . Any normal subgroup not containing V centralizes it; normality then forces that subgroup into the central C_2 . Normals containing V contribute exactly $Z_J(C_2 \times (L/V))$, whose comparator has faithful degree five. The two other quotients are L and $C_2 \times L$. Proposition 18.1, with repeated rank one and central rank at most one, bounds both their onto-map counts by a fixed constant times $2^{b/2}$. Thus

$$Z_J(U) \leq Z_J(C_2 \times (L/V)) + C_U 2^{b/2}.$$

Its exponent $1/2 < 6/8$ is strict. The finite comparator theorem applies with the original normalizer for each of the eight actions. $\square$

Theorem 18.9 (Every nonbinary action of degree sixteen). *The complete union of nonbinary transitive degree-sixteen actions has a contractive forward row and negligible scalar. In particular this covers every eight-pair action and every zero-ternary action on four four-point blocks.*

Proof. The finite action certificate contains the full nonbinary slice of the degree-sixteen transitive classification, with 527 original permutation representatives and their original normalizers. Its completeness premise is the stated transitive classification [Hul05, Hul23]. Each of its 9249 literal normal entries is retained separately. To justify normal completeness, the certificate lists all conjugacy classes, their exact sizes and normal closures. The supplied normals contain one and are closed under adjoining every such normal closure. Every normal subgroup is generated by the closures of the classes it contains, so this join closure proves that no original normal is omitted.

For 518 actions all quotients have a fixed mixed character tuple, and for all but a specified interval or head quotient of each remaining action the same is true. A mixed tuple has l linear and t general slots with literal kernel intersection equal to the original normal, and satisfies

$$3^{64l} 5^{64t} \leq 2^{381}.$$

Proposition 8.2 gives coefficient at most $381/192 = 2 - 1/64$, strictly below the physical reserve $16/8$. All target automorphism factors are fixed finite constants.

Two actions have a specified original derived subgroup with quotient C_2^4 , concretely represented on eight points. Precisely the normal interval above that subgroup is retained as the whole $Z_J(C_2^4)$; every other original normal has a mixed tuple. Theorem 7.3 applies with $v = 8$ and $\eta = 2 - 1/64$.

Seven actions have, among their original normals, eight exceptional quotients. Each is a central elementary binary extension of a repeated A_4 or S_4 model: the certified whole centre has rank c , and the quotient has repeated rank d , with $32 \max(c, d) \leq 127$. The maps and central preimages are literal; no splitting is inferred. Proposition 18.1 bounds each onto fibre by a fixed constant times $2^{\max(c, d)b/2} \leq 2^{3b/2}$. All other normals are mixed. Original-weight pointing and Lemma 7.2 give their direct exponential row. The finite union, with the two comparator scalars, proves the theorem. No certificate is selected after inspecting the complete source J . $\square$

19 The two zero-ternary frame exceptions

Put

$$z_2(G) = \max_{L \triangleleft G} \dim_{\mathbb{F}_2} L/[L, G]L^2, \quad \mu_T(M) = \max_{B \leq M \text{ invariant}} \dim_{\mathbb{F}_2} B/[B, T].$$

The second quantity is the maximum dimension of a trivial homogeneous section of the one actual module M .

Lemma 19.1 (A fixed-frame bound). *If $1 \rightarrow E \rightarrow U \rightarrow T \rightarrow 1$ has elementary binary kernel $E \leq M$ as a T -module, then $z_2(U) \leq \mu_T(M) + z_2(T)$. Moreover*

$$z_2(U) = \max_{N \triangleleft U} \text{rank } \Omega_1(Z(U/N))_2.$$

Proof. For $K \triangleleft U$, let L be its image in T , put $N = [K, U]K^2$ and $I = K \cap E$. The image of N is exactly $[L, T]L^2$, so there is an exact sequence

$$0 \longrightarrow I/(I \cap N) \longrightarrow K/N \longrightarrow L/[L, T]L^2 \longrightarrow 0.$$

Its first term is a trivial quotient of the actual submodule I of M , proving the bound. Each $K/[K, U]K^2$ is central elementary binary in its corresponding quotient of U . Conversely the preimage of any central elementary binary subgroup in U/N has relative head surjecting onto that subgroup. Taking maxima proves the identity. $\square$

Lemma 19.2 (Noncentral socle types in the two frames). *Suppose U has exact local A_4 or S_4 on $s \in \{8, 16\}$ blocks and zero ternary linear kernel. Let $M = (\mathbb{F}_2^2)^s$, $E = U \cap M$ and $T = U/E$, its actual faithful block top. Every nontrivial binary simple type in the socle of every quotient has Schur demand at most $5s/8$, and every odd-primary type has demand at most $\lfloor s/3 \rfloor$. Thus every quotient has a faithful tuple of at most $5s/8$ characters if $z_2(U) \leq 5s/8$.*

Proof. In $Q = U/N$, the whole binary socle A is a semisimple T -module: the normal binary $\bar{E}$ acts trivially on each simple type. Filter A by $A \cap \bar{E}$. For a type X of binary dimension d and Schur dimension e , its multiplicities satisfy

$$m_E \leq \lfloor 2s/d \rfloor, \quad m_T \leq \lfloor s/(2d) \rfloor.$$

The second bound uses the whole elementary outer section of T . These multiplicities add, since the types on the two sides may agree. For $d \geq 4$, their sum is at most $5s/8$. For $d = 3, e = 3$, dividing the sum by three and rounding up also suffices. For $d = 3, e = 1$, the image is C_7 ; its normal

kernel is transitive, since its orbit number divides both seven and s . Restricted to that kernel, X^{m_E} is trivial of dimension $3m_E$. Lemma 16.1 bounds this dimension by s , giving $m_E \leq \lfloor s/3 \rfloor$ and $m_T \leq \lfloor s/6 \rfloor$.

For $d = 2, e = 2$ the demand is at most $\lceil (s + s/4)/2 \rceil = 5s/8$. If $d = 2, e = 1$, the image is C_3 and its normal kernel K is transitive. At a block stabilizer, restriction to K either keeps a nontrivial local C_3 or S_3 , or kills the local C_3 in the A_4 case. In the first case take a Sylow 3-subgroup P of K . At a P -fixed block it is Sylow in the stabilizer and its local image is C_3 , with no fixed vector. Every nonfixed orbit contributes at most two invariant dimensions. Exactness of odd-order invariants on sections gives $2m_E \leq 2\lfloor s/3 \rfloor$. Adding $m_T \leq s/4$ suffices at both counts. In the second case the induced frame restricted to K is two copies of the honest permutation module $\mathbb{F}_2^s$. Its transitive Sylow 2-subgroup gives the scalar submodule capacity $\binom{a}{\lfloor a/2 \rfloor} = 3s/8$ for $s = 2^a \in \{8, 16\}$. Filtering a trivial section through the two copies gives $m_E \leq 3s/8$, and again $m_E + m_T \leq 5s/8$.

For odd p , the entire p -socle meets $\bar{E}$ trivially and embeds in a quotient of T , so its rank is at most $\lfloor s/p \rfloor$. The only remaining binary type is the trivial type, whose maximum rank over the normal quotients is exactly $z_2(U)$ by Lemma 19.1. Apply the whole-socle criterion. $\square$

The following finite inputs concern tops and induced linear frames, and do not enumerate all extensions U above a frame. They are supplied with literal permutation, module and kernel witnesses.

Lemma 19.3 (Finite frame inputs). *The zero-ternary frame certificates have the following properties.*

- (i) *Every nonbinary transitive top T of degree eight satisfies $z_2(T) \leq 2$. Every frame induced from an onto map of a point stabilizer to C_3 or S_3 satisfies $\mu_T(M) \leq 2$.*
- (ii) *For each degree-sixteen top having four four-point blocks, exact local A_4 or S_4 , and zero internal ternary kernel, every induced C_3/S_3 frame admits a Sylow 3-subgroup P and a 2-element $g \in N_T(P)$ with $z_2(T) + \dim(M^P)^g \leq 10$.*
- (iii) *Every primitive degree-sixteen top and every eligible induced frame admit an odd subgroup A with $z_2(T) + \dim M^A \leq 10$.*
- (iv) *The primitive degree-eight actions are the three affine groups $2^3 : C_7$, $2^3 : (C_7 : C_3)$, $\text{AGL}_3(2)$ and $\text{PSL}_2(7)$, $\text{PGL}_2(7)$, A_8, S_8 . Their normal subgroup orders are respectively*

$$(1, 8, 56), (1, 8, 56, 168), (1, 8, 1344), (1, 168), (1, 168, 336), (1, 20160), (1, 20160, 40320).$$

Except for the first group, a Sylow 3-subgroup has orbit lengths 1, 1, 3, 3 in each actual action.

Proof. The degree-eight top alphabet consists of the seventeen imprimitive classes constructed by the four-pair and two-four-block Goursat and lift certificates, and the seven primitive actions of the stated primitive classification. All normal subgroups are retained for the relative-rank check. All onto point-stabilizer maps are recovered by their literal kernels; for a fixed kernel, automorphisms of C_3 are realized by simultaneous Frobenius and automorphisms of S_3 are inner. This yields all thirty frames on twenty eligible tops.

For every one of these frames the supplied binary invariant-submodule list contains zero and is closed under adding every cyclic submodule. The latter are constructed from all vector orbits under the actual generator matrices. Any omitted invariant submodule would contain an included

one maximal under containment; adjoining the cyclic submodule of a vector outside it contradicts maximality. This proves completeness, and direct commutator ranks give the stated maximum two.

For (ii), the complete induced-frame, kernel and complement construction gives twenty-nine four-by-four top actions, including their nonsplit lifts. Each has exactly one eligible stabilizer kernel. The supplied P, g witnesses give the inequality directly; no invariant-submodule enumeration in dimension 32 is required. For (iii), all twenty-two primitive degree-sixteen entries are checked, giving nine eligible stabilizer kernels on seven tops and the stated odd-subgroup witnesses. The primitive completeness premise is [RD05, HRDR23]; parts (i), (ii) use the separate constructive completeness arguments just described. The normal orders and actual Sylow orbits in (iv) are direct checks on the same seven primitive representatives. Thus these are finite mathematical premises with independently checkable completeness certificates, rather than an inference from their cardinalities. $\square$

Theorem 19.4 (Complete zero-ternary eight and sixteen frames). *Every exact-local A_4/S_4 action with zero ternary kernel and eight or sixteen blocks has a faithful character cover on every normal quotient of length five or ten, respectively. Their entire original-normal menus satisfy Corollary 8.4.*

Proof. By Lemmas 19.1 and 19.2, it suffices to bound $\mu_T(M) + z_2(T)$ by five or ten. At degree eight Lemma 19.3(i) gives four.

We record two tools for degree sixteen. Odd-order invariants are exact, so for an odd subgroup P , $\mu_T(M) \leq \mu_{N_T(P)}(M^P)$. If g is a 2-element normalizing P , the binary cyclic module $M^P \downarrow_{\langle g \rangle}$ is a sum of modules over a chain ring. The head and socle have equal dimension, and submodules cannot have larger socle. Hence

$$\mu_T(M) \leq \dim(M^P)^g.$$

Also, if P is Sylow three in a transitive top of power-of-two degree, a P -fixed block contributes zero to M^P : the local onto C_3/S_3 map has nontrivial C_3 image there. Each nonfixed P -orbit contributes at most two invariant dimensions.

Suppose first T preserves eight pairs, with kernel K and top $S \leq S_8$. If $K \neq 1$, it projects onto every pair. The local map from a point stabilizer kills its intersection with K , because C_3, S_3 have no nontrivial normal 2-subgroup. It descends onto the stabilizer of a pair in S , giving one of the degree-eight frames M_0 . Induction in stages and the length-two filtration of the binary two-point permutation module give a filtration of M with two inflated copies of M_0 . Thus $\mu_T(M) \leq 2\mu_S(M_0) \leq 4$. Furthermore Lemma 19.1 on $K \leq \mathbb{F}_2^8$ gives $z_2(T) \leq 3 + z_2(S) \leq 5$, using the scalar induced capacity three in degree eight. The sum is at most nine. If $K = 1$, then $T \cong S$ and $z_2(T) \leq 2$. A Sylow three has at most two nonfixed orbits on the eight pairs, each lifting to two orbits on the sixteen points, since an odd pair stabilizer fixes its pair pointwise. Hence $\mu_T(M) \leq 8$, giving ten.

Otherwise a minimal nontrivial block of T has size four or eight, or T is primitive. For size four let F be its internal binary translation kernel, let D be its internal ternary kernel and let $S \leq S_4$ be the top. If $D \neq 1$, retained- F quotients have character cover two by Theorem 16.5. For quotients killing F , the central elementary binary socle injects after removing the odd D into a quotient of $C_2 \wr S \leq S_8$. Therefore $z_2(T) \leq 4$. If $S = A_4$ or S_4 , a Sylow three has at most three nonfixed orbits on the sixteen points: over its three cycled four-blocks the moving local triples give one orbit of length nine and their fixed points one orbit of length three; the fourth block contributes at most one further orbit. Thus $\mu_T(M) \leq 6$. If S is binary, let P be Sylow three of the full block kernel. It is also Sylow in T , and Frattini gives a normalizer transitive on the four

blocks. Each block contributes the same dimension $d \leq 2$ to M^P . The normalizer modulo P is binary, so the local d -space has a series of trivial factors. Induction gives d copies of the four-point permutation module in a filtration of M^P . Their scalar capacity is two, giving $\mu_T(M) \leq 2d \leq 4$. If instead the internal $D = 1$, use Lemma 19.3(ii) and the normalizer tool.

For a minimal block of size eight, the two-block kernel is a Goursat subdirect product of one of the seven primitive locals in Lemma 19.3(iv). The first, $2^3 : C_7$, has no eligible point-stabilizer map. For each other local a Sylow three has four nonfixed orbits on the sixteen points, so $\mu_T(M) \leq 8$. For $\text{PGL}_2(7)$ or S_8 , the intersection of the kernel with the square of the simple socle is semisimple, and the quotient embeds in $C_2 \wr C_2$ of degree four; hence $z_2(T) \leq 2$. For the simple locals the block kernel is semisimple, so its relative binary rank is zero. For the two eligible affine locals, the displayed normal menus show that every intermediate Goursat kernel is either a graph of the local group, the full product, or contains both translations with odd or common simple quotient. The translations are then a semisimple sum of nontrivial simple modules, including all automorphism twists. They have no trivial section, and the quotient has relative binary rank zero. Thus the kernel has $z_2 = 0$ and $z_2(T) \leq 1$. Every case gives a sum at most ten. The primitive degree-sixteen case is Lemma 19.3(iii).

These cases exhaust the top: a primitive binary group has prime degree, so no omitted binary primitive local occurs at sizes four or eight. The original extension U is arbitrary throughout. Finally $5^{64 \cdot 5} < 2^{24 \cdot 32 - 3}$ and $5^{64 \cdot 10} < 2^{24 \cdot 64 - 3}$ give the character-forward margin at the two physical widths. $\square$

20 Exhaustion of the nonbinary orbit actions

Theorem 20.1 (Complete nonbinary action reduction). *Let $\mathcal{X}_n$ consist of the actual subgroups of S_n with an orbit action outside the following list:*

$$\text{binary transitive actions,} \quad \text{natural } C_3, A_4, S_3, \quad \text{fixed points.}$$

There are positive constants c, c' and complete kernels $K(n, b)$, with $K(n, b) = 0$ for $b \geq n$, such that eventually

$$\frac{|\mathcal{X}_n|}{L_n} \leq \eta(n) + \sum_{b < n} K(n, b) \frac{s_b}{L_b}, \quad \sum_{b < n} K(n, b) = O(2^{-cn}), \quad \eta(n) = O(2^{-c'n^2}).$$

Every forward term retains the complete complementary subgroup and the original permutation-action weight. Thus every subgroup outside this counted union has only the displayed orbit actions.

Proof. We first prove that every transitive action outside the list meets one of the proved consumers on its entire original normal menu. This is an action-by-action exhaustion, preceding the counting union.

For a primitive action, a nonaffine socle is covered by Theorem 13.3. An affine action of degree at least 1024 is covered by Theorem 14.2. At smaller degrees at least five, the soluble and nonsoluble cases are Theorems 14.3 and 14.5, with the exceptional degree-five Frobenius action covered by Proposition 14.4. At degree four the only primitive groups are the natural A_4, S_4 , the latter again covered by that proposition. Degrees three and two leave precisely the allowed natural actions and the binary group.

Now let U be imprimitive. Choose a minimal nontrivial block of size r , with s blocks and faithful top $T \leq S_s$. The full block stabilizer induces an exact primitive local group $L \leq S_r$:

any smaller block for that induced action and all its translates would give a smaller block for U . Transport coordinates along representatives of the block action to obtain the actual embedding $U \leq L \wr T$. This does not impose a splitting or full block kernel.

If L is nonaffine, use Theorem 13.3. If it is affine with $r \notin \{2, 3, 4, 5, 8, 9, 16\}$, use Theorem 11.2. For the four degrees 5, 8, 9, 16, Theorem 12.3 covers every count except its explicit finite lists; these lists are consumed, respectively, by Theorems 18.7, 16.3, 16.4 and 16.3. Each is an entire-normal statement on arbitrary complete sources.

For $r = 3$, the exact local group is C_3 or S_3 . Theorem 12.1 and Proposition 15.3 cover all C_3 counts. The same capacity theorem and Theorem 15.9 cover all S_3 counts.

For $r = 4$, the local group is natural A_4 or S_4 . Theorem 12.2 leaves only its displayed finite count sets, all at most 3072. In the actual affine frame put $E = U \cap (\mathbb{F}_2^2)^s$, $R = U/E$ and $D = R \cap C_3^s$. For $D \neq 1$, Theorem 16.5 applies. For $D = 1$, Theorem 12.4 covers all counts except 4, 8, 16, with counts two and three impossible. At four blocks use Theorem 18.9; at eight and sixteen use Theorem 19.4. This disposes of every exact four-point component.

Only $r = 2$, $L = C_2$ remains. Its pair kernel is binary, so its top is nonbinary whenever U is nonbinary. The prime-capacity theorem leaves exactly

$$\{2^a : 1 \leq a \leq 8\} \cup \{3 \cdot 2^a : 0 \leq a \leq 10\}.$$

At count two the whole action would be binary. Counts three, four and six are Proposition 18.8 and Propositions 18.6, 18.4. Count eight is Theorem 18.9, and counts twelve and sixteen are Theorems 17.7 and 17.15. The remaining necessary counts are

$$24, 32, 48, 64, 96, 128, 192, 256, 384, 768, 1536, 3072.$$

Every one is even and at least 24. For each original $N \triangleleft U$, Corollary 6.7 applies to the actual section $K/(K \cap N)$, where K is the pair kernel, with quotient $U/(KN)$. There is no assumption that $N \leq K$. Since these physical degrees form a fixed finite set, Theorem 6.5 covers the union of all their normal entries with a strict gap. This proves the required exhaustion.

Finally choose one of the proved certificates on each original action class, before inspecting its complementary group, and give the finitely many consumer families a fixed order. For a subgroup in $\mathcal{X}_n$, at least one orbit and its actual Goursat normal entry belong to the first eligible family. Apply that family's complete incidence sum. For a family with a hot/cold split, the existence of any hot eligible occurrence defines its hot part; all eligible occurrences in its remaining part are cold. Thus a chosen cold pointing cannot hide another hot one.

The incidence factor for deleting the original w -point orbit is $n!/(b!a(U))$, with $b = n - w$. It changes the occurrence factorial once. None of the block systems, chief factors, comparator degrees or character tuples acquires a physical support or normalizer. The full-fibre theorems are evaluated first on the same actual complete $J \leq S_b$, so the cold bounds multiply s_b/L_b and the hot bounds are complete scalars at n . Every finite family has strict positive margin, and every unbounded family has its proved uniform margin and subquadratic error. A finite sum of their exponential rows and quadratic scalars gives the asserted K, η, c, c' . No bound on s_b/L_b was assumed. $\square$

21 Binary sections and joint character incidence

The nonbinary hypothesis in Lemma 5.3 is essential. For binary tops it is replaced by a joint budget indexed by actual character kernels. Write $B_a = (\binom{a}{\lfloor a/2 \rfloor})$, including $B_0 = 1$.

Lemma 21.1 (Locally dependent maps over $\mathbb{F}_2$). *Let $\mathcal{M} \leq \text{Hom}(V, W)$ have dimension at least two, and suppose that $\dim \mathcal{M}(v) \leq 1$ for every $v \in V$. All maps in $\mathcal{M}$ have image in one common line of W .*

Proof. If $f \in \mathcal{M}$ has rank at least two, then for every $g \in \mathcal{M}$ one has $\ker f \leq \ker g$. To see this, take $u \in \ker f$ and two vectors with independent f -images. Applying pointwise dependence to each vector and its translate by u forces $g(u)$ to lie in both independent lines. Thus g induces a map on $\text{im } f$ taking each x to either 0 or x . Linearity on independent pairs forces this map to be zero or the identity. Hence $g = 0$ or f , contradicting $\dim \mathcal{M} \geq 2$. All nonzero maps therefore have rank one. Two maps with distinct image lines have the form $u\alpha$ and $v\beta$ for independent u, v . There is an input on which both nonzero functionals α, β are one, contradicting pointwise dependence. $\square$

Lemma 21.2 (A deterministic separator). *Let $\mathcal{L} \leq H \otimes S$ have dimension ℓ , and put*

$$j_0 = \max_{0 \neq \lambda \in H} \dim(\mathcal{L} \cap (\lambda \otimes S)),$$

with empty maximum zero. There are q independent functionals on S whose simultaneous evaluation is injective on $\mathcal{L}$, where

$$q \leq \left\lfloor \frac{\ell + \max(1, j_0)}{2} \right\rfloor. \quad (21.1)$$

For $\mathcal{L} = 0$ take $q = 0$.

Proof. View tensors as maps $S^* \rightarrow H$. On the current common kernel, select an evaluation of rank at least two whenever one exists. Each step removes at least two dimensions, and its functional is independent of those already selected. If a remainder of dimension $r \geq 2$ has only evaluations of rank at most one, Lemma 21.1 puts it in one character line, so $r \leq j_0$. Separate its support with r more functionals. A remainder of dimension one requires only one functional. If k first-stage steps occurred, then $\ell \geq 2k + r$ and $q = k + r$, proving (21.1) in all cases. The final functionals remain independent: earlier ones vanish on the remaining support. $\square$

Let $T \leq S_{2^a}$ be transitive and binary and A a section of its permutation module whose action factors through B . Set

$$S = A^B, \quad t = \dim S, \quad H = H^1(B, \mathbb{F}_2), \quad \mathcal{L} = \ker(H^1(B, S) \rightarrow H^1(B, A)), \quad \ell = \dim \mathcal{L}.$$

The soluble module bound gives $t, \ell \leq B_a$, by applying it to A^B and the actual section $(A/S)^B \cong \mathcal{L}$.

Lemma 21.3 (Joint character-kernel budget). *Inflate H into $\text{Hom}(T, C_2)$ and fix a point x . Let $H_0 = \{\chi \in H : \chi(T_x) = 0\}$. For $W \leq H$ put $j = \dim(W \cap H_0)$. Then*

$$t + \dim(\mathcal{L} \cap (W \otimes S)) \leq 2^j B_{a-j}. \quad (21.2)$$

Proof. Let $K_W = \bigcap_{\chi \in W} \ker \chi$ in T . Its number of orbits is $[T : K_W T_x] = 2^j$, since the annihilator of the image of T_x in T/K_W is $W \cap H_0$. The orbit lengths are 2^{a-j} . The preimage in A of the connecting-map slice $\mathcal{L} \cap (W \otimes S)$ has dimension equal to the left side of (21.2). The group K_W fixes it: the corresponding cocycles into the trivial module S vanish on K_W . Filter this actual trivial section by the K_W -orbit modules and apply the soluble bound B_{a-j} to each. This proves the inequality. $\square$

Theorem 21.4 (Binary capacity from degree 64). *For $a \geq 5$ there is $C \leq S$ with*

$$2 \dim C + 4 \dim(A/C)^B \leq 2B_a + 2B_{a-1}. \quad (21.3)$$

For $a \geq 6$ the right side is at most $15 \cdot 2^a/16$.

Proof. A nonzero character has one or two kernel orbits, so Lemma 21.3 gives $j_0 \leq J = 2B_{a-1} - t$, using $B_a \leq 2B_{a-1}$. Apply Lemma 21.2 and let C be the common kernel in S of the selected functionals. Then $\dim C = t - q$ and $\mathcal{L} \cap (H \otimes C) = 0$. Equation (5.4) gives $\dim(A/C)^B = q$, so the cost is $2t + 2q$. If $J \geq 1$, this is at most $2t + \ell + J = t + \ell + 2B_{a-1} \leq 2B_a + 2B_{a-1}$. If $J = 0$, then $t = 2B_{a-1} \leq B_a$. The binomial identities imply a even and $t = B_a = 2B_{a-1}$; this central binomial coefficient is even. Therefore $q \leq \lceil \ell/2 \rceil \leq B_a/2$, giving the same result. Finally, for $a \geq 6$, both central-binomial proportions $B_a/2^a$ and $B_{a-1}/2^{a-1}$ are at most $5/16$. $\square$

Theorem 21.5 (The degree-32 boundary). *If $a = 5$, there is $C \leq S$ with $2 \dim C + 4 \dim(A/C)^B \leq 30$.*

Proof. Here $t, \ell \leq 10$ and $j_0 \leq 12 - t$. The separator gives cost at most $t + \ell + 12 \leq 32$. Unless $t = \ell = 10$, it is at most 31 and hence at most 30, since the cost is even.

Assume $t = \ell = 10$. Put H_0 as in Lemma 21.3 and choose a complement $H = H_0 \oplus H_1$. For

$$(M_0, M_1, M_2, M_3, M_4, M_5) = (10, 12, 12, 16, 16, 32)$$

the joint budget is $\dim(\mathcal{L} \cap (W \otimes S)) \leq M_j - 10$. In particular $\mathcal{L} \cap (H_1 \otimes S) = 0$, so projection embeds $\mathcal{L}$ into a ten-dimensional space $\bar{\mathcal{L}} \leq H_0 \otimes S$. For $W_0 \leq H_0$,

$$\dim(\bar{\mathcal{L}} \cap (W_0 \otimes S)) \leq M_{\dim W_0} - 10.$$

We have $\dim H_0 \leq 5$ from the orbit count; taking $W_0 = H_0$ forces equality, because $M_j - 10 \leq 6$ for $j \leq 4$.

Let R_q be the number of nonzero tensors of rank q in $\bar{\mathcal{L}}$, $1 \leq q \leq 5$. Double count containment of their row support in r -spaces of H_0 . The instances $r = 2, 4$ give the simultaneous constraints

$$\begin{aligned} 15R_1 + R_2 &\leq 465, \\ 15R_1 + 7R_2 + 3R_3 + R_4 &\leq 1953, \\ R_1 + R_2 + R_3 + R_4 + R_5 &= 1023. \end{aligned} \quad (21.4)$$

Indeed the exact double count is $\sum_{\dim W=r} (2^{\dim(\bar{\mathcal{L}} \cap (W \otimes S))} - 1) = \sum_q R_q \binom{5-q}{r-q}_2$.

For a uniform five-space $C \leq S = \mathbb{F}_2^{10}$, the probability of containing a fixed q -dimensional image is

$$p_q = \frac{\binom{10-q}{5-q}_2}{\binom{10}{5}_2}, \quad (p_1, \dots, p_5) = \left(\frac{1}{33}, \frac{5}{5621}, \frac{1}{40953}, \frac{1}{1733677}, \frac{1}{109221651} \right).$$

Set $z = p_5$, $y = (p_3 - p_5)/3$, and $x = (p_1 - 15y - z)/15$. These are positive, and the coefficient vector of x times the first row of (21.4), y times the second, and z times the third dominates $(p_1, \dots, p_5)$ coordinatewise. The slacks in coordinates two and four are respectively $1932416/1638324765$ and $2480/327664953$; the other three are equalities. Consequently

$$\sum_q p_q R_q \leq 465x + 1953y + 1023z = \frac{103925485}{109221651} < 1. \quad (21.5)$$

Projection from H to H_0 never increases tensor rank, and is injective on $\mathcal{L}$. Since containment probability decreases with rank, the expected number of nonzero tensors of the original $\mathcal{L}$ contained in $H \otimes C$ is also less than one. This number is an integer, so some C has zero intersection. Equation (5.4) then gives $\dim(A/C)^B = 10 - 5 = 5$, of cost $10 + 20 = 30$. $\square$

Every simple binary module for a 2-group is trivial. Hence these costs are precisely $2 \dim C + 4r(A/C)$. A binary transitive pair action of width $2s \geq 64$ therefore has an entry certificate for every normal axis, with $g \geq s/16$, by Proposition 6.3. To sum over all widths one must also bound the number of actions, axes, pair systems, cuts, and constants D_C uniformly. The finite-menu theorem alone does not perform that unbounded sum.

22 Reversible simultaneous quotient transport

The construction below permits nonabelian common quotients and proper subdirect replacement groups. These are actual group maps, rather than identifications of abstract quotient types.

Proposition 22.1 (Simultaneous reconstruction). *Let $H \leq \prod_i U_i$ be full subdirect. For selected coordinates let $N_i = H \cap U_i$ be the exact embedded coordinate axis, and specify maps*

$$\alpha_i : U_i \twoheadrightarrow Q_i = U_i/N_i, \quad \beta_i : P_i \twoheadrightarrow Q_i, \quad P_i \leq \prod_{V \in W_i} V$$

with P_i full on the displayed word W_i . Use identity maps on the untouched coordinates. Set $\alpha = \prod_i \alpha_i$, $\beta = \prod_i \beta_i$, and

$$\overline{H} = \alpha(H), \quad H_{\text{new}} = \beta^{-1}(\overline{H}). \quad (22.1)$$

Then H_{new} is full on every replacement factor. The decorated map is injective, with inverse

$$H = \alpha^{-1}(\beta(H_{\text{new}})). \quad (22.2)$$

Proof. The embedded N_i commute and each is contained in H . Hence $\ker \alpha = \prod_i N_i \leq H$, proving $H = \alpha^{-1}(\overline{H})$. The projection of $\overline{H}$ onto each Q_i is full. Any element of P_i has an image appearing in some element of $\overline{H}$; surjectivity of the other β_j permits a simultaneous lift. Thus the projection of H_{new} onto the whole P_i is full, which implies fullness on its displayed factors. Surjectivity of β gives $\beta(H_{\text{new}}) = \overline{H}$, proving the inverse formula. No step replaces P_i by the full product of its projections. $\square$

Proposition 22.2 (Bounded decoration fibre). *Suppose the original actions, words, and maps range over fixed finite menus of bounded degree, each replacement has the same physical degree as its original coordinate, and original critical coordinates are untouched. If the original noncritical support has degree b , the number of decorated preimages of any target subgroup on n points is at most $(n+2)^{Kb}$ for a menu-dependent constant K .*

Proof. Realize each replacement on the same labelled points as its original orbit. To reconstruct the decorated block structure, select and group the target orbits into the changed bounded blocks, and record the finitely many local action types, coordinate bijections, normal axes, and map choices. There are $O(b)$ selected orbits, grouping slots, and finite labels. Allowing $n+2$ choices at each slot bounds all these possibilities by $(n+2)^{Kb}$. For fixed decorations the original subgroup is unique by (22.2). This counts labelled groups directly, so division by $n!$ retains the original action-normalizer weights and occurrence factorials. $\square$

The original denominator can also be retained directly when only an upper bound is needed; uniqueness of the orbit presentation is unnecessary.

Lemma 22.3 (Upper assembly with original normalizers). *Fix nonempty finite sets Ω_i , actions $U_i \leq S(\Omega_i)$, and multiplicities m_i , and let X be the disjoint union of these occurrences, of total size n . Put*

$$W = \prod_i (N_{S(\Omega_i)}(U_i) \wr S_{m_i}) \leq S(X).$$

If $\mathcal{P}$ is any family of subgroups of $S(X)$ invariant under conjugation by W , then its assembled family $\mathcal{A} = \{gKg^{-1} : g \in S(X), K \in \mathcal{P}\}$ satisfies

$$|\mathcal{A}| \leq \frac{n! |\mathcal{P}|}{\prod_i |N_{S(\Omega_i)}(U_i)|^{m_i m_i!}}.$$

Proof. For every pair (g, K) and every $w \in W$, the pair $(gw, w^{-1}Kw)$ has the same assembled subgroup. It remains in $S(X) \times \mathcal{P}$ by invariance, and distinct w give distinct first coordinates. Every nonempty assembly fibre therefore has at least $|W|$ elements. Summing the fibre sizes gives $|\mathcal{A}||W| \leq n! |\mathcal{P}|$. The displayed denominator is the order of the original internal-symmetry group W . $\square$

No transitivity, separation of the action types, or fullness of the local subgroups is required for this upper bound. Invariance is a condition on the whole family; individual subgroups need not be fixed.

Here is the quantitative form needed to use such a transport. Suppose a target binary word has critical rank R , a regular C_4 factors, and noncritical carrier support of half-degree $4T$, so $N = R + 2a + 4T$. Put $C_{\text{new}} = 2a + 4T$ and write the original noncritical support as $2C_{\text{old}}$. Assume each replacement retains at least a fixed proportion $\delta > 0$ of noncritical points. Then

$$\delta C_{\text{old}} \leq C_{\text{new}} \leq C_{\text{old}}. \quad (22.3)$$

Proposition 22.4 (Absorbing decorations). *Assume the complete normalized target mass at each profile satisfies the two uniform estimates*

$$\begin{aligned} T \leq a/140, \quad a \geq 1 : \quad & 2^{-aN/25 + O(a \log(N+2))}, \\ T > a/140 : \quad & 2^{-(29/656)TN + O(N \log(N+2))}. \end{aligned} \quad (22.4)$$

With the finite-menu assumptions of Proposition 22.2 and (22.3), the full original family with a noncritical orbit has normalized mass $2^{-\Omega(N)}$.

Proof. First remove the original range $0 < C_{\text{old}} \leq \sqrt{N}/4$ by Theorem 4.2, before charging any decorations. Indeed its noncritical degree is at most $\sqrt{N}/2$, and its critical degree is at least N for large N . The remaining range satisfies $C_{\text{new}} > \delta\sqrt{N}/4$. By Proposition 22.2, after summing over original supports the additional factor is $2^{O_\delta(C_{\text{new}} \log(N+2))}$.

If $T \leq a/140$, then $C_{\text{new}} < 3a$. This factor is absorbed by the first error in (22.4), and the negative term is at most $-N/25$ before the smaller error. If $T > a/140$, then $C_{\text{new}} < 284T$, so $T > \delta\sqrt{N}/1136$. The negative term in the second envelope has order at least $N^{3/2}$ and absorbs both the displayed error and the decoration cost. There are only polynomially many profiles and support sizes. Their sum retains the exponential gain. $\square$

The inequalities (22.4) are the stronger profile bounds proved for the full twelve-carrier mixture in Theorem 23.15. Multiplying only a final $2^{-\gamma N}$ mixture bound by the decoration factor would be insufficient. An all-critical replacement also fails (22.3), and is outside the negligible-mass conclusion even though the reconstruction map remains valid.

23 The complete critical, cyclic-four, and carrier mixture

All groups and all direct-product coordinates in this section are labelled. Write $\text{SD}(G_1, \dots, G_t)$ for the number of subgroups full on every displayed coordinate, and put

$$\begin{aligned}\mathcal{K}_8 &= \{8T18, 8T26, 8T27, 8T28, 8T29, 8T31, 8T35\}, \\ \mathcal{K}_{16} &= \{16T1082, 16T1083, 16T1084, 16T1332, 16T1547\}.\end{aligned}$$

The names specify the literal permutation actions in the finite data. A carrier of degree eight has weight one, and one of degree sixteen has weight two. A carrier word of weight T has degree $8T$. Let D be a critical word of elementary quotient rank R .

23.1 Three simultaneous parameters on a full projection

For a finite 2-group H set

$$d(H) = \dim H/\Phi(H), \quad \rho(H) = \max_{M \triangleleft H, M \leq H'} d_H(M), \quad d_H(M) = \log |M : M^2[M, H]|.$$

The invariant $\tau(H)$ is the one in (3.2); the five-term sequence identifies it with $d_H(\Phi(H))$.

Lemma 23.1 (Joint transition). *Let $K \leq L \times G$ be full on both factors and let $N = K \cap G$. Put*

$$k = d_G(N), \quad n = \log |N|, \quad m = \max_{M \triangleleft G, M \leq N \cap G'} d_G(M), \quad R_N = N^2[N, G], \quad a_2 = d_G(R_N).$$

Then $\delta = d(K) - d(L)$ and $\epsilon = \max(0, \rho(K) - \rho(L))$ satisfy

$$0 \leq \delta \leq k, \quad 0 \leq \epsilon \leq m, \quad \delta + \epsilon \leq n, \quad \tau(K) - \tau(L) \leq k - \delta + a_2. \quad (23.1)$$

Proof. The map on Frattini quotients is onto, and $\delta = \log |N : N \cap \Phi(K)|$. Fullness gives $[N, K] = [N, G]$, so $R_N \leq N \cap \Phi(K)$ and $\delta \leq k$. For $M \triangleleft K$ with $M \leq K'$, let M_0 be its projection to L and $M_1 = M \cap N$. Lifting normal generators gives

$$d_K(M) \leq d_L(M_0) + d_K(M_1).$$

Here $M_1 \triangleleft G$, $M_1 \leq N \cap G'$, and $d_K(M_1) = d_G(M_1) \leq m$. Also $M_1 \leq N \cap \Phi(K)$, so $d_K(M_1) \leq n - \delta$. Maximization proves the two bounds on ϵ .

The map $\Phi(K) \rightarrow \Phi(L)$ is onto. After quotienting by squares and ambient commutators, its kernel is generated by $N \cap \Phi(K)$. Indeed products of squares and commutators in $\Phi(L)$ lift to such products in $\Phi(K)$. Consequently

$$\tau(K) \leq \tau(L) + d_K(N \cap \Phi(K)).$$

The quotient $(N \cap \Phi(K))/R_N$ is trivial elementary of dimension $k - \delta$, and the image of R_N has dimension at most a_2 . This proves the last inequality. $\square$

For a row $r = (k, n, m, a_2, c, g)$ define

$$h_r(x, y) = \max\{x\delta + y\epsilon : 0 \leq \delta \leq k, 0 \leq \epsilon \leq m, \delta + \epsilon \leq n\}, \quad q_r = \max(m, a_2).$$

The quantity a_2 cannot in general be replaced by m . The following criterion makes that replacement available for an entire ambient group and reduces the radical certificates to generator tests.

Lemma 23.2 (Finite radical certificates). *Let $G = \langle s_1, \dots, s_u \rangle$ be a finite 2-group and $N = \langle x_1, \dots, x_v \rangle \triangleleft G$. Then*

$$R_N = \langle x_i^2, [x_i, s_j] : 1 \leq i \leq v, 1 \leq j \leq u \rangle^G.$$

If $\Phi(G) \leq G'$, then $R_N \leq N \cap G'$ and $a_2 \leq m$ for every such N . If also $G' \leq N$, then $m = \rho(G)$ and G/N is abelian.

Proof. Write U for the displayed normal closure. Its generators lie in R_N , and $R_N \triangleleft G$, so $U \leq R_N$. In G/U , each $x_i U$ has square one and commutes with every $s_j U$, hence with the whole quotient. The image of N is therefore central elementary abelian, which gives $N^2[N, G] \leq U$ and proves equality. An ordinary closure can replace this normal closure when its normality in the original G has been certified.

Every square and ambient commutator from N lies in $\Phi(G)$. Under the stated containment this puts the same G -normal subgroup R_N inside $N \cap G'$. Its relative head is one of the terms in the maximum defining m , so $a_2 \leq m$. Finally $G' \leq N$ gives $N \cap G' = G'$, proving the remaining assertions. $\square$

Lemma 23.3 (One centralizer controls all normal heads). *Let G be a finite group, p a prime, $N \triangleleft G$, and $u \in G$. Write $R = N^p[N, G]$ and $d_{G,p}(N) = \log_p |N : R|$. Then*

$$p^{d_{G,p}(N)} \leq |C_N(u)|.$$

Consequently, for a finite 2-group G and any subgroup $S \leq G$,

$$2^{\max_{M \triangleleft G, M \leq S} d_G(M)} \leq |C_S(u)|, \quad 2^{\rho(G)} \leq |C_{G'}(u)|.$$

Neither N nor G' is required to be abelian.

Proof. The set map $f : N \rightarrow R$ given by $f(x) = [x, u]$ need not be a homomorphism. If $f(x) = f(y)$, then $y^{-1}x \in C_N(u)$. Choose one representative y_r of each nonempty fibre. The map

$$x \mapsto (f(x), y_{f(x)}^{-1}x)$$

injects N into $R \times C_N(u)$. Thus $|N| \leq |R| |C_N(u)|$, proving the first bound after division by $|R|$. Apply it to every original G -normal subgroup $M \leq S$ and use $C_M(u) \leq C_S(u)$. Taking the maximum gives the second bound; putting $S = G'$ gives the last one. $\square$

Lemma 23.4 (Joint commutator radicals). *Let G be a finite 2-group with $\Phi(G) = D = G'$, and put*

$$V = G/D, \quad E = D/(D^2[D, G]).$$

For $\lambda \in E^$ let b_λ be the alternating form on V given by $b_\lambda(xD, yD) = \lambda([x, y])$. For every original normal subgroup $N \geq D$, write $W = N/D$ and*

$$J_W = \{\lambda \in E^* : W \leq \text{rad}(b_\lambda)\}.$$

Then

$$d_G(N) \leq \dim W + \dim J_W. \tag{23.2}$$

Let q, a be nonnegative integers. If the forms associated with every $q + 1$ independent characters in E^ have common radical of dimension at most a , then*

$$\dim W > a \implies d_G(N) \leq \dim W + q.$$

In particular, zero common radicals for independent character pairs give $d_G(N) \leq \dim W + 1$ for $W \neq 0$. If each nonzero form has radical dimension at most a , then $d_G(N) \leq \dim W$ whenever $\dim W > a$. The subgroup D need not be abelian.

Proof. Elements of E^* are exactly the G -invariant characters of D . The commutator identities and this invariance make $\lambda([x, y])$ additive in both original variables. The resulting maps vanish on G' in each variable, so they descend to the displayed alternating forms on V .

Restriction of a G -invariant character of N to D has kernel the characters of N/D , of dimension $\dim W$. Every character in the restriction image annihilates $[N, G]$, so that image lies in J_W . Rank-nullity proves (23.2). This uses only a necessary condition on the restriction image; no assertion that every member of J_W extends is needed.

If $\dim J_W > q$, choose $q + 1$ independent characters in J_W . The same W lies in all their radicals, contradicting $\dim W > a$. Thus $\dim J_W \leq q$. The two special cases follow by taking $(q, a) = (1, 0)$ and $q = 0$, respectively. $\square$

Corollary 23.5 (Star commutator family). *In the setting of Lemma 23.4, suppose that there are linear identifications $V = U \oplus \mathbb{F}_2$ and $E^* = U^*$ under which the actual commutator forms are*

$$b_\lambda((u, t), (v, s)) = \lambda(u)s + t\lambda(v).$$

Then every original normal subgroup $N \geq D$ satisfies

$$d_G(N) \leq \max\{\dim U, \dim(N/D)\}.$$

Proof. Put $W = N/D$. If W contains (u, t) with $t \neq 0$, then every $\lambda \in J_W$ satisfies $t\lambda(v) = 0$ for all $v \in U$, by testing against $(v, 0)$. Thus $J_W = 0$. Otherwise $W \leq U \oplus 0$, and first projection identifies W with a subspace $S \leq U$. Testing against $(0, 1)$ shows that $J_W = S^\perp \leq U^*$, so $\dim W + \dim J_W = \dim U$. Both cases give the result through (23.2). The required identifications concern the complete original character space and the actual quotient; ranks of selected numerical forms alone do not supply them. $\square$

Lemma 23.6 (Derived-intersection directions). *Keep G, D, V, E as in Lemma 23.4, but let $N \triangleleft G$ be arbitrary. Put*

$$B = N \cap D, \quad W = ND/D, \quad L_B = \{\lambda \in E^* : \lambda|_B = 0\}.$$

Then $L_B \leq J_W$ and $|N| = |B|2^{\dim W}$. If $B < D$, then $L_B \neq 0$. Consequently, a radical bound a for every nonzero b_λ gives $\dim W \leq a$. If independent pairs have zero common radical, then $\dim L_B \geq 2$ implies $N \leq D$. For the star family of Corollary 23.5, $L_B \neq 0$ gives the sharper bound

$$\dim W + \dim L_B \leq \dim U.$$

Proof. Original normality gives $[N, G] \leq N \cap D = B$, so every character in L_B annihilates all commutators from W . The second isomorphism theorem identifies W with N/B , proving the order identity. If $B < D$, choose a maximal proper G -normal subgroup J of D containing B . Then D/J is a minimal nontrivial normal subgroup of the finite 2-group G/J , and hence has order two and is central. Its nonzero character gives a member of L_B .

The single-form bound follows by choosing a nonzero member of L_B . Two independent members force $W = 0$ under the stated pair hypothesis. For the star family, a nonzero member of L_B forces every vector of W to have last coordinate zero. First projection identifies W with a subspace of the common annihilator of L_B in U , whose dimension is $\dim U - \dim L_B$. This proves the last inequality. These are constraints from characters of D vanishing on B ; the extendibility and power constraints on characters of B remain separate. $\square$

Lemma 23.7 (Exact annihilator and center detection). *Keep G, D, V, E as above and write $R = D^2[D, G]$. For every subgroup $B \leq D$, the common kernel in D of L_B is BR . If $R \leq B$, put*

$$C_B = \{x \in G : [x, G] \leq B\}, \quad U_B = \bigcap_{\lambda \in L_B} \text{rad}(b_\lambda).$$

Then $B \triangleleft G$, $D \leq C_B$, and the original quotient map induces

$$C_B/D \cong U_B, \quad |C_B| = |D|2^{\dim U_B}.$$

Suppose in addition that independent character pairs have zero common radical and that $N \triangleleft G$ satisfies neither $N \leq D$ nor $D \leq N$. For $B = N \cap D$, the space L_B has a unique nonzero member λ . If $R \leq B$, then, with $w = \dim(ND/D)$ and $r = \dim \text{rad}(b_\lambda)$,

$$|D : B| = 2, \quad 1 \leq w \leq r, \quad |Z(G/N)| = 2^{r+1-w}, \quad |(G/N)'| = 2. \quad (23.3)$$

Proof. The invariant characters of D are the complete dual of D/R . The ordinary double-annihilator identity in this vector space says that their common kernel on B is the inverse image of BR/R , which is BR . This does not identify B itself unless $R \leq B$.

Under that containment, B/R lies in the central elementary subgroup D/R of G/R , so B is normal in G . For every original $x, y \in G$, detection gives $[x, y] \in B$ exactly when $\lambda([x, y]) = 0$ for every $\lambda \in L_B$. Thus $x \in C_B$ exactly when $xD \in U_B$. In particular $D \leq C_B$, and the asserted equivalence and order formula follow from the original quotient map.

In the crossing case, Lemma 23.6 gives $L_B \neq 0$ and $W = ND/D \neq 0$. Two independent members of L_B would force $W = 0$, so $\dim L_B = 1$. Its unique nonzero member detects BR ; when $R \leq B$, this is its literal kernel B in D , of index two. Also $W \leq \text{rad}(b_\lambda)$, so $1 \leq w \leq r$. Since $|N| = |B|2^w$ and $Z(G/N) = C_B/N$, the center formula follows. Finally the original derived group maps onto $(G/N)'$ with kernel B , proving its order. $\square$

Lemma 23.8 (Saturation and exact crossing heads). *Keep the notation of Lemma 23.7. Suppose $\dim E \geq 2$, independent character pairs have zero common radical, and every original G -normal subgroup $M \leq D$ satisfies*

$$M \leq R \quad \text{or} \quad R \leq [M, G].$$

Then every crossing normal N satisfies, with $B = N \cap D$,

$$[N, G] = B = N^2[N, G], \quad d_G(N) = \dim(ND/D).$$

For every original normal $B \leq D$ with $B \not\leq R$,

$$B^2[B, G] = R, \quad \max_{M \triangleleft G, M \leq B} d_G(M) = \max \left\{ \max_{M \triangleleft G, M \leq R} d_G(M), d_G(B) \right\},$$

where $d_G(M) = \dim \text{Hom}(M, \mathbb{F}_2)^G$.

Proof. Put $H = [N, G] \leq B$. If $H \leq R$, every character of D annihilates H , so every form annihilates $W = ND/D$ against V . Two independent characters would then force $W = 0$, a contradiction. Thus saturation gives $R \leq H$. The full space L_H is the joint annihilator of W , and the pair hypothesis makes its dimension at most one. Since $0 \neq L_B \leq L_H$, these spaces are equal. Exact detection now gives $H = B$. Every original square in N lies in $N \cap D$, since G/D has exponent two, proving the relative radical identity and the head formula.

If $M \leq D$ and $M \not\leq R$, saturation and monotonicity of the relative radical give $R \leq [M, G] \leq M^2[M, G] \leq R$. For $M \leq B$ outside R , the resulting quotient inclusion $M/R \leq B/R$ gives $d_G(M) \leq d_G(B)$. Subgroups inside R contribute the other maximum. Both terms are attained within B , which proves equality without enumerating its normal subgroups. $\square$

For the original pair-family masters 16T1082, 16T1083, 16T1084 and 16T1547, the saturation hypothesis is certified by original commutator words on their 16 or 64 derived rows. Every word has positive commutator depth, so an element of a normal subgroup outside R generates all of R through its original ambient commutators. The same finite certificates give saturation for 16T1332, whose star family is treated separately from the pair hypothesis. For the four pair-family masters, every nonzero form has radical dimension two, $d_G(D) = 3$, and $\max_{M \triangleleft G, M \leq R} d_G(M) \leq 2$. Consequently every crossing normal has exactly one of the following rows, in the order (k, n, m, a_2, c, g) :

G	$w = 1$	$w = 2$
16T1082, 16T1083, 16T1084	(1, 4, 2, 2, 2, 1)	(2, 5, 2, 2, 1, 1)
16T1547	(1, 6, 2, 2, 2, 1)	(2, 7, 2, 2, 1, 1).

Indeed B has index two in D , its relative radical is R , and $d_G(B) = 2$; hence $m = a_2 = 2$. The remaining coordinates follow from Lemma 23.7 and $|N| = |B|2^w$. These are exhaustive possibilities for crossing normals; no realization of every listed row or identification of different normals is required.

For 16T1547, the remaining normals inside R also admit a uniform argument. Here $|R| = 8$, and the original commutator tests on its eight rows give a central subgroup Z of order two with $R^2[R, G] = [R, G] = Z$. Every original normal $M \leq R$ either lies in Z or contains Z ; if $M \not\leq Z$, the same saturation words give $M^2[M, G] = Z$. Thus M is 1, Z , an order-four subgroup above Z , or R . This argument requires no list or count of the order-four subgroups. Their exact rows are

M	(k, n, m, a_2, c, g)
1	(0, 0, 0, 0, 1, 6)
Z	(1, 1, 1, 0, 2, 5)
$Z < M < R$	(1, 2, 1, 1, 1, 4)
R	(2, 3, 2, 1, 3, 3).

For completeness, the original form separation first gives $C_M \leq D$. If $M < R$, saturation strengthens this to $C_M \leq R$. The identity $[R, G] = Z$ then gives $C_M = R$ for $Z \leq M < R$; the eight-row fixed tests give $C_1 = Z$, and $C_R = D$. These are the center orders in the table. The derived orders are $|D|/|M|$, and the maximum defining m ranges over every original normal below M , using the same relative-radical identities.

Lemma 23.9 (Crossing capacities for the full star family). *Keep G, D, V, E as above, and suppose the complete original form family is the star family on $V = U \oplus \mathbb{F}_2$, with $E^* = U^*$ and $d = \dim U$. Assume every original normal subgroup of D is comparable with R . For a crossing normal N , put $B = N \cap D$, $H = [N, G]$, $w = \dim(ND/D)$ and $g = \dim L_B$. Then*

$$R \leq H \leq B, \quad d_G(N) + g \leq d, \quad |(G/N)'| = 2^g, \quad |Z(G/N)| = 2^{d-w}.$$

Proof. The complete space L_H is exactly the joint annihilator of $W = ND/D$. Since it contains the nonzero space L_B , the star formula forces $W \leq U \oplus 0$. Its first projection identifies L_H with the annihilator of a w -dimensional subspace of U , so $\dim L_H = d - w$. If $H \leq R$, then $L_H = E^*$, forcing $w = 0$; comparability therefore gives $R \leq H$. Exact character detection and the order identity now give

$$|D| = 2^{d-w}|H| = 2^g|B|, \quad |N| = 2^w|B|.$$

All original powers remain in the relative radical $N^2[N, G]$, which contains H . Hence $2^{d_G(N)}|H| \leq |N|$, and the displayed identities yield $d_G(N) + g \leq d$.

The full common radical of $L_B \neq 0$ is its annihilator in $U \oplus 0$, of dimension $d - g$. Thus $|C_B| = |D|2^{d-g}$ and $|Z(G/N)| = |C_B|/|N| = 2^{d-w}$. The image of D in G/N has order $|D : B| = 2^g$. This proof uses neither a one-dimensional parameter space nor an equality between H and the full relative radical of N . $\square$

Lemma 23.10 (The remaining original normal branches). *For each of the five degree-sixteen masters put $D = G'$, $R = D^2[D, G]$, $d = \log |D|$, $s = \log |G/D|$ and $e = d_G(D)$. Their parameters are*

G	d	s	e	$\log R $
16T1082, 16T1083, 16T1084	4	6	3	1
16T1332	6	5	4	2
16T1547	6	6	3	3.

If $D \leq N$, with $w = \dim(N/D)$, then $0 \leq w \leq s$ and the six invariants are bounded coordinatewise by

$$(\max(e, w), d + w, e, e, s - w, 0).$$

Here the order, the maximum $m = e$, and both quotient slopes are exact. For $R < N < D$ the exhaustive exact alternatives are

G	(k, n, m, a_2, c, g)
16T1082, 16T1083, 16T1084	$(1, 2, 1, 1, 2, 2), (2, 3, 2, 1, 3, 1)$
16T1547	$(1, 4, 2, 2, 2, 2), (2, 5, 2, 2, 3, 1)$
16T1332	$(t, 2 + t, t, 1, 4, 4 - t), \quad t \in \{1, 2, 3\}.$

Inside R , the first three masters have the two exact rows $(0, 0, 0, 0, 1, 4)$ and $(1, 1, 1, 0, 3, 3)$. For 16T1332 there is a central subgroup Z of order two, and every original normal inside R is $1, Z$ or R , with respective rows

$$(0, 0, 0, 0, 1, 6), \quad (1, 1, 1, 0, 1, 5), \quad (1, 2, 1, 1, 4, 4).$$

Together with the four small-radical rows of 16T1547 and the crossing results above, these alternatives cover every original normal subgroup.

Proof. For $D \leq N$, the image dimension gives $|N| = 2^{d+w}$ and the abelian quotient has order 2^{s-w} . The original derived-normal maximum is e : its upper bound follows from the checked original centralizer ranks, and D itself attains the lower bound. The full-form head estimate gives $d_G(N) \leq \max(e, w)$, while $N^2[N, G] \leq N \cap D$ gives $a_2 \leq m = e$.

For $R < N < D$, nonempty original commutator saturation gives $N^2[N, G] = R$. Put $t = d_G(N)$ and $\ell = \dim L_N$. Then $t, \ell > 0$, $t + \ell = e$, $|N| = |R|2^t$, and $m = \max(m(R), t)$. In a pair family, $e = 3$; the full common radical of L_N has dimension zero when $\ell = 2$, and dimension two when $\ell = 1$. Lemma 23.7 gives $c = \ell + \dim U_N$ and $g = \ell$. In the complete star family the nonzero full parameter space L_N has common radical dimension $4 - \ell = t$, so $c = 4$ and $g = 4 - t$. The excluded case $\ell = 0$ is $N = D$ and belongs to the first branch.

For the first three masters, $|R| = 2$ makes R central, its relative radical trivial, and its head and normal-head maximum both one. Full pair-form separation and saturation give $C_1 = R$ and $C_R = D$, proving the two small rows. For 16T1332, the four original radical rows, with all five original generators acting, give $R^2[R, G] = [R, G] = Z$, with $|R| = 4$ and $|Z| = 2$. Every original normal $M \leq R$ lies in Z or contains Z ; outside Z it has relative radical Z and head at most one. Cardinality therefore forces $M = R$. The same tests and full star separation give $C_1 = Z$, $C_Z = R$, and $C_R = D$, yielding the three rows and $m(R) = d_G(R) = 1$.

Finally split an arbitrary $N \triangleleft G$ first by $D \leq N$, then by $N \leq D$. In the latter case saturation makes N comparable with R , giving either $N \leq R$ or $R < N < D$. The remaining case is crossing. This proof never counts normals by counting their possible rows and does not assume that every displayed possibility is realized. $\square$

There is a useful simplification when comparing these rows numerically. The feasible pairs satisfy $\delta \leq k$, $\epsilon \leq m$ and $\delta + \epsilon \leq n$, so their polygon depends on n only through $\min(n, k + m)$. Thus a comparison row (K, N_0, M, A, C, G_0) dominates all pair supports and both marks whenever

$$k \leq K, \quad m \leq M, \quad a_2 \leq A, \quad \min(n, k + m) \leq N_0, \quad c \leq C, \quad g \leq G_0.$$

This comparison preserves the original order n in the group-theoretic identities. It simply includes its entire feasible polygon in the comparison polygon. In particular it remains valid for a negative first mark, with a nonnegative terminal multiplier.

For $Q = G/N$ take $(c, g) = (\log |Q|, 0)$ if Q is abelian, and $(c, g) = (\log |Z(Q)|, \log |Q'|)$ otherwise. For a full subdirect product L of the bounded carrier factors,

$$|\text{Epi}(L, Q)| \leq C(d(L) + 2)^{C_{cd(L)+g\rho(L)}}. \quad (23.4)$$

The following elementary bound proves (23.4) for every finite 2-group source, without a restriction on its direct-product factors. Its constant is absolute on the fixed target alphabet.

Proposition 23.11 (Structured epimorphisms). *Let G, Q be finite 2-groups, and put $d = d(G)$, $\rho = \rho(G)$ and $q = \log |Q|$. Then*

$$|\text{Epi}(G, Q)| \leq (d + 1)^q |Q'|^\rho |Q|^{|Q'|^q} |Z(Q)|^d. \quad (23.5)$$

In particular, (23.4) holds with

$$C = C_Q = \max\{q, |Q|^{|Q'|^q}\} \geq 1, \quad c = \log |Z(Q)|, \quad g = \log |Q'|.$$

These choices also give the stated abelian case.

Proof. Choose an actual generating tuple of G of length d , by lifting a basis of $G/\Phi(G)$. For every epimorphism to Q , at most q of these same generators have images generating Q : choose a basis from their images in $Q/\Phi(Q)$ and lift back to the selected indices. Padding by identities gives at most $(d + 1)^q$ ordered tuples. Fix one such tuple $x = (x_1, \dots, x_q)$ and count the maps f for which $f(x)$ generates Q .

Set $D = [\langle x \rangle^G, G] \leq G'$ and $H = D \cap \ker f$. These are normal subgroups of the original G , and $f(D) = Q'$, so $[D : H] = |Q'|$. There are at most $|Q'|^\rho$ possible H . Indeed, for every $M \triangleleft G$ inside G' , the number of G -normal subgroups of index at most 2^a in M is at most $2^{a\rho}$. To see this, every proper normal interval in a finite 2-group admits a bottom step of index two: choose a central involution in its nontrivial normal quotient. Each normal $J \leq M$ has at most $2^{d_G(J)} - 1 \leq 2^\rho - 1$ normal index-two children, since each is the kernel of a nonzero G -invariant character of J . Keeping the previous subgroups as well as these children multiplies the cumulative count by at most 2^ρ at each step, starting with the single subgroup M .

Now fix H and let C be the inverse image in G of the centralizer of the tuple xH in G/H . The conjugates of that tuple lie coordinatewise in $x_i D/H$, so

$$[G : C] \leq |D/H|^q = |Q'|^q.$$

Every counted f sends C into $Z(Q)$, because it kills H and $f(x)$ generates Q . Thus $G \rightarrow Q/Z(Q)$ is constant on left C -cosets, giving at most $|Q|^{|Q'|^q}$ quotient maps. Normality of C is not required. In a fixed quotient-map fibre, choose one map f_0 when the fibre is nonempty. The pointwise differences $f(g)f_0(g)^{-1}$ are central and form a homomorphism $G \rightarrow Z(Q)$; they determine f . There are at most $|Z(Q)|^d$ such homomorphisms, by the original generating tuple. Empty fibres contribute zero. Multiplying the tuple, normal-subgroup, quotient-map and central-fibre bounds proves (23.5). Finally $(d+1)^q \leq (d+2)^{C_Q}$ and $|Q|^{|Q'|^q} \leq C_Q$. $\square$

23.2 The finite rows and the recurrence

Use the masters $X = 8T26$, $J = 8T27$, $P = 8T35$ and the five degree-sixteen carriers. The finite normal-kernel certificate supplies their actual rows and their multiplicities. It includes every literal normal kernel, by the central-involution closure argument in Lemma 24.5 below. The numerical reduction stated and certified in Appendix B gives the matrix

$$B = \frac{1}{64} \begin{pmatrix} 5 & 6 & 34 & 28 & 32 & 30 \\ 6 & 0 & 44 & 36 & 40 & 36 \\ 34 & 44 & 0 & 16 & 20 & 40 \\ 28 & 36 & 16 & 12 & 16 & 32 \\ 32 & 40 & 20 & 16 & 16 & 32 \\ 30 & 36 & 40 & 32 & 32 & 24 \end{pmatrix}. \quad (23.6)$$

$$\alpha = (5/8, 3/4, 0, 1/4, 1/4, 1/2), \quad \beta = (3/8, 1/2, 0, 1/4, 1/4, 1/2).$$

Here rows have physical scale $s_r = 1$ or 2. Before domination their pair coefficient is

$$\frac{\max\{h_r(c_s, g_s), h_s(c_r, g_r)\}}{s_r s_s},$$

and their marked coefficients are $k_r/s_r, q_r/s_r$. Each dominated row is moved to a row dominating all these coefficients simultaneously. Finally scale its accumulated mass by four, obtaining $p_i \geq 0$ with $\sum_i p_i = 4T$; this accounts for the factor $1/64$ in (23.6).

The crossing branch of $16T1332$ can be assigned directly to the sixth colour, without distinguishing its six possible pairs (g, w) . Indeed Lemma 23.9 and saturation give the coordinatewise upper row

$$(4 - g, 6 - g + w, \max(2, 4 - g), \max(2, 4 - g), 4 - w, g), \quad g, w \geq 1, \quad g + w \leq 4.$$

All these rows lie below $r_* = (3, 8, 3, 3, 3, 3)$ at scale two. For any displayed row $s = (k, n, m, a_2, c, g)$, with nonnegative slopes,

$$\max\{h_s(3, 3), h_{r_*}(c, g)\} \leq \max\{3 \min(n, k + m), 3(c + g)\}.$$

Checking this last expression on the forty-one displayed rows of Appendix B, with their stated colours j and scales s_s , gives the bound $8(4s_s)B_{6j}$ in every column. The self support is at most $18 \leq 64B_{66} = 24$, and both marks are $3 \leq 8\alpha_6 = 8\beta_6 = 4$. This handles all crossing normals of that master at once, while retaining their separate original weights. The remaining branches are covered by Lemma 23.10. Its effective-polygon comparison sends every pair-family row, every above-derived row, and the small-radical rows into the twelve displayed H16 rows. The first two star intermediate

rows go to H16 rows 2 and 4. The last star intermediate row is $r_{\dagger} = (3, 5, 3, 1, 4, 1)$, at scale two, and can also be assigned directly to colour six. Indeed

$$\max\{h_s(4, 1), h_{r_{\dagger}}(c, g)\} \leq \max\{4k + m, 3(c + g)\}.$$

The forty-one displayed comparisons satisfy the same bound $8(4s_s)B_{6j}$. Its self support is at most $15 \leq 24$ and its pair support with r_* is at most $18 \leq 24$; both marks are at most 4. Thus these two extra envelopes use the existing sixth colour and do not change the cone matrix or its physical normalization. No step replaces the original weighted normal family by a sum over distinct numerical labels.

Lemma 23.12 (The complete marked recurrence). *For every such master word, the logarithm of its full-subdirect count with D is at most the maximum, over $0 \leq j \leq R$, $0 \leq \ell \leq R/2$ and its row masses, of*

$$\mathcal{E} = \ell(R/2 - \ell) + (j - \ell)(R - j) + \frac{1}{2}p^T Bp + \sum_i p_i \{\ell(\alpha_i + \beta_i) + \alpha_i \max(j - 2\ell, 0)\}, \quad (23.7)$$

plus $O((R + T) \log(R + T + 2))$.

Proof. Project first onto the complete carrier subgroup H . For this same H , Proposition 3.4 gives the sum of the marks $2^{(j-\ell)d(H)+\ell\tau(H)}$, with coefficient

$$\phi^{-2} \begin{bmatrix} c \\ \ell \end{bmatrix}_2 72^\ell 2^{(j-\ell)(R-j)}, \quad c \leq R/2.$$

For one carrier peel the lower projection L , literal axis N and literal epimorphism $L \rightarrow G/N$ recover the subgroup uniquely. Thus (23.4) is used once, with no automorphism divisor. Lemma 23.1 shows that an arbitrary current mark $2^{xd+y\rho+z\tau}$, for $y, z \geq 0$, incurs at most

$$z(k + q_r) + h_r(x - z, y).$$

This remains true for negative x . For the initial mark $(x, y, z) = (j - \ell, 0, \ell)$ it is at most $\ell(k + q_r) + k \max(j - 2\ell, 0)$. Every later epimorphism contributes the nonnegative tilt (c_s, g_s) . Subadditivity of the support function bounds the additional cost of an unordered pair of peels by the maximum in the definition of the pair coefficient. Summing pairs gives at most the quadratic term $\frac{1}{2}p^T Bp$; adding diagonal self-pairs only enlarges it. The simultaneous row domination gives the last term of (23.7). Finally use $\begin{bmatrix} c \\ \ell \end{bmatrix}_2 \leq \phi^{-1} 2^{\ell(c-\ell)}$.

There are a fixed number of normal choices per coordinate, hence at most $2^{O(T)}$ histories. The polynomial factors in (23.4) cost $O(T \log(T + 2))$; 72^ℓ , the Gaussian constants and the outer (j, ℓ) sum cost $O(R + \log(R + 2))$. This proves the stated uniform error. The initial projection H is recovered intrinsically, so the argument attaches its critical factors exactly once. $\square$

Theorem 23.13 (Critical-carrier reserve). *Uniformly over all original carrier colour words of weight $T \geq 1$,*

$$\log \text{SD}(D, \mathbf{G}) \leq \frac{(R + 4T)^2}{4} - \frac{25}{82}RT - \frac{29}{164}T^2 + O((R + T) \log(R + T + 2)). \quad (23.8)$$

The sum of their original action weights is

$$A_T = [z^T] \exp \left(\frac{3}{64}z + \frac{65}{688128}z^2 \right). \quad (23.9)$$

Proof. Appendix B proves, for every point of the two exhaustive cones $j \geq 2\ell$ and $j \leq 2\ell$,

$$\mathcal{E} \leq R^2/4 + \frac{139}{328}R \sum_i p_i + \frac{627}{2624} \left(\sum_i p_i \right)^2.$$

With $\sum_i p_i = 4T$, this is (23.8) for masters. There are fixed epimorphisms

$$X \rightarrow 8T26, \quad J \rightarrow 8T27, 8T28, \quad P \rightarrow 8T18, 8T29, 8T31, 8T35.$$

The inverse image under their product, acting identically on D and all degree-sixteen factors, injects each original full-subdirect family into its master family. Its image recovers the original subgroup, and fullness of each replaced coordinate follows from surjectivity. This proves the bound for each original word.

The degree-eight normalizers have orders 192 for 8T18, 384 for 8T31, and 128 for the other five; their reciprocals sum to $3/64$. The degree-sixteen orders, in the displayed order, are 344064, 49152, 49152, 24576, 98304, whose reciprocals sum to $65/688128$. Summing $\prod_U 1/(a(U)^{m_U} m_U!)$ proves (23.9). Each injection is used profile by profile with its original weight. Overlap between the images of different profiles causes no identification of the original colours. $\square$

23.3 Adding every cyclic-four orbit

Let a be the number of regular C_4 orbits, let T be the carrier weight and put $N = R + 2a + 4T$. The exact original profile weight is

$$\frac{c_R}{8^a a!} A_T, \quad c_R = [y^R] \exp(y/2 + y^2/6 + y^4/384). \quad (23.10)$$

Lemma 23.14 (Cyclic-four comparison and Hall bound). *For any finite 2-group B of order 2^u ,*

$$\text{SD}(D, C_4^a, B) \leq \sum_{k=0}^{2a} \binom{2a}{k} \text{SD}(D, C_2^k, B), \quad (23.11)$$

$$\log |\text{Sub}(C_2^R \times C_4^a \times B)| \leq \frac{u^2 + (R + a + u)^2 + a^2}{4} + O((a + u) \log(a + u + 2) + \log(R + 2)). \quad (23.12)$$

In the first inequality B may be a displayed word, with fullness required on each of its factors.

Proof. For a central extension by C_2 , the subgroups above a fixed subgroup of the quotient consist of its unique full preimage and its complements to C_2 . The latter are absent or form a torsor of size $|\text{Hom}(H, C_2)|$. The split extension realizes the upper bound exactly. Apply this successively to the central involutions in the C_4 coordinates, retaining the projection to $D \times B$ throughout. After dropping fullness only on these coordinates, the comparison group is C_2^{2a} . Its uniquely determined set of nonzero coordinate projections has size k , giving (23.11). For the terminal upper bound, this support decomposition is unnecessary: the terminal family already permits every image in its binary abelian coordinates. Regroup the new $2a$ binary coordinates with the original binary abelian part, retaining fullness on each nonabelian critical coordinate and each carrier factor. The resulting terminal family has critical rank $R + 2a$, with no binomial prefactor. This regrouping concerns the comparison family; the original orbit profile and its weight remain unchanged.

The same central comparison gives, for every finite group E ,

$$|\text{Sub}(E \times B)| \leq |\text{Sub}(E \times C_2^u)|.$$

Indeed, choose a central subgroup $C \leq B$ of order two and compare $E \times B \rightarrow E \times (B/C)$ with its split extension. Apply induction to B/C , keeping $C_2 \times E$ as the unchanged exterior, and regroup the binary factors. This induction preserves every condition on the exact E -image; it does not require E to be abelian.

Set $E = C_4^a \times C_2^R$ and $r = R + u$. Reduce the C_4 coordinates of $C_4^a \times C_2^r$ modulo two. Its kernel has binary dimension $r + a$. Above an actual image U of dimension j , the squares span exactly the copy of U in the doubled C_4 coordinates. The square obstruction therefore has dimension j . The exact central-lift formula, with k the retained annihilator dimension, gives

$$|\text{Sub}(C_4^a \times C_2^r)| = \sum_{j=0}^a \begin{bmatrix} a \\ j \end{bmatrix}_2 \sum_{k=0}^{r+a-j} \begin{bmatrix} r+a-j \\ k \end{bmatrix}_2 2^{jk}.$$

Use $\begin{bmatrix} n \\ k \end{bmatrix}_2 \leq \phi^{-1} 2^{k(n-k)}$, where $\phi = \prod_{i \geq 1} (1 - 2^{-i})$. The exponent of each summand is at most

$$j(a-j) + k(r+a-j-k) + jk = j(a-j) + k(r+a-k) \leq \frac{a^2 + (r+a)^2}{4}.$$

There are at most $(a+1)(r+a+1)$ summands. In particular we have the stronger explicit bound

$$|\text{Sub}(C_2^R \times C_4^a \times B)| \leq \frac{(a+1)(R+a+u+1)}{\phi^2} 2^{((R+a+u)^2 + a^2)/4}. \quad (23.13)$$

The logarithm of the prefactor is $O((a+u) \log(a+u+2) + \log(R+2))$, uniformly also when $a = u = 0$. Adding the nonnegative $u^2/4$ recovers (23.12) with the stated constants. $\square$

Theorem 23.15 (Complete mixture). *The normalized mass, relative to L_{2N} , of all fixed-point-free binary subgroups whose orbit actions belong to the critical actions, regular C_4 , and the twelve carriers, with some noncritical orbit, is $2^{-\Omega(N)}$. At a fixed profile the following stronger bounds hold uniformly:*

$$\begin{aligned} T \leq a/140, a \geq 1 : & \quad 2^{-aN/25 + O(a \log(N+2))}, \\ T > a/140 : & \quad 2^{-(29/656)TN + O(N \log(N+2))}. \end{aligned} \quad (23.14)$$

Proof. The carrier product B has $u = \log |B| \leq 7T$. Use the exterior-preserving central comparison in Lemma 23.14, and regroup its unrestricted binary abelian coordinates as above. The terminal reserve from Theorem 23.13, applied directly at critical rank $R + 2a$, gives

$$\begin{aligned} \log \text{SD}(D, C_4^a, B) & \leq N^2/4 - \frac{25}{82}(R+2a)T - \frac{29}{164}T^2 + O(N \log(N+2)) \\ & \leq N^2/4 - \frac{29}{656}TN + O(N \log(N+2)). \end{aligned}$$

The original noncritical profile weights have a uniform finite-sum bound without evaluating their normalizers. If w_i is the original normalizer order of colour i , then $w_i \geq 1$, and any finite set $\mathcal{M}$ of multiplicity profiles satisfies

$$\sum_{\mathbf{m} \in \mathcal{M}} \prod_{i=1}^{13} \frac{1}{w_i^{m_i} m_i!} \leq \prod_{i=1}^{13} \exp(1/w_i) \leq e^{13}.$$

All thirteen colours remain separate, including those with the same master. Together with $c_R/c_N \leq (2N)^{2a+4T}$, division by $c_N G_N$ and insertion of (23.10) proves the second envelope.

For the other envelope first project onto an arbitrary actual $H \leq C_4^a \times B$. Its sectional binary rank is at most $a + u$: intersect an elementary section with $H \cap C_4^a$ and project the quotient to B . Hence $\tau(H) \leq a + u$. Proposition 3.4, with the nonnegative δ and $d/2$ dropped, gives

$$N_D(H) \leq X_R(d(H)) 2^{h_R(a+u+\log 72)+O(\log(R+2))}, \quad h_R(x) = \max_{0 \leq \ell \leq R/2} (x\ell - 3\ell^2/4) \leq x^2/3.$$

The graph identity $\sum_{H \leq C_4^a \times B} X_R(d(H)) = |\text{Sub}(C_2^R \times C_4^a \times B)|$ then allows a single use of (23.12). Put $N_0 = R + 2a$. Subtracting the resulting quadratic exponent from $N_0^2/4$ leaves exactly

$$aR/2 + a^2/6 - uR/2 - 7au/6 - 5u^2/6.$$

If $T \leq a/140$, then $u \leq a/20$, so this is at least $(19/40)aR + (17/160)a^2 \geq aN_0/20 \geq aN/25$; here $N \leq 71N_0/70$. The fixed $\log 72$ correction costs $O(a + 1)$, and all profile and coefficient costs are $O(a \log(N + 2))$. This proves the first envelope, including $T = 0$.

Let $C = 2a + 4T$ be the noncritical half-support. For this finite alphabet, the entire range $0 < C \leq \sqrt{N}/4$ has a direct estimate. Each original carrier action has binary character rank at most three times its physical scale: apply the certified head bound to the top normal subgroup and inflate characters along its fixed master epimorphism. This is strictly less than half its original degree. The regular C_4 has rank one. For an actual full noncritical tail J , restrict to one of its original blocks. The restriction kernel acts faithfully on the original complement, so the half-degree bound for that kernel preserves the local gap and gives $d_2(J) \leq C - 1$, even when J is a proper subdirect product. Applied to the actual evaluation kernel, the same bound gives $\tau(J) \leq C$. These marks concern J itself, before any master pullback.

To count such tails, the master pullback is injective, and its source has order 2^q with $q = 2a + u \leq 7C/4$. The central comparison above bounds its subgroup count by G_q . The terminal Hall estimate therefore bounds the model count by

$$2\phi^{-3}(R+1)(c+1)X_R(C-1)G_q2^{(C+7)^2/3},$$

where c is the number of nonabelian critical factors. The elementary Gaussian estimates give $X_R(C-1)/G_N \leq \phi^{-2}(R+1)2^{-N/2+1/2}$ and $G_q \leq \phi^{-1}(q+1)2^{q^2/4}$. Sum the critical profiles at their exact rank $R = N - C$, retaining c_R , and the noncritical profiles with their original weights as above. The logarithm of the normalized mass at this support is at most

$$-\frac{N}{2} + \frac{1}{2} + \frac{211}{192}C^2 + \frac{14}{3}C + \frac{49}{3} + C \log(2N) + O(\log(N+2)).$$

For $C \leq \sqrt{N}/4$ the quadratic terms leave at most $-1325N/3072$, and the remaining error is uniformly $o(N)$. Thus this range, including its support-size sum, is exponentially small. This finite-alphabet argument uses neither the general equality classification nor the coarse ordinary subgroup-count estimate from Theorem 4.2.

If $C > \sqrt{N}/4$ and $T > a/140$, then $C < 284T$, so $T > \sqrt{N}/1136$. The negative term of the second envelope therefore has order at least $N^{3/2}$ and absorbs its error. The first envelope is exponentially small uniformly for $a \geq 1$. After summing the original multiplicity weights, at most $(N+1)^2$ parameter pairs (a, T) remain. Their sum proves the theorem. $\square$

The same weighted estimates give a direct extension to one natural S_3 orbit, with critical orbits and the original thirteen noncritical colours on the remaining points. Write $\mathbf{p}$ for the critical profile of rank R , keep the noncritical profile $\mathbf{q}$ fixed, and put $C = 2a + 4T > 0$ and $N = R + C + 1$. The physical degree is $2N + 1$. The exterior group is a 2-group. If (g, d) lifts a three-cycle and $d^{2^k} = 1$, its 4^k th power is $(g, 1)$. Thus every subgroup full on the S_3 coordinate contains A_3 on that coordinate alone. The sign quotient, followed by its inverse full preimage, identifies its full models bijectively with those of $(\mathbf{p} + \mathbf{e}_{C_2}, \mathbf{q})$. Every original critical and noncritical coordinate remains full; no full projection onto either entire class product is required.

The physical weight is still the original one. If $\omega(\mathbf{p}, \mathbf{q})$ denotes the reciprocal product of the original normalizers and occurrence factorials, then

$$\frac{\omega(\mathbf{p}, \mathbf{q})}{6} = \frac{p_{C_2} + 1}{3} \omega(\mathbf{p} + \mathbf{e}_{C_2}, \mathbf{q}).$$

The factor includes both the marker normalizer six and the new C_2 occurrence factorial. The map $\mathbf{p} \mapsto \mathbf{p} + \mathbf{e}_{C_2}$ is injective into the critical profiles of rank $R + 1$, so positivity bounds the original weighted sum by $(R + 1)/3$ times the complete shifted sum, with $\mathbf{q}$ unchanged. Its half-degree is exactly $(R + 1) + C = N$. Moreover, $L_{2N+1} \geq (2N + 1)!c_N G_N$, retaining the nonnegative additional term of the odd coefficient. The preceding weighted estimates therefore give a polynomial times $2^{-\Omega(N)}$ for each such bin. There are at most $(N + 1)^2$ pairs (a, T) with $C + 1 \leq N$, and the weight correction is at most $(N + 1)/3$. Their cubic polynomial cost is absorbed by the exponential decay. This bounds the actual union of physical subgroups, allowing overlapping profile presentations. It is a specialization to this finite alphabet and leaves the general complete-family marker transfer of Section 28 unchanged.

Combining (23.14) with Propositions 22.1–22.4 proves the same negligible-mass conclusion for every fixed finite menu of simultaneous equal-degree quotient replacements retaining a positive fraction of the original noncritical support. The original small-support range is removed before paying its decoration fibre.

24 Complete binary counting

Let F_N be the number of all fixed-point-free 2-subgroups of S_{2N} divided by $L_{2N} = (2N)!c_N G_N$, and let E_N be the same normalized count restricted to groups with a noncritical orbit. Thus $F_N = C_{2N}^{\text{crit}} + E_N$ exactly. We prove the complete recurrence needed in Section 32.

24.1 The original degree-thirty-two boundary

Theorem 24.1 (Adaptive cover at degree sixteen). *Let $T \leq S_{16}$ be transitive and binary, and let A be a section of its original permutation module whose action factors through a specified quotient B of T . There are $C \leq A^B$ and a specified permutation cover of B of degree $v_0 \in \{8, 16\}$ such that*

$$v_0 + 2 \dim C + 4 \dim(A/C)^B \leq 30. \quad (24.1)$$

Proof. Use $S, t, H, \mathcal{L}, \ell, H_0$ from Lemma 21.3. Put $d_0 = \dim H_0 \leq 4$. For $a = 4$ the joint budgets $2^j B_{4-j}$, for $0 \leq j \leq 4$, are $(6, 6, 8, 8, 16)$. In particular $t, \ell \leq 6$. If $d_0 \leq 3$, taking $W = H$ gives $t + \ell \leq 8$; taking a character line gives $j_0 \leq 6 - t$. The separator in Lemma 21.2 supplies q evaluations with kernel $C \leq S$, and (5.4) gives $\dim C = t - q$, $\dim(A/C)^B = q$. If $t \leq 5$, their cost is $2t + 2q \leq t + \ell + 6 \leq 14$. If $t = 6$, then $\ell \leq 2$ and $j_0 = 0$, so $q \leq 1$ and the same bound holds. The original degree-sixteen cover $T \rightarrow B$ therefore proves (24.1).

If $d_0 = 4$, the common kernel of H_0 has sixteen orbits on sixteen points. Faithfulness makes it trivial. Thus $T = C_2^4$ is regular; as every character in H_0 comes from B , also $B = T$. Four independent characters give a faithful action of this same B on four disjoint pairs. Use it as the degree-eight cover. The original module still has $t, \ell \leq 6$, $j_0 \leq 6 - t$. For $t \leq 5$ the separator cost is at most $t + \ell + 6 \leq 17$; for $t = 6$ it has $q \leq \lceil \ell/2 \rceil \leq 3$ and cost at most 18. Adding the new cover degree gives at most 26. The module A was never replaced by a section of the eight-point permutation module. $\square$

For every transitive binary action U of degree 32 and every $N \triangleleft U$, choose an invariant pair system with kernel K and top $T \leq S_{16}$. The groups $Q = U/N$, $A = KN/N$ and $B = U/(KN)$ meet the theorem's hypotheses. Since every simple binary module of a 2-group is trivial, $r(A/C) = \dim(A/C)^B$. Proposition 6.3 gives the same-source envelope with $v = v_0 + 2 \dim C$, $\lambda = r(A/C)/2$ and $g = 32 - v - 8\lambda \geq 2$. The finite-menu proof of Theorem 6.5, restricted on its physical side to fixed-point-free binary sources, consequently gives

$$T_{32}(N) \leq \eta_{32}(N) + K_{32}(N, N - 16)F_{N-16}, \quad \eta_{32}(N) = 2^{-\Omega(N^2)}, \quad K_{32}(N, N - 16) = 2^{-\Omega(N)}. \quad (24.2)$$

Here T_{32} denotes the family with a degree-32 orbit and none larger. The auxiliary graph may have fixed points, so its moment is bounded by the complete count s_{b+qv} as in Proposition 6.3. Only the physical cold complement is fixed-point-free. At this fixed width all action, normal, frame and cut choices form a finite menu, and may all be retained at a fixed constant cost. The coefficient is the sum of $\Delta(2N, 2N - 32)D_C 2^{(4-g/16)(2N-32)}/a(U)$; its logarithm is at most $-g(2N - 32)/16 + O(\log(N + 2))$.

24.2 Uniformity over unbounded binary widths

Put $A_k = \sum_{j < k} B_j$, with $A_0 = 0$. These cumulative Boolean widths give the binary normal-count input directly.

Lemma 24.2 (Original normal heads). *For a transitive 2-group $U \leq S_{2^k}$ and every original normal subgroup $N \triangleleft U$,*

$$\dim \text{Hom}(N, C_2)^U \leq A_k.$$

Consequently $d(U) \leq A_k$, and the number of original normal subgroups of U is at most $2^{(2^k-1)A_k}$. Moreover $A_k = O(2^k/\sqrt{k+1})$, in particular $A_k = o(2^k)$.

Proof. At degree one the group is trivial. Otherwise choose an original invariant pair system, with kernel K and transitive top T of degree 2^{k-1} . The exact subgroup $N \cap K$ is a submodule of the pair permutation module, so its invariant character dimension is at most B_{k-1} by the binary module bound. Restriction of invariant characters from N has kernel the characters of the actual image NK/K , invariant under T . Thus the required head is at most $B_{k-1} + A_{k-1} = A_k$. Surjectivity of restriction is neither needed nor asserted.

Apply this to the original top subgroup to bound the ordinary character rank of U ; Frattini lifting gives A_k generators, padding by identities if necessary. To count normals, let c_a count the original U -normal subgroups of index at most 2^a . Every new subgroup at the next cutoff is an index-two child of one at the previous cutoff: for an original normal H , use an order-two subgroup of $Z(U/H)$. Each parent has at most $2^{A_k} - 1$ such children, from its nonzero invariant characters. Hence $c_0 = 1$ and $c_{a+1} \leq 2^{A_k} c_a$. The bound follows from $|U| \leq 2^{2^k-1}$. Finally the elementary central-binomial recurrence gives $(\lfloor j/2 \rfloor + 1)B_j^2 \leq 4^j$. Splitting the cumulative sum at $k/2$ proves the stated estimate for A_k . $\square$

Lemma 24.3 (Zero cuts at large binary widths). *Let U be transitive and binary of degree $w = 2s$, where $s = 2^j$, and fix an original pair kernel K and $N \triangleleft U$. For $A = KN/N$ and $B = U/(KN)$, the literal cut $C = 0$ has*

$$v = s, \quad r(A) = \dim A^B \leq B_j, \quad g = w - v - 4r(A) \geq w/128 > 0 \quad (j \geq 9).$$

For $j \geq 11$ the stronger bound $g \geq w/32$ holds. The corresponding coefficient remains $D = |A||H^1(B, A)|$, and the same-source moment weight is exactly $\Phi(J) = e_B(J)$.

Proof. The invariant bound follows by taking the full preimage of A^B in the original pair kernel and applying the binary module bound to that trivial quotient. Since B is binary, its simple modules over $\mathbb{F}_2$ are trivial, so this invariant dimension is the Schur capacity. The original top $T \leq S_s$ maps onto this same B ; Proposition 6.3 applies to that faithful cover, without assuming a faithful degree- s action of B or a split extension. With $C = 0$ it gives the stated D and Φ .

Pascal's identity gives $B_{j+1} \leq 2B_j$. The exact values $B_9 = 126$ and $B_{11} = 462$ therefore imply, respectively, $256B_j \leq 63 \cdot 2^j$ for $j \geq 9$ and $64B_j \leq 15 \cdot 2^j$ for $j \geq 11$. Substitution into $g = s - 4r(A)$ proves both bounds. $\square$

Thus physical widths 1024 and 2048 already have zero-cut parameters $e = g/16 \geq 1/2$ and $e \geq 1$, respectively, and can be included in a fixed finite direct-fusion prefix. At every width at least 4096, the zero cut meets the uniform margin used below. Only widths 64, 128, 256, 512 in the large-width route require the sharper cut construction; degree 32 remains the separate adaptive boundary above.

Theorem 24.4 (Complete large-orbit deletion). *Let $T_{64}(N)$ be the normalized family with some actual orbit of width at least 64. There are nonnegative η_{64} and K_{64} , with $K_{64}(N, M) = 0$ unless $M \leq N - 32$, such that eventually*

$$T_{64}(N) \leq \eta_{64}(N) + \sum_{M < N} K_{64}(N, M) F_M, \quad \eta_{64}(N) \leq 2^{-(2N)^2/2^{22}}, \quad \sum_M K_{64}(N, M) \leq 2^{-(2N)/512}. \quad (24.3)$$

Proof. Every nontrivial transitive 2-group has an invariant pair system: a central involution is fixed-point-free, since a central element fixing a point fixes its entire transitive orbit. An invariant pair system is determined by the partner of one point, so there are at most $w - 1$ systems. Retain all of them. Choose one auxiliary coordinate chart per system; in the sums below a frame means this chosen realization. Write $w = 2s \geq 64$. For every literal normal $N \triangleleft U$ and every pair kernel K , put

$$Q = U/N, \quad A = KN/N, \quad B = U/(KN), \quad S = A^B.$$

For $s = 32$ use Theorem 21.5; the remaining finite widths $64 \leq s < 2048$ are covered by Theorem 21.4. For $s \geq 2048$ the zero cut in Lemma 24.3 already suffices. Retain all cuts $C \leq S$ satisfying $2c + 4r(A/C) \leq 15s/16$, where $c = \dim C$. This menu is nonempty and invariant under the original action normalizer. The original pair top is a specified degree- s cover of B . Proposition 6.3 gives

$$e_Q(J) \leq D_C 2^{\lambda b} \Phi_C(J), \quad \Phi_C(J) = e_B(J) 2^{cd_2(J)}, \quad D_C = |A/C| |H^1(B, A/C)|, \quad \lambda = r(A/C)/2.$$

The module bound in Lemma 21.3 gives $t = \dim S \leq B_{\log s} \leq 5s/16$. Therefore

$$w/2 \leq v = s + 2c \leq 13w/16, \quad g = w - v - 8\lambda \geq w/32, \quad g \leq w - v \leq v. \quad (24.4)$$

We next bound the entire varying menu. Write $w = 2^k$. Lemma 24.2 bounds the number of literal normal subgroups of each action by $2^{(w-1)A_k}$ and gives an actual generating tuple of length A_k . Every such action is conjugate into a fixed Sylow subgroup of S_w , of order 2^{w-1} . Choose one original tuple for each action class after this conjugation. Equal tuples generate the same subgroup, so the number of transitive action classes is also at most $2^{(w-1)A_k}$. This argument requires no separate soluble normal-generation estimate. Furthermore

$$t \leq B_{\log s} = O(s/\sqrt{\log s}), \quad \log G_t = O(s^2/\log s),$$

$$\log D_C \leq \dim(A/C)(1 + d(B)) = O(w^2/\sqrt{\log w}).$$

The last inequality uses $d(B) \leq d(T) \leq A_{k-1}$ for the original transitive top T . More explicitly, every subspace of $\mathbb{F}_2^t$ has a spanning t -tuple, so there are at most 2^{t^2} cuts. Using the frame bound $w - 1 \leq 2^w$ and $1/a(U) \leq 1$, set

$$h(2^k) = 2(2^k - 1)A_k + B_{k-1}^2 + 2^{k-1}(1 + A_{k-1}) + 2^k = O(w^2/\sqrt{\log w}).$$

The cocycle bound here evaluates actual cocycles on generators of the same B before passing to their cohomology quotient. These estimates give

$$\sum_{[U], \deg U=w} \frac{1}{a(U)} \sum_{\text{frames}, N, C} D_C \leq 2^{h(w)}. \quad (24.5)$$

Every normal and every cut is included before taking this bound.

The graph proof of Proposition 6.3 uses the same source J for all maps. For binary J and binary covers its graph is binary, but may have fixed points; in particular $\sum_{J \leq S_b, J \text{ binary}} \Phi_C(J)^q \leq s_{b+qv}$. Put $e = g/16$, $a_0 = v/8 + e$, and declare the entry hot if $\Phi_C(J) > 2^{a_0 b}$. Use $q = \max(1, \lceil 8eb/v^2 \rceil)$ in (6.4). For $b > 0$ its rounding square is at most $v^2/16$, while (24.4) gives

$$g/v \geq 1/26, \quad b + qv \leq 3b/2 + v \leq 3n/2.$$

Consequently the main exponent of each hot term is at most

$$-2^{-18}b^2 - 87w^2/4096. \quad (24.6)$$

For $b = 0$ the strict hot event is empty. The normalized hot sum has the original factor

$$\frac{D_C}{b!a(U)c_{n/2}G_{n/2}} 2^{\lambda b - (q-1)a_0 b} s_{b+qv}.$$

Thus the coarse subgroup bound contributes $O(n^{3/2})$ and $c_{n/2}^{-1}$ contributes $O(n \log(n+2))$ to its logarithm; dropping $1/b! \leq 1$ is harmless for this upper bound. For every $\delta > 0$ there is $H_\delta < \infty$ such that $h(w) \leq \delta w^2 + H_\delta$ for all $w \geq 64$. Take $\delta = 1/128$ in (24.5). The retained negative terms in (24.6), together with $b^2 + w^2 \geq n^2/2$, absorb all errors and the at most n widths, yielding the stated hot scalar.

On the complement of the union of all hot events, the complete cold coefficient for $b = n - w$ is

$$\Delta(n, b) \sum_{[U], \deg U=w} \frac{1}{a(U)} \sum_{\text{frames}, N, C} D_C 2^{(w/8 - g/16)b}.$$

The source factorial cancels the $b!$ in the pointing factor once. The coefficient inequality $c_{b/2}/c_{n/2} \leq (2N)^{w/2}$ and $\log(G_{b/2}/G_{n/2}) = -wb/8 - w^2/16 + O(1)$ give its logarithm at most

$$h(w) + (w/2) \log n + O(1) - wb/512 - w^2/16.$$

Use $h(w) \leq w^2/128 + H$ and, uniformly for $64 \leq w \leq n$, absorb $(w/2) \log n + H + O(1)$ into $wn/8192$ for large n . The result is at most $-wn/1024$. Summing at most n widths proves (24.3), conservatively with rate $n/512$.

The hot union is selected first over all actual entries; its complement is then bounded by cold pointing. Deleting an actual orbit leaves a complete fixed-point-free binary subgroup, so the cold target is precisely $F_{b/2}$. This establishes the inequality before any bound on F has been assumed. $\square$

24.3 Finite witness coverage and its universal interpretation

Let $\mathcal{A}$ be the seventeen-action alphabet in Theorem 23.15.

Lemma 24.5 (Finite coverage principles). *Fix a power-of-two degree w . A list of literal transitive 2-groups containing a Sylow subgroup of S_w , and closed up to supplied permutation conjugacy under all transitive index-two subgroups, contains every transitive binary action up to conjugacy. For any finite 2-group U , a list of literal normal subgroups containing 1 and closed under all inverse images of order-two subgroups of $Z(U/N)$ contains every normal subgroup of U .*

Proof. The iterated binary wreath group on w points has order 2^{w-1} , which is the full 2-part of $w!$. Hence every binary action is conjugate into this explicit Sylow group. In a finite 2-group any subgroup lies at the bottom of a chain of index-two subgroups: choose successive maximal proper overgroups. Every overgroup of a transitive subgroup is transitive. Induction down the chain proves the first claim. All index-two subgroups are exactly the kernels of the nonzero functionals on $U/\Phi(U)$, so this closure can be checked exhaustively.

If $N < L$ are normal in U , the nontrivial normal subgroup L/N of U/N meets its centre nontrivially, by the class equation. This intersection contains an involution. Its inverse image is a normal index-two extension of N lying inside L . Repetition starting at 1 reaches every L , proving the second claim. $\square$

Lemma 24.6 (Uniform order stops at power-of-two degrees). *Let $k \geq 4$, $w = 2^k$, and let $U \leq S_w$ be a faithful transitive binary action with $|U| \leq 2^{w/2}$. For every original normal subgroup $N \triangleleft U$, put $Q = U/N$, $a = \log_2 |Q|$, and $z = \dim \Omega_1 Z(Q)$. Then at least one of the sufficient fusion conditions*

$$2a < w, \quad 38^{8z} < 2^w 25^{8z}$$

holds. Consequently, for any fixed finite collection of such original actions, all their normal axes admit the original-weight fusion estimate of Theorem 6.5, with constants that may depend on that collection. In particular this applies at degree sixteen to every transitive binary action of order at most 256.

Proof. If $N \neq 1$, then $|N| \geq 2$, so $|Q| = |U|/|N| \leq 2^{w/2-1}$ and $2a \leq w - 2 < w$. If $|U| \leq 2^{w/2-1}$, the same bound holds for every quotient, including $N = 1$.

It remains to consider $N = 1$ and $|U| > 2^{w/2-1}$. The elementary inequality $k + 1 \leq 2^{k-1}$ for $k \geq 4$ gives $2^{w/2-1} \geq 2^k = w$, so $|U| > w$. Every central subgroup acts freely on the original w points: a central element fixing one point fixes its entire U -orbit, and faithfulness makes that

element trivial. Hence $A = \Omega_1 Z(U)$ has order at most w . Equality would make A transitive. Any element of U fixing one point would then fix every point by commuting with A , so U would act regularly and have order w , a contradiction. Thus $|A| < 2^k$ and $z < k$. For $k = 4$, the exact integer inequality $38^{24} < 2^{16} 25^{24}$ gives $38^{8z} < 2^{16} 25^{8z}$ for every $0 \leq z \leq 3$. For $k \geq 5$, induction from $k = 5$ gives $8(k - 1) \leq 2^k = w$. If $z > 0$, use $38 < 50$ to obtain

$$38^{8z} < 2^{8z} 25^{8z} \leq 2^w 25^{8z},$$

since $8z \leq 8(k - 1) \leq w$. The case $z = 0$ is immediate.

For the order condition, Proposition 6.4 supplies $D = 1$, $v = 2a$, $\lambda = 0$, and $\Phi(J) = 2^{ad_2(J)}$. These are graph-moment marker coordinates; they do not assert that Q has a faithful action on $2a$ points. For the character condition, the actual-source estimate (24.7) below, with its stated nilpotent conjugacy-class bound, supplies $D = |\text{Aut } Q|$, $v = 0$, $\Phi(J) = 1$, and $\lambda = z \log_2(38/25) < w/8$. This estimate also holds for arbitrary $J \leq S_b$: restriction to an original Sylow 2-subgroup preserves surjectivity to Q and determines each map, since that Sylow subgroup maps onto the joint binary image of any two maps in $Q \times Q$. In both cases $w - v - 8\lambda > 0$, and all moments concern the same original J . Sum over the complete fixed finite collection of these original pairs (U, N) and apply Theorem 6.5. Its pointing weights retain $|N_{S_w}(U)|$, while the character branch retains $|\text{Aut } Q|$. No quotient normalizer or smaller faithful action is substituted for the original action. $\square$

Lemma 24.7 (Finite binary entry theorem). *For every transitive binary action U of width $w \in \{2, 4, 8, 16\}$ outside $\mathcal{A}$, and every literal normal $N \triangleleft U$, at least one of the following witnesses exists for $Q = U/N$:*

1. *An invariant pair system with kernel K , an actual central cut $C \leq (KN/N)^{U/(KN)}$, and an actual permutation cover of $B = U/(KN)$, such that $v_B + 2 \dim C + 4 \dim((KN/N)/C)^B < w$.*
2. *A faithful permutation representation of Q of degree $v < w$.*
3. *With $z = \dim \Omega_1 Z(Q)$, the exact integer inequality $38^{8z} < 2^w 25^{8z}$.*
4. *An equal-degree full-subdirect replacement by a word in $\mathcal{A}$, with actual maps onto Q , retaining at least one quarter of its points in noncritical factors.*

The last branch needs only the following four entries:

U	$ N $	Q	replacement word
8T16	2	SmallGroup(16, 3)	8T27
8T20	2	SmallGroup(16, 3)	8T27
8T21	2	SmallGroup(16, 3)	8T27
16T1086	4	SmallGroup(256, 7667)	$C_4[4], D_8[4]^3$

In the last row the cover is a proper full-subdirect subgroup of order 256 in the displayed product, not that whole product.

Proof. This is the finite computer-assisted lemma recorded by the literal binary entry certificate. We describe its mathematical coverage and the statements checked at each vertex; these are essential to its use. There are explicit iterated wreath roots in each of the four degrees. Each action record gives every transitive index-two child, an actual conjugator to its destination, and a checked basis of its Frattini quotient. Nonregular base actions retain their children. Regular

transitive actions have no transitive proper subgroup. For an index-two child $K \leq U$, a proposed conjugator to T needs only the checks $g^{-1}t_jg \in K$ on generators t_j of T , together with $2|T| = |U|$. They give $T \leq gKg^{-1}$, and the actual index forces equality. No full element-by-element conjugacy table is needed. Each nonbase action record gives 1 and every central-involution extension of each normal vertex. The two closure checks prove exhaustive action and normal coverage by Lemma 24.5; catalogue identifiers play no role in these implications.

For the first branch the record supplies the pair partition, kernel K , intersection $M = K \cap N$ and lifted cut $M \leq C_0 \leq K$. The literal equalities $[C_0, U] \leq M$ and $\ker(U \rightarrow \text{cover image}) = KN$ certify the same central section and quotient. Fixed sections are computed by commutators modulo M and C_0 , giving the integers c, r and the strict gap. For a direct witness the supplied generator images define a map with kernel exactly N and moved degree strictly below w . For a character witness the centre and its involution subgroup are reconstructed in the literal quotient, and the displayed integer inequality is checked. For a transport witness both specified maps are onto the same literal quotient, the original map has kernel N , and each displayed projection of the actual cover group equals its assigned action. The degree and noncritical fraction are then checked on the actual orbit partition. The four exceptional charts satisfy these conditions, including the retained proper cover in degree sixteen. The committed finite records and checker in Appendix A supply these finite identities.

There is also a uniform proof for regular actions of width $w \geq 8$: choose a central involution and its coset pairs. Then $K = C_2$, $A = KN/N$ is central of dimension at most one, and one may take $C = A$. The quotient B has its regular cover of degree at most $w/2$; $w/2 + 2 \dim A < w$. All binary actions of degrees two and four are already in $\mathcal{A}$. $\square$

The first two witness types have the graph envelopes of Proposition 6.3, with $g > 0$. The third has a direct slope strictly below $w/8$. Indeed a 2-group Q has a faithful tuple of z irreducible complex characters: choose constituents whose central characters span the dual of $\Omega_1 Z(Q)$. A nontrivial normal subgroup meets $\Omega_1 Z(Q)$, so their common kernel is trivial. For a binary $J \leq S_b$, inflation along an epimorphism yields a tuple from $\text{Irr}(J)^z$. Every tuple determines at most one kernel, and each kernel has exactly $|\text{Aut } Q|$ epimorphisms to Q . The nilpotent conjugacy-class bound [Mar05, Theorem 1.5] therefore gives

$$|\text{Epi}(J, Q)| \leq |\text{Aut } Q| k(J)^z \leq |\text{Aut } Q| 2^{z \log(38/25)b}. \quad (24.7)$$

The strict integer test is exactly $z \log(38/25) < w/8$. For $Q = 1$ use $e_Q(J) = 1$. These assertions concern the actual source J ; no faithful action of an abstract quotient on its points is assumed.

24.4 Small widths and the complete recurrence

Theorem 24.8 (Complete binary recurrence). *There are nonnegative $\eta_B(N)$ and $K_B(N, M)$, supported on $0 \leq M < N$, and constants $a_B, C_B > 0$, such that eventually*

$$E_N \leq \eta_B(N) + \sum_{M < N} K_B(N, M) F_M, \quad \eta_B(N) + \sum_{M < N} K_B(N, M) \leq C_B 2^{-a_B N}. \quad (24.8)$$

In particular F_N is bounded and $E_N = O(2^{-a_B N})$.

Proof. Power-of-two orbit sizes give the exact disjoint partition

$$E_N = T_{64}(N) + T_{32}(N) + S_{16}(N),$$

where S_{16} has all actual orbit widths at most sixteen and at least one noncritical orbit. The first two parts are bounded by (24.3) and (24.2).

For S_{16} consider a nonbase orbit U and its actual axis $N = H \cap U$. Include a fixed finite invariant menu of all witness transports in Lemma 24.7; choose the transports before counting the complementary source. First take the union in which some actual graph entry is hot. Theorem 6.5 bounds this complete union by a scalar $2^{-\Omega(N^2)}$. On its complement, point any orbit with an accepted strict local entry. For graph entries the cold coefficient is

$$\Delta(2N, 2N - w) \frac{D}{a(U)} 2^{(w/8 - g/16)(2N - w)}.$$

For the direct character entry it is the same coefficient with $D = |\text{Aut } Q|$ and exponent $z \log(38/25)(2N - w)$. Their original Gaussian ratio leaves respectively $-g(2N - w)/16$ or $-(w/8 - z \log(38/25))(2N - w)$, with error $O(\log(N + 2))$. The fixed finite menu has a positive minimum gap. Summing every literal normal and every witness thus yields a kernel K_s with row $2^{-\Omega(N)}$, supported at $N - 4, N - 8$; only nonnegative targets are included. Entries with zero prefix degree have a direct bound and need no hot moment. Deleting the original orbit again leaves the complete fixed-point-free binary count $F_{N-w/2}$.

In the remaining family every nonbase orbit has a transport witness for its exact axis. Apply all these maps simultaneously, leaving $\mathcal{A}$ factors untouched. The product of the embedded axes lies in H , so Proposition 22.1 recovers H as the full preimage of the joint quotient image. Every replacement has the same physical degree, and retains at least one quarter of its noncritical support. Theorem 23.15 and the stronger profile envelopes (23.14), inserted in Proposition 22.4, give a scalar $2^{-\Omega(N)}$ for the entire remaining family. This also includes words already contained in $\mathcal{A}$, by identity transport. The small original noncritical-support range is removed before charging the decoration factor.

Consequently $S_{16}(N) \leq \eta_s(N) + \sum_M K_s(N, M) F_M$, with both its scalar and complete row exponentially small. Add this to the two disjoint large-width bounds to obtain (24.8). No estimate of F_M has been used to prove the kernel rows.

Finally C_{2M}^{crit} is bounded by the independently proved critical-family theorem. Substitute $F_M = C_{2M}^{\text{crit}} + E_M$ in the recurrence. Beyond an onset its row is at most $1/2$, and its forcing is bounded. A constant dominating twice that forcing and all finitely many initial E_M bounds E_N by induction. Thus F_N is bounded. Substituting this established bound back into the same recurrence proves exponential decay. $\square$

25 Direct fusion for original binary orbits

The following cut-or-character alternative gives a shorter route to original local entries from width 32. It uses the deterministic separator and the actual action kernel, without the rank-moment calculation or the adaptive degree-32 cut. Its recurrence has complete ordinary subgroup counts at smaller degrees as targets. Thus it can be inserted directly in the ordinary recurrence; the independent binary recurrence of the preceding section has a different target and remains a separate statement.

25.1 An original cut-or-character alternative

Write $B_j = \binom{j}{\lfloor j/2 \rfloor}$ and $\gamma = 5^{2/7}$. The permutation class estimate used in Proposition 8.3 implies the weaker bound $k(P)^7 \leq 5^{2b}$ for every actual binary subgroup $P \leq S_b$. Consequently, for a finite

binary group Q , with $z = \dim \Omega_1 Z(Q)$, the original character envelope is

$$|\text{Epi}(J, Q)| \leq |\text{Aut } Q| 2^{\alpha(z)b}, \quad \alpha(z) = \frac{2z}{7} \log_2 5, \quad J \leq S_b. \quad (25.1)$$

Here J is arbitrary: restriction to its original Sylow 2-subgroup is injective on epimorphisms to Q and remains surjective onto Q . The same bound holds for any specified surviving subfamily by inclusion. This is the separate γ envelope, not a replacement of every previously accepted 38/25 criterion.

Theorem 25.1 (Original cut-or-character alternative). *Let $U \leq S_{2h}$ be transitive and binary, where $h = 2^k$ and $k \geq 4$. Fix an original invariant pair frame, with flip kernel K and faithful transitive top $T \leq S_h$. For each original $N \triangleleft U$, retain*

$$Q = U/N, \quad A = K/(K \cap N), \quad B = U/(KN),$$

including the actual map $T \twoheadrightarrow B$ and the actual kernel chart $A \longrightarrow Q \twoheadrightarrow B$. At least one of the following alternatives holds.

1. *There is a literal subspace $C \leq A^B$ such that*

$$2 \dim C + 4 \dim(A/C)^B \leq h - 2.$$

2. *The action of T on A is faithful, the whole center of Q lies in A , and*

$$\Omega_1 Z(Q) = A^B, \quad z = \dim \Omega_1 Z(Q) \leq B_k, \quad 5^{16z} < 2^{7(2h)}.$$

In particular each original N has a same-source fusion entry with an even prefix $v < 2h$ and a strictly positive gap. No splitting of $Q \rightarrow B$ is assumed.

Proof. Put $S = A^B$, $t = \dim S$, $\ell = \dim(A/S)^B$, and $D_0 = 2B_{k-1}$. These are dimensions of the same actual section, so $t + \ell \leq \dim A$. The original permutation-section bound and its one-character kernel restriction give $t \leq B_k \leq D_0$ and the complete displacement line-slice bound $D_0 - t$. The deterministic separator supplies q independent functionals on S , with annihilator C , such that

$$\dim C = t - q, \quad \dim(A/C)^B = q, \quad 2q \leq \ell + \max\{1, D_0 - t\}.$$

The fixed-quotient identity uses the full displacement slice; it does not merely test each scalar line separately.

If $2 \dim A \leq h$, the resulting even cost satisfies

$$2 \dim C + 4 \dim(A/C)^B = 2t + 2q \leq t + \ell + D_0 + 1.$$

The inequalities $8D_0 \leq 3h$ and $h \geq 16$ make the last expression at most $h - 1$. Evenness gives the first alternative. The inequality $t \leq D_0$ is retained here; truncating a negative difference would not justify the displayed estimate.

Suppose instead that $2 \dim A > h$. Let H be the actual normal kernel of T acting on A . If $H \neq 1$, its orbits on the original h points have a common size $2^a \geq 2$: normality gives uniformity, and faithfulness of T excludes singleton orbits. The original permutation preimage of A , restricted to H , maps onto the trivial H -module A . Applying the orbit-section bound and $2B_a \leq 2^a$ therefore

gives $2 \dim A \leq h$, a contradiction. Thus T acts faithfully on A . Since this action factors through the actual surjection $T \rightarrow B$, that map is injective as well.

For any central element of the original Q , its image in B acts trivially on the literal kernel A . Faithfulness places the element in A . The actual kernel chart now identifies $Z(Q)$ with A^B ; in particular this center is elementary. Finally $8B_k \leq 3h$ gives $16z \leq 3(2h)$, and $125 < 128$ gives the strict integer test. $\square$

For the first alternative put $c = \dim C$, $r = \dim(A/C)^B$ and choose the entry

$$v = h + 2c, \quad \lambda = r/2, \quad D_C = |A/C| |H^1(B, A/C)|, \quad \Phi_C(J) = |\text{Epi}(J, B)| 2^{cd_2(J)}. \quad (25.2)$$

The original central extension and its actual quotient by C give the envelope $D_C 2^{\lambda b} \Phi_C(J)$. Its full cocycle factor is retained. The faithful original cover $T \leq S_h$ mapping onto the same B gives all common-source moments

$$\sum_{J \leq S_b} \Phi_C(J)^q \leq s_{b+qv} \quad (q \geq 1).$$

There is no assertion that B itself has a faithful action of degree h . In the character alternative take

$$v = 0, \quad \lambda = \alpha(z), \quad D_C = |\text{Aut } Q|, \quad \Phi_C(J) = 1.$$

Its moments equal s_b . In both alternatives define

$$e = \frac{2h - v - 8\lambda}{16} > 0.$$

The cut alternative has $2h - v - 8\lambda \geq 2$, but no such uniform lower bound is asserted for the character alternative. For each fixed finite collection of widths, positivity supplies a finite-menu minimum. All entries are chosen before the exterior degree and the exterior subgroup; the original normalizer is unchanged.

25.2 The ordinary forward recurrence for binary orbits

Let s_n denote the number of all subgroups of S_n , let $L_n > 0$ be the exact parity-dependent benchmark, and put $\sigma_n = s_n/L_n$. Let $\mathcal{A}_n$ be the family of literal $H \leq S_n$ having an actual orbit of size at least 32 whose restriction image is binary. The entire H , its complementary restriction, and their correlation are unrestricted.

Theorem 25.2 (Original ordinary binary-orbit recurrence). *There is a nonnegative row $K(n, m)$, with $K(n, m) = 0$ when $m \geq n$, such that for $n \geq 2048$,*

$$\frac{|\mathcal{A}_n|}{L_n} \leq \sum_{m < n} K(n, m) \sigma_m. \quad (25.3)$$

There are constants $C, \kappa > 0$ for which

$$\sum_{m < n} K(n, m) \leq C 2^{-\kappa n}$$

for all sufficiently large n . In particular the row sum is eventually at most $1/2$. Any subfamily defined by additional conditions on the original H obeys the same recurrence by inclusion.

Proof. An actual binary transitive orbit has power-of-two degree. For each width $w = 2h \in \{32, 64, 128, 256, 512\}$, choose one representative of each original action-conjugacy class and one invariant pairing for each representative. Apply Theorem 25.1 to every literal original normal subgroup. Relabelling the actual orbit and its actual complement gives physical coverage by these families. This is an upper union of original subgroups; no uniqueness of the chosen orbit or of its chart is required.

For a fixed representative U , normal axis N and exterior degree $b = n - w$, use only the first same-source moment. Its exact contribution at the target degree $m = b + v$ is

$$\frac{n!}{b! |N_{S_w}(U)|} D_C 2^{\lambda b} \frac{L_{b+v}}{L_n}. \quad (25.4)$$

Every original normal is retained, including distinct summands with the same target degree. Since $v < w$, every such target is smaller than n . The exact benchmark ratio and $e > 0$ give exponential decay for this genuinely fixed finite menu, without an a priori estimate on the sequence s_m .

The widths 1024 and 2048 use their already proved positive zero-cut entries and the same finite first-moment argument. For widths at least 4096 use the original wide zero-cut installation and its proved normalizer-weighted menu aggregate. The actual zero entries are bounded by that full nonnegative row; numerical acceptance of extra entries is not used as a substitute for a local envelope. Its row sum has the proved uniform exponential bound.

Add these rows and cover the literal family $\mathcal{A}_n$ by the finite-width and wide sectors. Overlap only enlarges the upper bound. Adding their exponential estimates gives the asserted C, κ and eventual contraction. All denominators in (25.4) remain the original symmetric normalizers. $\square$

26 A joint cyclic-four character moment

For a finite group J , let $d(J), c(J)$ be the first two conjugate columns of the binary part of J^{ab} , and put

$$q(J) = d(J) + c(J) = \log |\text{Hom}(J, C_4)|, \quad M_4(b, r) = \sum_{J \leq S_b} 2^{r q(J)}, \quad F(b, r) = \frac{b^2}{16} + \frac{br}{4} + \frac{r^2}{2}.$$

The two source columns must be retained jointly; the second column is not separately additive in an extension.

Theorem 26.1. *As $b \rightarrow \infty$,*

$$\log M_4(b, r) \leq F(b, r) + o((b + r)^2),$$

uniformly for r/b in any fixed bounded interval. The range $r \geq b/2$ also satisfies the bound directly, without an upper restriction on r/b . Literal homomorphisms are counted, without a target automorphism divisor.

26.1 The retained normal columns

We use the bounded-word counting construction of [RDT25, Sections 3–5 and Proposition 7.4], specifying the changes needed for the character weight. Fix a physical orbit bound C . In its nonexcess case, let Q_i be quotients of actual transitive binary groups of degrees $n_i < C$, such that every Q_i -normal subgroup has normal-generator number at most $n_i/4$. Put $n = \sum_i n_i$. For an

efficiently ordered upper normal tableau write N_i for its diagonal normal subgroup, $d_i = d_{Q_i}(N_i)$, and $c_i = \log |N_i|/n_i$. The efficient-generator division before the final maximization in that construction gives

$$E = \sum_{j < i} (c_i - c_j) n_j d_i, \quad 0 \leq c_1 \leq \dots \leq c_t \leq 1, \quad d_i \leq \min(n_i/4, c_i n_i) \quad (26.1)$$

as an upper bound on the logarithm of the number of groups with a fixed tableau. All tableau and ordering choices cost $O_C(n \log(n+2))$ by its normal-chain bound.

Lemma 26.2. *For a full subgroup L with this tableau,*

$$d(L) \leq D := \sum_i d_i, \quad d(L) + c(L) \leq S := \sum_i \min(2d_i, c_i n_i).$$

Proof. Filter L by the kernels of successive tail projections. The i th quotient is N_i with conjugation action supplied by the onto projection to Q_i . Restrict global characters $L \rightarrow C_4$ along this filtration. Two characters with the same previous restriction differ on the next term by an invariant character of N_i , hence by a character of $N_i/[N_i, Q_i]$. The latter abelian group has at most d_i generators and order at most $2^{c_i n_i}$, so contributes at most $2^{\min(2d_i, c_i n_i)}$ maps. Multiplying the fibre bounds proves the second inequality. The same argument with C_2 proves the first. It bounds extendable global restrictions and does not presume that every invariant character extends. $\square$

Put $T(t) = \sum_{c_i \geq t} n_i$, $0 < t < 1$. Layer-cake identities and (26.1) imply

$$E \leq \frac{1}{4} \int_0^1 T(t)(n - T(t)) dt, \quad D \leq \int_0^{1/4} T(t) dt, \quad S \leq \int_0^{1/2} T(t) dt. \quad (26.2)$$

For example the first integral before its factor $1/4$ is $\sum_{j < i} (c_i - c_j) n_i n_j$. It follows already that $E + r q(L) \leq n^2/16 + nr/4 + r^2/2$, by maximizing separately on $0 < t < 1/2$ and $1/2 < t < 1$.

26.2 The abelian packet in the same tableau

Let A be the product of the abelian physical coordinate quotients, on total support a , and put $B = A \times C_4^r$. The abelian quotient bound gives $\log |A| \leq a/2$. For a fixed remaining projection L , the exact number of subgroups of $B \times L$ onto L is

$$X_B(L) = \sum_{D \leq B} |\text{Hom}(L, B/D)|. \quad (26.3)$$

This follows by taking the actual intersection D and the literal graph map. Write x_j for the conjugate columns of L_2^{ab} . The Hall subgroup formula, after dropping only the monotonicity restriction on quotient columns, bounds its logarithm by

$$\sum_j \Psi(B_j, x_j) + O_C(\log(a+r+2)), \quad \Psi(v, x) = \max_{0 \leq y \leq v} y(v - y + x) = \begin{cases} vx & v \leq x, \\ (v+x)^2/4 & v \geq x. \end{cases}$$

There are boundedly many columns since the physical degrees are bounded. The Gaussian factors and the number of column profiles give the stated logarithmic error.

The derivative of Ψ in its first argument increases in both arguments. Since $B_1 \geq B_j$ and $x_1 \geq x_j$, moving each unit of the original A -column mass into the first column can only increase the bound. Fill any remaining mass up to $a/2$. This gives

$$\log X_B(L) \leq \Psi(a/2 + r, d(L)) + \Psi(r, c(L)) + O_C(\log(a+r+2)). \quad (26.4)$$

Let the two maximizing Hall ranks be $\alpha \geq \beta$; this ordering is possible because both the ambient and source columns are ordered. Lemma 26.2 gives the joint inequality

$$\alpha d(L) + \beta c(L) \leq (\alpha - \beta)D + \beta S \leq \alpha \int_0^{1/4} T(t) dt + \beta \int_{1/4}^{1/2} T(t) dt.$$

Define

$$g_n(x) = \max_{0 \leq z \leq n} \{z(n-z)/4 + xz\} = \begin{cases} n^2/16 + nx/2 + x^2 & x \leq n/4, \\ nx & x \geq n/4. \end{cases}$$

The tableau entropy and the full abelian fibre together are bounded by

$$\frac{1}{4}g_n(\alpha) + \frac{1}{4}g_n(\beta) + \frac{1}{2}g_n(0) + \alpha(a/2 + r - \alpha) + \beta(r - \beta).$$

Direct differentiation on the two branches proves

$$\max_{0 \leq x \leq v} \{x(v-x) + g_n(x)/4\} = (v + n/4)^2/4.$$

The maximizer is $x = v$ when $v \leq n/4$, and $x = (v + n/4)/2$ otherwise. Relaxing the order between α, β now gives the upper bound

$$\frac{n^2}{32} + \frac{(a/2 + r + n/4)^2 + (r + n/4)^2}{4} = F(n + a, r) - \frac{na}{16} \leq F(n + a, r). \quad (26.5)$$

This is the required abelian/nonabelian splice on one source.

26.3 Excessive quotients and bounded binary words

For a product D of actual transitive binary actions of degree less than C , of total degree b , we claim

$$\log |\text{Sub}(D \times C_4^r)| \leq F(b, r) + O_C((b + r) \log(b + r + 2)). \quad (26.6)$$

First condition on coordinate projections, refining any intransitive projection into its actual orbits. This costs $2^{O_C(b)}$ and does not increase physical support. Follow the literal normal-intersection construction in [RDT25, Proposition 7.4] on the physical coordinates, keeping the auxiliary factor C_4^r throughout. Quotienting the embedded axes is injective because their product lies in the subgroup. In particular those axes are already killed by every auxiliary graph coordinate; no character is lost by quotienting first. The nonexcess branch is (26.5).

In the excessive branch choose the physical coordinate of degree w in the priority of cases (IIa)–(IIc) of that proof. Its nonabelian quotient Q has, by [RDT25, Theorem 5.7], a normal subgroup F_0 and parameters e, z, g satisfying

$$z = \log |C_Q(F_0)|, \quad g = \log |[F_0, Q]|, \quad ez + g \leq w/2, \quad z \leq w/4.$$

The three values of e are 3 for the degree-eight excess-two case, 1 for excess one, and $5/4$ for the remaining excess-two case; the excess-one branch retains the generator-equality hypothesis of that theorem.

In the remaining nonabelian projection select a maximal fully independent tail. Let b_3 be the degree of its dependent physical coordinates, and b_4 the degree of the independent tail together with every abelian physical coordinate. Thus $b_3 + b_4 = b - w$. The normal filtration in [RDT25, Theorem 5.5] gives, on this same source, the bounds $eb_3/4 + b_4/2 + r$ for ordinary generators and

$b_3/4$ for the derived-normal term in its epimorphism estimate. To check the auxiliary modification: the abelian-coordinate kernel contributes at most $b_4/2 + r$ generators, whereas commutators have trivial abelian coordinates, so the derived group projects isomorphically onto the derived nonabelian projection. Passing to quotients does not increase the relative normal-generator caps. No faithful action of Q at degree w is assumed.

The epimorphism estimate [RDT25, Theorem 4.6] consequently becomes

$$\log |\text{Epi}(\text{source}, Q)| \leq \frac{w(b_3 + b_4)}{8} + \frac{wr}{4} + O_C(\log(b + r + 2)).$$

This is paid by the exact reserve $F(b, r) - F(b - w, r) = w(b - w)/8 + wr/4 + w^2/16$. Induct on the number of physical coordinates. The lower normal-tableau chains absorb the normal-axis sums exactly as in equation (7.1) of the cited proof; their count and the reorderings have logarithm $O_C(b \log(b + 2))$. The accumulated epimorphism errors are $O_C(b \log(b + r + 2))$. This proves (26.6).

26.4 The weighted generation reduction

Proof of Theorem 26.1. If $G = \langle K, x_1, \dots, x_l \rangle$, restriction and the displayed generator values determine a character. Thus

$$|\text{Hom}(G, C_4)|^r \leq 4^{rl} |\text{Hom}(K, C_4)|^r. \quad (26.7)$$

Choose one encoding for each G and bound by the sum of all encodings. This avoids requiring any unproved closure property of a marked family.

For soluble groups in bounded orbit containers, the nilpotent generation theorem and the socle-length bound of [RDT25, Theorem 7 and Lemma 8.5] supply a nilpotent subgroup and $O_C(\sqrt{b})$ additional generators. The ambient soluble group has order $2^{O(b)}$. The tuple choices and (26.7) cost $2^{O_C(b^{3/2} + r\sqrt{b})}$. For a bounded-orbit nilpotent group, the Cartesian Sylow orbit grids of [RDT25, Lemma 2.7] compress the binary and odd actions to degrees m, d with $m + d \leq b$. The grid choices cost $2^{O_C(b \log(b + 2))}$. The odd factor has no nontrivial C_4 character. Graph injection and (26.6) bound the binary moment by $2^{F(m, r) + o_C(b^2)}$ for $r = O(b)$; (1.1) bounds the odd factor by $2^{d^2/16 + O(d^{3/2})}$. Since $F(m, r) + d^2/16 \leq F(b, r)$, the bounded soluble case follows.

For the large-orbit step fix $\eta > 0$. The soluble generator estimate [RDT25, Proposition 6.2] supplies a cutoff C such that the large-orbit projection requires at most $\eta b/\delta$ generators, where $\delta = (\log 24)/3$ is the soluble order constant. Its kernel is an actual subgroup on the small support. Working in a product of maximal transitive soluble containers, the choices of lifts and their marked values cost at most $2^{\eta b^2 + 2\eta r b/\delta}$ by (26.7). The orbit partitions and container choices cost $2^{O(b \log b)}$ by [RDT25, Theorem 8.6]. The bounded-source estimate is at most $F(b, r)$ since F increases with physical degree. First take $b \rightarrow \infty$ with η, C fixed, and then $\eta \rightarrow 0$. This proves the soluble result uniformly for $0 \leq r \leq K_0 b$. It does not assert a uniform $O(b^{3/2})$ error as C varies.

Finally every finite group is generated by a soluble subgroup and one additional element by [RDT25, Theorem 8.4]. The additional costs $b!$ and 4^r are $2^{o(b^2)}$ in this range. For $r \geq b/2$, the direct bound $q(J) \leq b/2$ gives $\log M_4(b, r) \leq b^2/16 + br/2 + O(b^{3/2})$. Its leading expression is at most $F(b, r)$ because their difference is $r(2r - b)/4 \geq 0$. $\square$

27 Markers, binary moments, and the one-triple audit

27.1 Exact natural- S_3 transport

Proposition 27.1. *Let $X \leq T \times S_3^g$ be full on T and every marker. Put $K_X = X \cap C_3^g$, let $\overline{X}$ be its image in $T \times C_2^g$, and let $L = \overline{X} \cap C_2^g$, of dimension ℓ . Then*

$$\overline{X} \cong T \times L, \quad X/O^2(X) \cong (T/O^2(T)) \times C_2^\ell, \quad d_2(X) = d_2(T) + \ell, \quad \tau_2(X) = \tau_2(T).$$

For one marker the numbers of possible X with $\ell = 1, 0$ are respectively 1 and $2^{d_2(T)} - 1 + |\text{Epi}(T, S_3)|$.

Proof. The graph map $T \rightarrow C_2^g/L$ factors through $A_2(T)$ and has a linear lift, proving the first isomorphism. The kernel K_X has odd order, so inflation is an isomorphism in binary cohomology. The maximal binary quotient is therefore that of $T \times L$. Künneth and the identical elementary factor show that the kernel defining τ_2 is unchanged. For one marker Goursat's lemma gives: the full product; one common- C_2 graph for each nonzero binary character of T ; and one literal graph for each epimorphism $T \rightarrow S_3$. These are disjoint. $\square$

If $p_{T,g,\ell}$ counts the actual marker subgroups with this value of ℓ , their exact attachment to a critical product D , with f fixed points, has weight

$$\frac{1}{6^g g! f!} \sum_{\ell} p_{T,g,\ell} N_D((T/O^2(T)) \times C_2^\ell). \quad (27.1)$$

This identity does not license replacing an arbitrary complete exterior projection by its unmarked count.

27.2 A character moment on the complete binary error

Let $\mathcal{E}_N$ be all fixed-point-free 2-subgroups of S_{2N} with at least one actual orbit outside Table 1, and put

$$E_N = \frac{|\mathcal{E}_N|}{L_{2N}}, \quad pE_N = \frac{1}{L_{2N}} \sum_{Y \in \mathcal{E}_N} t(Y),$$

where $t(Y)$ is the number of its actual two-point orbits. Set $E_0 = pE_0 = 0$.

Proposition 27.2 (Complete binary pair moment). *For $N \geq 2$,*

$$pE_N \leq [7 + (6N)^{1/4}]E_N + r_N E_{N-1}, \quad r_N = \frac{N-1}{4} \frac{c_{N-1} G_{N-1}}{c_N G_N}. \quad (27.2)$$

The same estimate holds for the family of all fixed-point-free binary subgroups, with its corresponding counts.

Proof. Each pair orbit supplies a nonzero character of Y . Let $q(Y)$ be the number of distinct such characters, $D(Y)$ the number of unordered pairs of equal characters, and $J_4(Y)$ the number of ordered selections of four pair orbits with independent characters. Then

$$t(Y) - q(Y) \leq D(Y), \quad q(Y) \leq 7 + J_4(Y)^{1/4}.$$

For the second inequality, choose one original pair orbit for each distinct character. The span of at most three independent characters contains at most seven nonzero characters, so each of four

successive choices has at least $q - 7$ possibilities. Thus $J_4(Y) \geq (q - 7)^4$ when $q \geq 7$; the case $q < 7$ is immediate.

There is an injection from groups Y with four ordered independent pair orbits to groups H with one selected E_8 orbit and an ordering of its four center-pair orbits, at the same degree. For each ordered pair partition $\mathcal{P}$, fix a point chart once and use it to place an extraspecial group $E(\mathcal{P})$ on the eight selected points. Explicitly, inside $C_2^4 \rtimes S_4$, take the even flips semidirect the regular block V_4 . Its centre has exactly these four pair orbits. Fix, once for each ordered partition, an isomorphism $E(\mathcal{P})/Z(E(\mathcal{P})) \cong C_2^4$. Replace $Y \leq C_2^4 \times J$ by its full preimage in $E(\mathcal{P}) \times J$. Independence gives the full E_8 projection. The centre of this local projection, its retained pair order, and the fixed isomorphism recover Y . The complete exterior projection is unchanged, so its noncritical orbit is retained. There are at most $\lfloor N/4 \rfloor$ possible selected E_8 orbits, each with 4! possible orders. Consequently

$$\sum_{Y \in \mathcal{E}_N} J_4(Y) \leq 6N|\mathcal{E}_N|.$$

Concavity of $x^{1/4}$ gives $\sum_Y q(Y) \leq [7 + (6N)^{1/4}]|\mathcal{E}_N|$.

Collapse two pair orbits carrying the same character to one pointed pair orbit. This operation and its inverse preserve a noncritical orbit. The exact labelled weight ratio is

$$\frac{1/(2^2 2!)}{1/2} = \frac{1}{4}.$$

Thus

$$\frac{1}{(2N)!} \sum_{Y \in \mathcal{E}_N} D(Y) = \frac{1}{4(2N-2)!} \sum_{Y \in \mathcal{E}_{N-1}} t(Y) \leq \frac{N-1}{4(2N-2)!} |\mathcal{E}_{N-1}|.$$

Divide by $c_N G_N$ and combine the two bounds. □

For $n = 2j + 1$, put $p_{2j+1} = c_j + c_{j-1}/6$ and

$$\alpha_j = \frac{c_j}{p_{2j+1}}, \quad \beta_j = \frac{c_j}{3p_{2j+1}}, \quad u_j = \alpha_j + \beta_j[7 + (6j)^{1/4}], \quad v_j = \beta_j r_j. \quad (27.3)$$

A single fixed point contributes $\alpha_j E_j$. Replacing a pointed pair by a natural S_3 orbit has weight ratio $(1/6)/(1/2) = 1/3$ and contributes $\beta_j p E_j$. Thus their combined binary-error contribution is at most $u_j E_j + v_j E_{j-1}$. Section 33 gives $c_{j-1}/c_j \sim (96j)^{1/4}$, so $u_j \rightarrow 1$ and $v_j \rightarrow 0$. For the recurrence closure one can avoid this sharper asymptotic input: the elementary coefficient and Gaussian-ratio bounds give, uniformly,

$$0 \leq u_j \leq 4(j+1), \quad 0 \leq v_j \leq 2\Phi^{-2}(j+1)^3.$$

Either fixed polynomial is absorbed by half of any positive exponential rate. Thus the odd transfer introduces no additional analytic frontier. In particular this is not a strict contraction on unrestricted complete counts. In the final assembly, native forward recurrences for E_j and E_{j-1} are substituted first; the two target rows are padded to the ambient degree and these polynomial factors are absorbed into their exponential rates.

27.3 The split one-triple incidence

For a finite group K , a nonzero character $\chi : K \rightarrow C_3$ is split if it has a section. This is equivalent to being nonzero on an element of order three. The nonsplit characters, including zero, are

therefore the annihilator of the span of the order-three elements in $A_3(K)$. They form a vector space. Write its dimension as $h_3(K)$ and set

$$\sigma_3(K) = 3^{d_3(K)} - 3^{h_3(K)}. \quad (27.4)$$

This counts oriented split nonzero characters exactly.

Proposition 27.3 (Inverse-complement incidence). *For any collection $\mathcal{P}$ of pairs (K, χ) with $K \leq S_m$ and χ split and nonzero, define*

$$q(K, \chi) = |\{x \in K : |x| = 3, \chi(x) = 1\}|.$$

Then

$$|\mathcal{P}| = \sum_{|x|=3} \sum_{\substack{N^x=N, \ N \cap \langle x \rangle = 1 \\ (N \langle x \rangle, \chi_{N,x}) \in \mathcal{P}}} \frac{1}{|\{y \in Nx : |y| = 3\}|}, \quad (27.5)$$

where $\chi_{N,x}$ kills N and sends x to $1 \in C_3$. Furthermore, if $P \triangleleft K$ is a 2-group, inflation from K/P preserves d_3 , h_3 , and σ_3 .

Proof. For each (K, χ) insert the identity $1 = \sum_{x:|x|=3, \chi(x)=1} q(K, \chi)^{-1}$. With $N = \ker \chi$, the pair (N, x) determines both K and χ , and the denominator is precisely the one displayed in (27.5). Every ternary character kills P . A split character upstairs has an order-three representative whose image is a representative downstairs. Conversely, choose an order-three element y downstairs on which the character is nonzero and lift it to $x \in K$. Then $x^3 \in P$, so $(x^3)^{2^a} = 1$ for some a . The element $z = x^{2^a}$ satisfies $z^3 = 1$, and its character value is $2^a \chi(y) \neq 0$ in $\mathbb{F}_3$. Thus z has order three and the inflated character splits. This proves splitness in both directions and hence all three equalities. $\square$

These identities are required tests for applying a counting estimate to the one-external-triple case, customarily denoted $c = 1$. One must bound the sum of (27.4), not an unmarked subgroup sum, and must retain the reciprocal in (27.5). For example, on the shared-top group $J = \mathbb{F}_4^r \rtimes C_3$ and target $\mathbb{F}_4^m \rtimes C_3$, above the fixed top map the exact cocycle and onto-lift counts are

$$4^{m(r+1)}, \quad 4^m \prod_{i=0}^{m-1} (4^r - 4^i),$$

respectively. Maschke's theorem gives $H^1(C_3, \mathbb{F}_4^m) = 0$; $\text{Hom}_{C_3}(\mathbb{F}_4^r, \mathbb{F}_4^m)$ has 4^{mr} elements, and the 4^m coboundaries are literal distinct graphs. A second mark on the same J produces a joint moment on that source. It cannot be replaced by two independently chosen source groups. These exact tests do not themselves prove that the high-incidence $c = 1$ domain has been exhausted.

28 The complete marker sum

Here the binary exterior includes every even orbit, including the critical orbits and their already assembled attachments. Thus no terminal function is left to be multiplied into the counting comparisons below.

For $a \geq 1$ define the full-coordinate ternary fibre

$$L_3(a) = \sum_{\substack{K \leq \mathbb{F}_3^a \\ \text{pr}_i K = \mathbb{F}_3 \ (1 \leq i \leq a)}} 3^{a - \dim K}, \quad L_3(0) = 1.$$

For $g \geq q \geq 1$ put

$$H_{g,q} = \sum_{\substack{a_1 + \dots + a_q = g \\ a_i \geq 1}} \prod_{i=1}^q \frac{L_3(a_i)}{a_i!}, \quad H_{0,0} = 1,$$

and put $H_{g,0} = 0$ for $g > 0$.

Proposition 28.1 (Exact collapse of repeated signs). *Let $\mathcal{Y}_m$ be a collection of fixed-point-free binary subgroups on m points, closed under conjugation by S_m . Let $t_q(Y)$ count the q -element sets of actual pair orbits of Y whose characters on Y are distinct. For $n = m + 3g + f - 2q$, the number of subgroups obtained by replacing these selected pairs by g natural S_3 orbits in q sign classes, and adding f fixed points, is exactly*

$$\frac{n!}{m!} \frac{2^q H_{g,q}}{6^g f!} \sum_{Y \in \mathcal{Y}_m} t_q(Y). \quad (28.1)$$

Every group so counted recovers Y , the selected pairs, the sign classes, and all its ternary extension data.

Proof. Write the original group as $X \leq T \times S_3^g$, with T its complete projection on the other moved orbits. The image B of X in $T \times C_2^g$ is a binary group. Its marker signs are nonzero characters. Group equal signs into classes of sizes $a_1, \dots, a_q$. Deleting repeated signs is injective on B ; realize each retained sign on a pair. The resulting group is $Y \leq S_m$ and the newly introduced pairs give the selected set.

Every B -invariant subspace of C_3^g splits across these distinct sign isotypes by an explicit projection. Regard the signs as scalars in $\mathbb{F}_3$. For distinct signs χ, ψ , choose $b \in B$ with $\chi(b) \neq \psi(b)$. The operator

$$(\chi(b) - \psi(b))^{-1} (b - \psi(b)I)$$

preserves the subspace, is the identity on the χ -isotype and vanishes on the ψ -isotype. Multiplying one such diagonal operator for every other distinct sign gives the exact χ -projection. These projections sum to the identity, with each distinct sign counted once, regardless of its coordinate multiplicity. In an isotype of dimension a_i , the intersection K_i must have full coordinate projections. Indeed, lift an original marker 3-cycle and raise it to a sufficiently large power of 4: its binary image vanishes and the 3-cycle is unchanged. Conversely, a full ternary kernel coordinate together with a nontrivial sign generates the full original S_3 coordinate.

For a fixed invariant kernel $K = \bigoplus_i K_i$, let P be the whole original preimage of B . Subgroups of P with image B and kernel K correspond exactly to sections of $P/K \rightarrow B$. The fixed transpositions on the marker coordinates, together with the unchanged exterior coordinate, give one such section. Relative to this section the fibre is exactly $Z^1(B, \mathbb{F}_3^g/K)$, with no quotient-automorphism factor.

This cocycle count has an elementary inverse. On a nontrivial sign component $V_i = \mathbb{F}_3^{a_i}/K_i$, a cocycle z vanishes on $\ker \chi_i$: there it is a homomorphism, and each value is annihilated by both a power of 2 and 3. Choose an actual $s_i \in B$ with $\chi_i(s_i) = -1$. The cocycle identity then gives

$$z(b) = \chi_i(b)z(s_i) - z(s_i) \quad (b \in B).$$

Thus evaluation at s_i is inverse to the principal-cocycle map, since $-1 - 1 = 1$ in $\mathbb{F}_3$. The distinct-sign projections descend to the original quotient; summing their projected evaluations gives the inverse for the whole quotient. Consequently the fibre has exactly $3^{g - \dim K} = \prod_i 3^{a_i - \dim K_i}$ elements, and the literal lift count is $\prod_i L_3(a_i)$.

The original marker weight is $1/(6^g g! f!)$. Assigning the displayed marker coordinates to the q selected pairs contributes $g!/\prod_i a_i!$. A selected pair already has action weight $1/2$, so replacing the q pairs contributes 2^q . These give the coefficient in (28.1); the change of labelled support gives $n!/m!$; here closure under relabelling is essential. An ordering of the selected pairs only indexes the summation variables a_i ; it creates no $q!$ factor. Projection recovers all the displayed data, proving equality. $\square$

Proposition 28.2 (A presentation bound for collapsed pairs). *Under the hypotheses of Proposition 28.1, write $m = 2M$ and suppose $0 \leq q \leq M$. The same number of groups is at most*

$$\frac{n!}{m!} \frac{2^q H_{g,q}}{6^g f!} (M)_q |\mathcal{Y}_m|, \quad (M)_q = \frac{M!}{(M-q)!}.$$

This bound also follows directly by covering the groups with displayed collapsed presentations, without identifying equivalent presentations.

Proof. For each original sign image, enumerate its q distinct signs and assign each of the g marker coordinates to its sign. These presentations cover the original family; counting all enumerations gives an upper bound. For a fixed multiplicity profile, the assignments contribute $g!/\prod_i a_i!$ and the literal extension fibres contribute $\prod_i L_3(a_i)$. Thus their complete contribution is $g! H_{g,q}$.

Fix an exterior orbit profile with p existing pair orbits. Merge the q displayed collapsed pairs with those p pairs, and retain the condition that the resulting subgroup belongs to $\mathcal{Y}_m$. Dropping distinctness and any conditions referring to the distinguished q positions only enlarges this model family; its nonnegative presentation weights are kept unchanged. Relabelling invariance allows the ordinary full-profile labelling formula to be applied to it. The original pair divisor is $2^p p!$, whereas the merged pair divisor is $2^{p+q}(p+q)!$. Their ratio is

$$2^q \frac{(p+q)!}{p!} = 2^q (p+q)_q \leq 2^q (M)_q.$$

Every other exterior normalizer and occurrence factorial is unchanged. Together with the original marker divisor $6^g g!$, the fixed-point divisor $f!$, and the labelled-support ratio $n!/m!$, this gives the claimed bound. Summing over exterior profiles counts each eligible collapsed profile at most once for the fixed value of q . $\square$

Theorem 28.3 (Complete noncanonical marker row). *Write $n = 2N + \varepsilon$. For even $m = 2M < n$, define*

$$K_{\text{mark}}(n, m) = \frac{c_M G_M}{p_n G_N} \sum_{g,f,q} \frac{2^q H_{g,q}}{6^g f!} \binom{M}{q}, \quad (28.2)$$

where the sum is over nonnegative integers satisfying

$$n = m + 3g + f - 2q, \quad \begin{cases} q = 0, & g = 0, \\ 1 \leq q \leq \min(g, M), & g > 0, \end{cases} \quad D = \frac{g + f - \varepsilon}{2} \geq 1.$$

For all sufficiently large N , at both parities,

$$\sum_{\substack{m < n \\ m \text{ even}}} K_{\text{mark}}(n, m) \leq 2^{-N/10}. \quad (28.3)$$

For collections $\mathcal{Y}_m$ closed under conjugation by S_m , the complete normalized count of the corresponding marker groups whose collapsed group lies in $\mathcal{Y}_m$ is at most $\sum_m K_{\text{mark}}(n, m) |\mathcal{Y}_m| / L_m$. The same conclusions hold for the larger kernel $K_{\text{mark}}^+(n, m)$ obtained by replacing $\binom{M}{q}$ with $(M)_q$.

Proof. The counting comparison for K_{mark} follows from Proposition 28.1 and $t_q(Y) \leq \binom{M}{q}$. In particular it does not assume boundedness of any normalized subgroup count.

Put $\lambda = \log_2 3$ and $k = g - q$. Counting ordered bases bounds the number of d -dimensional subspaces of $\mathbb{F}_3^a$ by $3^{d(a-d+1)}$: there are at most 3^{ad} ordered independent d -tuples, and each such subspace has at least $3^{d(d-1)}$ ordered bases. Dropping the full-coordinate condition therefore gives

$$L_3(a) \leq (a+1)3^{a^2/4+a}.$$

For $a \geq 2$, use $a+1 \leq 3^{a-1}$ to obtain

$$L_3(a) \leq 3^{(a-1)^2/4+15(a-1)/4} \leq 2^{\lambda(a-1)^2/4+6(a-1)},$$

where $\lambda < 8/5$. The same bound holds at $a = 1$ because $L_3(1) = 1$ exactly. Consequently

$$H_{g,q} \leq 2^{\lambda k^2/4+6k+g}.$$

Here $\sum_i (a_i - 1)^2 \leq k^2$, and the number of compositions of g is at most 2^g ; dropping $\prod_i a_i!$ is harmless. For every term of (28.2),

$$M = N - D - k, \quad 0 \leq k \leq \min(2D, N - D), \quad g \leq 2D + 1.$$

The uniform Gaussian estimates leave quadratic exponent

$$Q = -\frac{N(D+k)}{2} + \frac{(D+k)^2}{4} + \frac{\lambda k^2}{4}.$$

We claim $Q \leq -13ND/60$. Since $\lambda < 8/5$,

$$60Q + 13ND \leq E := 15D^2 + 30Dk + 39k^2 - 17ND - 30Nk.$$

If $3D \leq N$, factor this polynomial as

$$E = -(17D + 30k)(N - 3D) - (2D - k)(39k + 18D) \leq 0.$$

If $3D \geq N$, use instead

$$E = -(3D - N)(9N - 8D) - (N - D - k)(39k + 9(N - D)) \leq 0.$$

All displayed factors are nonnegative under the stated constraints, including the endpoint $M = 0$.

The coefficient recurrence gives $c_M/p_n \leq (2N)^{D+k}$. Also $\binom{M}{q} \leq (M)_q \leq M^q \leq N^q$, $q \leq 3D$, and $6k + g \leq 15D$. The omitted factor $2^q/(6^g f!)$ is at most one. For fixed D there are at most $(2D+2)^2$ choices of g, f, q . The complete fixed- D logarithm is therefore at most $-13ND/60 + O(D \log(N+2))$, with one absolute implied constant. Eventually the logarithmic error is at most $ND/60$, uniformly in D , so the fixed- D logarithm is at most $-ND/5$. There are at most N positive defects, and therefore the complete sum is at most $N2^{-N/5} \leq 2^{-N/10}$ for all sufficiently large N . The same proof therefore bounds K_{mark}^+ ; its counting comparison follows directly from Proposition 28.2. In the $D = 0$ states used below, q is zero or one, so the falling factorial and binomial factors coincide. $\square$

Corollary 28.4 (Pure markers). *The normalized mass of all subgroups whose even orbits are critical and whose marker profile has $D \geq 1$ is $O(2^{-N/10})$. This sum includes all actual critical attachments and both parities.*

Proof. Collapse yields only critical binary orbits. Theorem 1.1 bounds their complete normalized count uniformly, including all finite initial degrees. Apply Theorem 28.3 to this conjugation-invariant family. This argument counts already assembled groups, so it also includes non-Frattini central attachments. $\square$

Corollary 28.5 (One typed binary-error transfer). *Suppose a family consists of fixed points, natural S_3 orbits, and binary orbits, at least one of which is noncritical. Its entire marker contribution is bounded by*

$$\sum_{\substack{m < n \\ m \text{ even}}} H_E(n, m) E_{m/2}, \quad H_E(n, m) = K_{\text{mark}}(n, m) + \mathbf{1}_{n=2N+1}(u_N \mathbf{1}_{m=n-1} + v_N \mathbf{1}_{m=n-3}),$$

with u_N, v_N from (27.3). There is no scalar remainder.

Proof. The noncritical binary orbit is unchanged by marker collapse, so the noncanonical part targets the complete conjugation-invariant family defining E . Any additional restrictions only decrease the original count. The only $D = 0$ marker states are no marker in even degree, one fixed point, and one S_3 in odd degree. The even state belongs to the separately counted complete binary family. The exact odd contributions are $\alpha_N E_N$ and $\beta_N p E_N$ by (28.1), and Proposition 27.2 gives the asserted two targets. Neither four-pair fusion nor duplicate collapse is required to preserve any earlier exclusion; they preserve a noncritical actual orbit, which is the defining condition of the complete target E . $\square$

For the final boundedness-first assembly we use a coarser but more compositional form of the same comparison. When $D \geq 1$, take $\mathcal{Y}_m$ to be the collection of all subgroups, so the marker kernel points directly to A_m . Only the zero-defect states retain the current and predecessor binary errors. Substituting their native forward recurrences then produces one strictly forward ordinary row.

29 The first split one-triple application

We consider subgroups with exactly one actual regular C_3 orbit and no actual natural A_4 orbit. On its complement of size $m = n - 3$, an external graph is determined by a literal nonzero character $\chi : K \rightarrow C_3$. Fix a generator of the displayed C_3 to specify its orientation. For split characters, let $\Theta(K)$ be the set whose complete graphs survive any preceding disjoint exclusions. The exact surviving character weight is

$$\sigma_3^{\text{surv}}(K) = |\Theta(K)| = \sum_{\substack{\chi \neq 0 \\ \chi \text{ split}}} \mathbf{1}_{\{\text{graph}(\chi) \text{ survives}\}}, \quad 0 \leq \sigma_3^{\text{surv}}(K) \leq 3^{d_3(K)} - 3^{h_3(K)}. \quad (29.1)$$

In particular the inequality need not be equality after exclusions. On a fixed labelled complement the surviving mass is precisely

$$\frac{n!}{6m!} \sum_{K \leq S_m} \sigma_3^{\text{surv}}(K). \quad (29.2)$$

The sum restricts to residuals without a regular C_3 or natural A_4 orbit. The factor six is the original action normalizer, and there is one selected triple because the actual orbit is unique.

Proposition 29.1 (Low one-triple cone). *The split graphs with $20d_3(K) \leq 3m$ have normalized mass at most*

$$K_{\text{lo}}(n, m) \frac{s_m}{L_m}, \quad K_{\text{lo}}(n, m) = \frac{1}{6} \Delta(n, m) 3^{3m/20} \leq 2^{-\eta m + O(\log(n+2))}, \quad \eta = \frac{1}{4} - \frac{3 \log_2 3}{20} > 0.$$

Proof. Apply (29.1) to each residual before enlarging its number to s_m . The labelled factorials cancel exactly as in (29.2). Here is an explicit bound covering every fixed deletion width w :

$$\Delta(b+w, b) \leq \phi^{-2} 2^{(w+1)/4} (b+w+1)^{w+2} 2^{-\lfloor w/2 \rfloor b/4}, \quad \phi = \prod_{j \geq 1} (1 - 2^{-j}). \quad (29.3)$$

Indeed, the two guarded parity coefficients at rank k both lie between A_k and $(k+1)A_k$, since $A_{k-1} \leq 2kA_k$. The rank shift $t = \lfloor (b+w)/2 \rfloor - \lfloor b/2 \rfloor$ lies between $\lfloor w/2 \rfloor$ and w . Apply the coefficient shift bound $A_k \leq (2(k+t))^t A_{k+t}$ and the Gaussian ratio bound; their polynomial factors are at most $(b+w+1)^{w+2}$. The Gaussian exponent is at most $(w+1)/4 - \lfloor w/2 \rfloor b/4$, proving the display also at rank zero. For $w = 3$ this gives the required loss $m/4$. Finally $3^5 < 2^8$ gives $3 \log_2(3)/20 < 6/25 < 1/4$, so the low-cone kernel is also $O(2^{-m/200})$ after absorbing its fixed polynomial factor. $\square$

We next prove the earlier orbit estimates actually needed for the high cone. Only finitely many physical degrees occur in this application; no growing action menu is needed. Put $\gamma = (\log_2 3)/3$ and

$$Z_J(U) = \sum_{N \triangleleft U} |\text{Epi}(J, U/N)|, \quad a(U) = |N_{S_w}(U)|, \quad J \leq S_b.$$

All target normal subgroups in this sum are literal. Fixing each target quotient map is the asymmetric Goursat convention, so there is no additional automorphism divisor.

We use one fixed-source reduction throughout these rows. Suppose an actual target extension has elementary p -kernel module M and top B , and that M embeds equivariantly in one copy of $\mathbb{F}_p[B]$. For an onto top map $\beta : J \twoheadrightarrow B$, evaluation at the identity coefficient injects the equivariant homomorphisms from $\ker \beta$ to M into $\text{Hom}(\ker \beta, \mathbb{F}_p)$. Indeed, equivariance and the surjectivity of β recover every regular coordinate from that one coefficient. Consequently every surviving lift fibre has size at most

$$|M| |H^1(B, M)| |\text{Hom}(\ker \beta, \mathbb{F}_p)|.$$

This remains valid for empty and nonsplit fibres. We first sum these character masses over the actual top epimorphisms, and only then over every literal original normal axis. The same complete source J is retained in all terms; the fixed module and cohomology factors are absorbed into D_U . Thus the row-specific numerical work below concerns only the top maps and the character ranks of their actual kernels.

Lemma 29.2 (Finite earlier orbit fibres). *For the following actual actions, $Z_J(U)/a(U)$ is at most $D_U(1+b)2^{\alpha_U b}$, with a fixed constant D_U :*

action	w	α_U
6T1, 6T4, 6T5, 6T6	6	γ
12T20, 12T85, 12T164	12	2γ
12T194	12	$16/15$
12T228, 12T229, 12T265	12	$1/2 + 5 \log_2(3)/9$
transitive 3-group	9	$5 \log_2(3)/9$
transitive 3-group	27	$5 \log_2(3)/3$

The degree-nine row and the three final degree-twelve actions require J to have exactly one regular C_3 orbit and no natural A_4 orbit. Every other row permits arbitrary J .

Proof. We give the fibre arguments, including the shared source budgets. We use the internally proved bounds

$$d_2(F) \leq b/2, \quad d_3(F) \leq b/3$$

for every literal permutation subgroup $F \leq S_b$. They follow by restriction to an actual Sylow subgroup and degree induction on faithful p -group actions, and hence apply unchanged to literal kernels of the top maps below. For the few whole abelian target terms we separately use the abelianization-order bound $|J/J'| \leq 3^{b/3}$ from [KP89]. Constants depending on a fixed target, including its finite normal menu, are absorbed only after its full quotient sum has been written.

For $6T1 = C_6$, the full fibre is $|\text{Hom}(J, C_6)|$. For $6T5 = C_3 \wr C_2 \cong C_3 \times S_3$, its normal quotients are $1, C_2, C_3, C_6, S_3, C_3 \times S_3$. Set $F = J'J^2$ and $E = J/F$. The semisimple $\mathbb{F}_3 E$ -module $H_1(F; \mathbb{F}_3)$ has trivial multiplicity $d = d_3(J)$ and sign multiplicities a_θ , with $d + \sum a_\theta \leq b/3$. Coprime inflation–restriction gives exactly

$$|\text{Epi}(J, S_3)| = 3 \sum_{\theta \neq 0} (3^{a_\theta} - 1), \quad |\text{Epi}(J, C_3 \times S_3)| = (3^d - 1) 3 \sum_{\theta \neq 0} (3^{a_\theta} - 1).$$

The product is onto because C_3 and S_3 have no common nontrivial quotient. Since $\sum (3^{a_\theta} - 1) \leq 3 \sum a_\theta - 1$, each expression is at most $3^{b/3+1}$. The central and twisted characters have used one module budget.

The actions $6T4, 6T6$ are respectively A_4 in degree six and $C_2 \times A_4$ in degree six. Their binary bases are cyclic $\mathbb{F}_2[C_3]$ -modules. Likewise $12T20, 12T85, 12T164$ have cyclic binary bases and elementary tops C_3^2 , with the top action on the base allowed to have a kernel. Here is the requisite general estimate for $M \rtimes C_3^r$, $r = 1, 2$, with cyclic binary M . Each quotient is $V \rtimes C_3^h$, $h \leq r$, with V a multiplicity-one semisimple module. Use the same source $F = J'J^3$, $E = J/F$, $D = H_1(F; \mathbb{F}_2)$ for every top map. Write the trivial multiplicity of D as r_0 and its nontrivial simple multiplicities as a_α . Each nontrivial type has dimension two, and

$$r_0 + 2 \sum_{\alpha} a_\alpha \leq b/2.$$

For a fixed onto top map $\theta : E \rightarrow C_3^h$, surjective restrictions to F number $(2^{r_0} - 1)^e \prod_i (4^{a_{\alpha_i}(\theta)} - 1)$, where $e \in \{0, 1\}$. If a nontrivial target constituent is retained, fix one of its two concrete ternary characters. Pulling it back to a specified source type permits at most $2 \cdot 3^{(h-1)d_3(J)}$ top maps. Bound the remaining factors using the one source budget and sum the retained type. The result is at most

$$(b/2) 3^{(h-1)d_3(J)} 2^{b/2}.$$

For each restriction and top map, the lift torsor has at most $|M|$ elements, by coprime cohomology on E . Abelian quotients are bounded by $|J/J'|^r$ (or by one such factor when $h = 0$). Thus all quotients obey the claimed $(1+b)2^{r\gamma b}$ bound. This argument treats nonfaithful top actions as well as faithful ones.

For $12T194 = C_3 \wr A_4$, use its full ternary block kernel. Every quotient is an extension of a quotient S of A_4 by a module subquotient X of the natural permutation module $\mathbb{F}_3^4$. Every such X embeds in $\mathbb{F}_3[S]$. To see this, first restrict to the normal V_4 . Over a splitting field its trivial character gives one trivial C_3 -module; its three nontrivial characters form one orbit and give a simple matrix-algebra component. Subquotients of the first embed in the regular C_3 block, and the second component is semisimple. The whole permutation module embeds in $\mathbb{F}_3[A_4]$ by coset sums. If the action factors through S , the image lies in the fixed space of the kernel, which is the inflated regular $\mathbb{F}_3[S]$ -module. For a concrete top map $J \rightarrow S$ with kernel F , the restriction

choices are therefore at most $3^{d_3(F)} \leq 3^{b/3}$. The compatible lift torsor has a fixed target-dependent size: after the restriction is fixed, differences descend to S .

The quotient menu itself has a shorter structural proof. If $q : A_4 \twoheadrightarrow S$ and $N = \ker q$, then $|N|$ divides 12. A normal subgroup of order two is central, whereas $Z(A_4) = 1$. For $|N| = 3, 4, 6$, the quotient has order 4, 3, 2, respectively, and is abelian. Thus

$$V_4 = [A_4, A_4] \leq N.$$

The orders three and six are impossible, and order four forces $N = V_4$. Hence S is A_4 , C_3 , or trivial. For the latter two targets the internal ternary character-rank bound gives the required top count. The full target has a shorter same-source argument. Put $D = [J, J]$, identify $[A_4, A_4]$ with V_4 , and fix one nonzero binary character $\eta : V_4 \rightarrow C_2$. Label an epimorphism $f : J \twoheadrightarrow A_4$ by

$$\chi_f = \eta \circ f|_D \in \text{Hom}(D, C_2).$$

This label determines the literal kernel of f . Indeed, if $\chi_f = \chi_g$ and $x \in \ker f$, then surjectivity of g gives, for every $a \in A_4$,

$$\eta([g(x), a]) = \eta(g([x, y])) = \eta(f([x, y])) = 0 \quad (a = g(y)).$$

The action of $A_4/V_4 \cong C_3$ permutes the three nonzero elements and the three nonzero characters of V_4 freely. Consequently $\eta([v, a]) = 0$ for every $a \in A_4$ forces $v = 1$. Thus $x \in \ker g$, and symmetry gives $\ker f = \ker g$.

Every fixed kernel supports exactly $|\text{Aut}(A_4)|$ literal epimorphisms. Applying the internal binary half-rank theorem to the same derived subgroup $D \leq S_b$ therefore gives

$$|\text{Epi}(J, A_4)| \leq |\text{Aut}(A_4)| |\text{Hom}(D, C_2)| \leq |\text{Aut}(A_4)| 2^{\lfloor b/2 \rfloor}.$$

Since $1/2 < 8/15$, this embeds directly in the required quotient-top exponent, without a polynomial loss. For a fixed top map, the literal kernel contributes at most $3^{d_3(F)} \leq 3^{b/3} \leq 2^{8b/15}$. Summing the three actual quotient targets therefore gives $D2^{16b/15}$, as recorded in the table.

For degree-nine 3-groups, the target chart has two forms: an elementary ternary group of rank at most two, or an extension of C_3 by a module embedding in $\mathbb{F}_3[C_3]$. The source pattern is aggregated before any target is counted. If o_0 is its unique regular C_3 orbit, then for every literal orbit image and every original normal axis the local bounds have the form

$$9\delta_o \leq 2|o| + 3\mathbf{1}_{\{o=o_0\}}.$$

The faithful orbit filtration sums these inequalities on the same complete source J . In the present one-regular- C_3 , no-natural- A_4 case, (9.5) reads

$$9d_3(J) \leq 2b + 3.$$

For an elementary target, the exact homomorphism count bounds onto maps by the corresponding power of $3^{d_3(J)}$. In the second case, summing first over the actual maps $J \twoheadrightarrow C_3$ and their literal kernels, then retaining the fixed module and H^1 factors, gives uniformly

$$|\text{Epi}(J, Q)| \leq 3^{d_3(J)+b/3+6}. \quad (29.4)$$

The constant 3^6 is target-local and does not affect the row exponent. Combining this with (9.5) gives the ternary exponent $5 \log_2(3)/9$; when the descended extension has a binary kernel, the internally proved literal-kernel bound $d_2(\ker \beta) \leq b/2$ gives the mixed exponent $1/2 + 5 \log_2(3)/9$.

For the three remaining degree-twelve actions, the actual action has three blocks of four points, binary translation base $W \leq V_4^3$, and a 3-group top P acting transitively on the nine nonzero local translations. The full module V_4^3 is cyclic: a nonzero local translation generates its local plane under the block stabilizer, and its top translates generate all three planes. Every invariant submodule is cyclic by Maschke's theorem. Consequently, for every normal quotient Q , its binary kernel M is cyclic for its top quotient S of P . Put $F = O^3(J)$ and $E = J/F$. For a fixed map $J \rightarrow S$, the pulled-back M embeds in the regular $\mathbb{F}_2[E]$ -module. It follows that

$$|\mathrm{Hom}_E(H_1(F; \mathbb{F}_2), M)| \leq 2^{b/2}.$$

Coprime lift torsors on E cost at most $|M| \leq 2^6$. Apply (29.4) to the same source J and the actual nine-point action of P , before enlarging the pattern class of sources. This gives the exponent $1/2 + 5 \log_2(3)/9$.

Finally we prove the degree-27 row without any classification of its 3-groups. More generally, for a transitive 3-group of degree $w = 3^k$, every quotient satisfies

$$|\mathrm{Epi}(J, Q)| \leq D_w 3^{(w+3)b/18}, \quad (29.5)$$

where D_w is independent of J and U . Choose a central semiregular C_3 in U ; its blocks have full elementary ternary kernel $V \leq \mathbb{F}_3^s$, $s = w/3$, and transitive 3-group top T . A central semiregular C_3 of T makes this permutation module free of rank $s/3$ on restriction. Every subquotient M therefore has socle dimension at most $s/3$. If its action factors through S , it embeds in at most $s/3$ copies of the regular $\mathbb{F}_3[S]$ -module: the group algebra is a self-injective algebra with its unique simple module trivial. Hence, over a fixed map $J \rightarrow S$, restrictions from its literal kernel number at most $3^{sb/9}$. After fixing a restriction, lift differences descend to S and have a bounded target-dependent cost. Induction on k gives (29.5), since $(s+3)/18 + s/9 = (w+3)/18$. If $S = 1$, use the simultaneous head bound $d_3(U) \leq (w+3)/6$, obtained from the same block filtration. The base $w = 3$ is the elementary rank bound. For $w \leq 27$ the normal menus and target sizes are finite, so their uniform constants are finite. At $w = 27$ the coefficient is $5/3$, as required. $\square$

Lemma 29.3 (Actual action membership). *Every exceptional action in Theorem 9.5, other than natural C_3, A_4 , is accepted by Lemma 29.2. The action descriptions used there are certificates on the original permutation sets.*

Proof. For the degree-six actions the witnesses are, in order, regular C_6 , the degree-six action of A_4 , $C_3 \wr C_2$, and $C_2 \times A_4$ with its cyclic three-dimensional binary base. These witnesses also show that the first and third have normal Sylow 3-subgroup of index two, the second is an irreducible coprime affine action outside degree four, and the fourth has cyclic binary module and elementary ternary top. Thus they belong to the earlier physical action domains.

The degree-six ownership statement also has a structural proof. Choose an actual minimal block system. Highness rules out primitivity, and the degree product leaves either fibres of size three over two blocks or fibres of size two over three blocks. In the first case the faithful top has order two. A nontrivial sign on a three-point fibre annihilates the complete relative ternary head by the signed-coordinate argument, contradicting highness. Hence all fibre signs are trivial, the literal block kernel is a normal 3-group, and its index is two. Thus the whole 3×2 branch enters the first earlier domain without finite action recognition.

In the 2×3 branch, highness forces the faithful three-block top to have order three. The literal block kernel is its actual invariant submodule of the binary permutation module $\mathbb{F}_2^3$. Every invariant submodule of this local C_3 -module is generated by the orbit of one vector. Schur–Zassenhaus

supplies an actual complement to the binary kernel, and restriction identifies that complement with the three-block top. The additive orbit words then translate through the faithful pair-coordinate chart into products of actual conjugates of the distinguished kernel element. This is precisely the cyclic binary-module earlier-owner witness. Consequently the entire degree-six high branch is accepted structurally. The $6T4$ and $6T6$ certificates remain independent audits of the two familiar actions, but the $c = 1$ ownership implication no longer uses the degree-six catalogue.

Degree twelve also admits a structural reduction before any finite recognition is used. For an internally selected minimal block system, write r for the fibre size, s for the number of blocks, v for the ternary weight of a chosen primitive-fibre chief series, and t for the relative ternary head of the original normal top image. The high pair has relative rank two and

$$2 \leq vB(s) + t, \quad v \leq \lfloor r/3 \rfloor, \quad t \leq C(s), \quad rs = 12.$$

The 2×6 orientation has $v = 0$ and $t \leq 1$, while the 6×2 orientation has $v \leq 1$ and $t = 0$; both are impossible. Thus every high degree-twelve pair has a 3×4 or 4×3 minimal block system. In either case $B(s) = 1$, and equality is forced throughout:

$$\delta_A(N) = 2, \quad v = 1, \quad t = 1.$$

The rank-one four-block top is the natural A_4 action, while the faithful transitive rank-one three-block top has order three. Hence these are literal prime-base and V_4 -block top geometries, rather than names assigned from the finite catalogue. Accordingly the remaining bounded recognition problem concerns only the two saturated prime-base and V_4 -block geometries. The complete degree-twelve census below audits that structural fork and identifies the owner realized in each branch.

The prime-base branch retains more than the numerical equality above. A nontrivial sign on any three-point fibre would bound the original relative head by the top relative head. Since their dimensions are two and one, respectively, every fibre sign is trivial and the literal block kernel is a 3-group. Applying the exact normal-chain restriction formula to

$$N \cap \ker(A \longrightarrow A_4) \leq N$$

shows that the retained restriction image on this physical intersection has dimension exactly one. In particular it contains an actual nonzero whole-ambient invariant character; this is the marking datum needed below, not merely a positive chief weight.

For reference, the natural A_4 -module $\mathbb{F}_3^4$ has exactly four invariant submodules:

$$0, \quad \langle (1, 1, 1, 1) \rangle, \quad \{x : \sum_i x_i = 0\}, \quad \mathbb{F}_3^4.$$

The augmentation module in the third position has full projection on every coordinate, so coordinate-surjectivity alone does not remove it. It has no nonzero A_4 -invariant linear functional, whereas the diagonal and full modules do. Thus, once the literal fibre-coordinate action is identified with this natural module, the retained character forces the block kernel to be diagonal or full. This identification is now intrinsic. Choose one actual triple-fibre chart and transport it to the other fibres by ambient group elements. The vanished fibre signs make every transition even, so the resulting coordinate map injects the original block kernel A_4 -equivariantly into $\mathbb{F}_3^4$. Its image and the image of every physical intersection with an original normal subgroup are invariant. The retained character therefore applies to the actual coordinate image, rather than to a substituted product module.

The same four-submodule classification removes a second, potentially case-by-case step. If $B, C \leq \mathbb{F}_3^4$ are A_4 -invariant, the section $B/(B \cap C)$ embeds A_4 -equivariantly back into $\mathbb{F}_3^4$: the required maps are the identity, the diagonal and augmentation projections, or zero. Suppose its action factors through an actual quotient $A_4 \twoheadrightarrow Q$. Evaluating an embedded vector at one coordinate and translating that functional through Q gives an injective matrix-coefficient map

$$B/(B \cap C) \hookrightarrow \mathbb{F}_3[Q].$$

Thus every such descended section lies in one copy of the regular module of its actual quotient; no classification of the quotients of A_4 is needed. The arbitrary-normal quotient construction retains the other presentation of the same physical section. From the checked original kernel chart $\pi : G \twoheadrightarrow A_4$ and any $N \triangleleft G$, it constructs the literal extension

$$G/N \longrightarrow G/(\ker \pi N)$$

with kernel $\ker \pi / (N \cap \ker \pi)$, its quotient module and the original conjugation action. The same construction identifies its module with the invariant coordinate section above, constructs the onto top map $A_4 \twoheadrightarrow G/(\ker \pi N)$, and proves that the coordinate action factors through this actual quotient. The matrix-coefficient map therefore gives an injective equivariant map from the physical kernel of G/N into $\mathbb{F}_3[G/(\ker \pi N)]$. This uses neither a splitting nor the assumption $N \leq \ker \pi$. Thus the remaining obligation in this orientation is the fixed-source owner sum, rather than quotient-module identification.

There is a parallel intrinsic reduction in the four-point-fibre branch. The physical intersection $N \cap \ker(A \rightarrow C_3)$ has relative ternary head one. An odd local image would propagate to every fibre by original ambient conjugation. Every normal subgroup of such an odd transitive four-point image has zero relative ternary head; the subdirect relative-head inequality would therefore force the displayed intersection head to vanish. Hence all literal fibre images are even. Set

$$W = \{g \in \ker(A \rightarrow C_3) : g^2 = 1\}.$$

Then $W \triangleleft A$ is an elementary binary group which embeds in the three actual Klein-four coordinates, so $|W| \leq 64$. The quotient A/W is a 3-group, and Schur–Zassenhaus gives an actual ternary complement. Moreover each block-kernel image is all of A_4 : the positive intersection head supplies one full local image, and ambient conjugation transports fullness to the other fibres. Cubing an arbitrary lift of a local Klein translation kills every ternary coordinate and leaves the prescribed translation, so each literal map $W \rightarrow V_4$ is onto. This statement deliberately allows the proper submodules occurring in the degree-twelve rows: it does not replace W by all of V_4^3 . Put

$$\mathcal{T} = X \times (V_4 \setminus \{1\}),$$

where X is the set of three actual four-point fibres. Ambient conjugation, written in arbitrary local point charts, defines an action on the nine elements of $\mathcal{T}$. Its kernel is exactly W : fixing all local translations forces the local permutation into the Klein group, and then faithfulness on the original points identifies the intrinsic core. Full local A_4 images make this action transitive. Hence the whole 3-group quotient A/W , rather than only the three-block top, has a faithful transitive degree-nine action. The elementary group W also supplies its intrinsic binary conjugation module. For every original $N \triangleleft A$, the arbitrary-normal construction retains the literal extension

$$A/N \longrightarrow A/(WN), \quad \ker = W/(W \cap N),$$

with the original representatives and conjugation action; the descended top is still a 3-group. No containment $N \leq W$, complement, full-product, or cyclic-module assumption is used. The remaining step in this orientation is its fixed-source top-character estimate.

The bounded degree-twelve certificate constructs $W = O_2(U)$ and a Sylow 3-complement for 12T20, 85, 164, 228, 229, 265. It verifies that W has three actual orbits of size four and local image V_4 . Restricting translations to the blocks constructs a literal ambient V_4^3 normalized by U . The complement acts faithfully and transitively on its nine nonidentity local translations, and transitively on the three blocks. Its order, and the base order, are

U	12T20	12T85	12T164	12T228	12T229	12T265
$ W $	4	16	64	64	64	64
$ P $	9	9	9	27	27	81.

In the first three rows the complement is C_3^2 , and an exhibited base vector has orbit spanning W ; these are exactly the cyclic module witnesses used in the proof. In the last three rows the actual nine-point complement is, respectively, 9T6, 9T7, 9T17; the proper/full binary-base domain applies. In 12T20, 12T194, the normal ternary subgroup instead has four actual orbits of size three, local image C_3 , and full natural A_4 block action. The standard transversal embedding puts the original group inside $C_3 \wr A_4$; the base is diagonal in the first case and full in the second. Thus 12T20 also belongs to the earlier prime-base domain, and 12T194 belongs to it directly.

These finite statements are part of the bounded action certificate: they verify actual blocks, local images, full projection and cyclic generators, rather than deciding membership from order or a catalogue label. They are separate from the exhaustive normal-pair computation which produced the high list. The structural three-by-three block argument in Theorem 9.5 places every degree-nine rank-two pair in the transitive 3-group owner, and its iterated reversible block-sign dichotomy places the degree-27 row in that same owner. Their counting proof therefore applies uniformly to every actual transitive 3-group in those degrees; no further finite action classification is introduced here. $\square$

Corollary 29.4 (Ternary owner or capacity). *Let (A, N) be a transitive normal pair of degree w . Then at least one of the following holds:*

1. *A is the natural C_3 or natural A_4 action;*
2. *the original action of A is accepted by an earlier physical family in Lemma 29.2;*
- 3.

$$20\delta_A(N) \leq 3w.$$

The owner in the second alternative is attached to the original permutation action, so deleting that actual orbit retains the complete complementary subgroup.

Proof. If the capacity inequality fails, Theorem 9.5 puts the actual action on its necessary high list. The two natural actions give the first alternative, and Lemma 29.3 gives the second for every other entry. The structural degree-six and degree-twelve arguments in that lemma identify the owner intrinsically; the finite certificates only audit the resulting action charts. $\square$

Theorem 29.5 (First split one-triple estimate). *The family of split one-triple graphs with exactly one actual regular C_3 orbit and no natural A_4 orbit admits a positive forward bound against complete lower-degree subgroup counts whose row tends to zero exponentially. After its low cone and the actual orbit families in Lemma 29.2 have been removed, its high cone is empty.*

Proof. Apply Proposition 29.1. For every remaining graph with an eligible actual orbit, point one such orbit of degree w . Its complete complement $J \leq S_b$, $b = n - w$, includes the external triple and every other orbit and relation. The exact pointed comparison is

$$\sum_H k(H) = \sum_U \frac{n!}{b!a(U)} \sum_J Z_J^{\text{surv}}(U), \quad 0 \leq Z_J^{\text{surv}}(U) \leq Z_J(U).$$

Here the surviving fibre retains the preceding exclusions on the complete pointed graph. The unrestricted fibre is used only for the upper bound. If required, the pattern is retained on each J before using its rank bound. Deleting any of the indicated nonmarker actions preserves the unique- C_3 , no-natural- A_4 pattern. Only after the fibre has been bounded do we enlarge the number of such J to s_b . Normalize once, obtaining terms

$$D_U(1+b)\Delta(n,b)2^{\alpha_U b} \frac{s_b}{L_b}.$$

For even w the Gaussian coefficient of b is $w/8$; for odd w its safe value in both parities is $(w-1)/8$. These are strictly larger than every corresponding α_U in the table. For a check using only rational numbers, $3^5 < 2^8$ bounds its five exponents by $8/15, 16/15, 25/18, 8/9, 8/3$, respectively. Each resulting gap is at least $1/9$. Equation (29.3) bounds the coefficient cost for each fixed deletion by a polynomial in n . There are only finitely many actions in degrees 6, 9, 12, 27, so the whole forward row is $O(2^{-\varepsilon n})$ for some $\varepsilon > 0$. To form a disjoint physical partition, assign a graph to the first eligible family in a fixed order. Summing its intersections before applying the full pointed upper bound charges each family once. Multiple eligible orbits give only the positive pointing multiplicity.

Suppose now that a high graph survives. Its residual K has $20d_3(K) > 3m$. Because projection of the complete graph onto K is an isomorphism, the actual nonexternal orbit actions of the graph are exactly those of K . Apply Corollary 29.4 to each original orbit pair. The retained pattern excludes its first alternative, and the preceding physical removals exclude its second. Thus every orbit pair satisfies the capacity inequality. Equation (9.2) now gives

$$20d_3(K) \leq \sum_i 20\delta_{A_i}(N_i) \leq \sum_i 3 \deg A_i = 3m,$$

a contradiction. Fixed points have head zero and keep their degree in this inequality. Thus every surviving character set $\Theta(K)$ in the high cone is empty. This is a statement about complete marked graphs; it introduces no additional numerical kernel or scalar term. $\square$

The two exact incidence tests. The preceding proof respects Proposition 27.3 on every disjoint piece. In particular, applying that proposition with $\chi \in \Theta(K)$ gives

$$\sum_K \sigma_3^{\text{surv}}(K) = \sum_{|x|=3} \sum_{\substack{N^x=N, \ N \cap \langle x \rangle=1 \\ \chi_{N,x} \in \Theta(N \langle x \rangle)}} \frac{1}{|\{y \in Nx : |y|=3\}|}.$$

The reciprocal is preserved. It is not harmless to drop it: for $K_0 = C_2^{3h} \rtimes \langle x \rangle$, with x cycling h triples of pair coordinates, and $K = K_0 \times A_5^2$ on $m = 6h + 10$ points, its indicated quotient coset contains exactly $4^h \cdot 21^2$ elements of order three. This already loses $m/3 - O(1)$ bits if the reciprocal is discarded.

Likewise the affine test on the same source $J = \mathbb{F}_4^r \rtimes C_3$ has, above the fixed top map to $\mathbb{F}_4^t \rtimes C_3$, exactly $4^{t(r+1)}$ cocycles and $4^t \prod_{i=0}^{t-1} (4^r - 4^i)$ onto lifts. A second mark stays on this same J . In

the proof above the external triple is part of J in every earlier fibre, including the degree-six affine fibre; it is never appended as an independent character maximum. The final zero-set argument makes all these joint high surviving incidences empty and therefore requires no factorization of their moments.

30 The complete macroscopic alternating packet

Write $A_4 = (\mathbb{F}_4, +) \rtimes C_3$ in its natural action on four points. Let $Z_{b,c,t}$ count subgroups in the fixed container $S_b \times C_3^c \times A_4^t$ full on the displayed C_3, A_4 coordinates and with residual projection having no natural C_3 or A_4 orbit. Put $n = b + 3c + 4t$.

Theorem 30.1 (All image and source parameters summed). *Uniformly for $b, c \geq 0$ and $t \geq 1$,*

$$\log Z_{b,c,t} \leq \frac{n^2}{16} - \frac{43t^2}{450} + O(b^{3/2} + t + \log(c+2)). \quad (30.1)$$

After the single original action factor $n!/(b!6^c c!24^t t!)$, the entire sector $t \geq n/1000$ has normalized mass at most $2^{-43n^2/(900 \cdot 10^6)}$ eventually.

Proof. Fix the literal residual group $J \leq S_b$. Put $F = J'J^3$, $E = J/F = \mathbb{F}_3^r$, $r = d_3(J)$, and $T = \mathbb{F}_3^t$. Conjugation makes $M = H_1(F, \mathbb{F}_2)$ a semisimple $\mathbb{F}_2 E$ -module. Since $\mathbb{F}_4$ contains all third roots of unity, its nontrivial simple modules have dimension two, are indexed by $h \in (E^* \setminus \{0\})/\{\pm 1\}$, and have endomorphism field $\mathbb{F}_4$. Write their multiplicities as a_h , and $A = \sum_h a_h$. The permutation section bound gives $A \leq d_2(F)/2 \leq b/4$; equation (9.5) gives $r \leq 2b/9$.

If $K \triangleleft J$ and J/K is elementary ternary, then $K/F = D$ for a unique subspace $D \leq E$. Odd-order homology gives naturally

$$H_1(K, \mathbb{F}_2) = M_D.$$

The D -coinvariants retain precisely those simple types whose characters vanish on D , with the same multiplicities. Distinct retained types do not merge: equality up to sign on E/D is already equality up to sign on E . This controls all image kernels from one fixed source module M .

An image $L \leq J \times T$ onto J contains $F \times 0$. Its reduction $\bar{L} \leq E \times T$ is therefore equivalent to its annihilator, the graph

$$\bar{L}^\perp = \{(\Gamma(w), w) : w \in C_T\}, \quad C_T \leq T^*, \quad \Gamma : C_T \rightarrow E^*.$$

This is a bijection: projection of $\bar{L}$ onto E is onto exactly when the annihilator meets $E^* \times 0$ trivially. Let $S = \{i : e_i \in C_T\}$, $s = |S|$, $u = t - s$, and $a = \dim C_T - s$. Fullness on packet coordinate i says $\Gamma(e_i) \neq 0$ whenever $i \in S$. The exact rank is

$$d_3(L) = r + u - a.$$

The coordinates in S are exactly those eligible for a residual source map; coordinate i has source type $[\Gamma(e_i)]$. No eligible and ineligible coordinate can have the same packet isotype: equality up to sign would put $e_i \pm e_j$ in $\ker \Gamma \leq C_T$, and hence put both in C_T .

We spell out the binary fibre above this same image. If $W \leq V^t$, $V = (\mathbb{F}_4, +)$, is its intersection with the packet kernel, write $Q = V^t/W$. Subgroups above L with intersection W are the graphs of literal cocycles in $Z^1(L, Q)$. For $K = L \cap J$, the odd quotient $L/K \leq T$ gives

$$H^1(L, Q) \cong \text{Hom}_{L/K}(H_1(K, \mathbb{F}_2), Q), \quad |B^1(L, Q)| = |Q|.$$

The second equality holds because all packet simple constituents are nontrivial and thus $Q^L = 0$. Coordinate fullness can only restrict this set. An isotype of packet multiplicity m and source multiplicity α consequently contributes at most

$$F_\alpha(m) = \sum_{x=0}^m \left[\begin{matrix} m \\ x \end{matrix} \right]_4 4^{(\alpha+1)x} \leq 2^{2\alpha m + m^2/2 + 4m}.$$

Indeed $\left[\begin{matrix} m \\ x \end{matrix} \right]_4 < 2 \cdot 4^{x(m-x)}$ and there are $m+1$ summands. The term $+1$ counts the literal coboundaries; it is not another source-module choice.

Fix C_T and extend its s coordinate axes to a basis. Values of Γ on the other a basis vectors contribute 3^{ra} . If the projective multiplicities of the nonzero axis values are m_h , the eligible fibre is at most $2^{2\sum_h a_h m_h + s^2/2 + 4s}$. Summing all its oriented axis values gives

$$2^{s^2/2 + 4s} \left(2 \sum_h 4^{a_h} \right)^s.$$

Integer convexity and the two physical rank bounds show

$$2 \sum_h 4^{a_h} \leq 2 \left(\frac{3^r - 1}{2} + 4^A - 1 \right) \leq 2^{b/2 + 2}.$$

The ineligible part has source multiplicity zero and contributes at most $2^{u^2/2 + 4u}$. Thus the whole fixed- C_T sum is at most

$$2^{bs/2 + (s^2 + u^2)/2 + (\log_2 3)ra + O(t)}.$$

For each exact axis set S , the number of possible C_T of dimension $s+a$ is at most $\left[\begin{matrix} u \\ a \end{matrix} \right]_3$: quotient by the contained coordinate space, and omit the requirement of no additional axes. There are $\binom{t}{s}$ choices of S . Defining

$$J_*(x, y) = \max_{0 \leq z \leq y} z(x + y - z) = \begin{cases} xy, & x \geq y, \\ (x + y)^2/4, & x \leq y, \end{cases}$$

and $\lambda = \log_2 3$, the all-image fibre over this J is at most

$$2^{\max_{s+u=t} \{bs/2 + (s^2 + u^2)/2 + \lambda J_*(r, u)\} + O(t)}. \quad (30.2)$$

This sums the image, source-column assignment, kernel subspace and cocycle on the same group.

The deficit in (30.2) from $bt/2 + t^2$ is at least $\delta_0 t^2$, with $\delta_0 = (2 - \lambda)/4$. For completeness, set $x = b/t$, $v = u/t$. The deficit divided by t^2 is at least

$$d(x, v) = 1/2 + v - v^2 + xv/2 - \lambda J_*(2x/9, v).$$

When $2x/9 \geq v$ this increases in x , so its minimum is at $x = 9v/2$, where it is $1/2 + v + (5/4 - \lambda)v^2 \geq 1/2$. On the other region it is concave in x , so check $x = 0$ and $x = 9v/2$. At zero it is $1/2 + v - (1 + \lambda/4)v^2$, a concave quadratic whose minimum on $[0, 1]$ is δ_0 .

Adjoin the c external regular ternary coordinates. The binary packet extension does not change $d_3(L)$, and $d_3(L) \leq P = 2b/9 + t$. Its one exact graph fibre is at most $2^{\lambda J_*(P, c) + O(\log(c+2))}$. Its physical increment in $n^2/16$ is $3bc/8 + 3tc/2 + 9c^2/16$. If $c \leq P$, the difference is

$$(3/8 - 2\lambda/9)bc + (3/2 - \lambda)tc + 9c^2/16 \geq -\frac{4}{9}(\lambda - 3/2)^2 t^2 > -t^2/225.$$

If $c \geq P$, the difference increases in c and at $c = P$ is $P[(1/2 - 2\lambda/9)b + (33/16 - \lambda)t] > 0$. Summing J uses the independent coarse bound $s_b \leq 2^{b^2/16 + O(b^{3/2})}$ exactly once. Since $\delta_0 - 1/225 > 43/450$, this proves (30.1).

Every actual group recovers the sets of its natural C_3 and A_4 orbits. Choosing these supports and actions contributes exactly $n!/(b!6^c c! 24^t t!)$. The residual J excludes those orbit types, so they are not selected a second time. Normalization by L_n costs only $O(n \log(n+2))$ beyond the $n^2/16$ term. For $t \geq n/1000$ the main loss dominates this and the $O(n^{3/2})$ error uniformly; the $O(n^2)$ possible (c, t) values are absorbed as well. This proves the complete scalar claim. $\square$

31 The small natural alternating packet

Let $\mathcal{U}$ be the class of allowed transitive actions consisting of natural C_3, A_4 and actions U for which every literal normal subgroup N satisfies $\delta_U(N) \leq 3 \deg(U)/20$. This is an explicit intrinsic action predicate. The following theorem does not assert that an arbitrary residual already lies in $\mathcal{U}$.

Theorem 31.1 (Complete low-packet transfer). *For subgroups all of whose actions lie in $\mathcal{U}$, the sector with fewer than $n/1000$ natural A_4 orbits and at least one natural C_3 or A_4 orbit has a complete ordinary forward row $O(2^{-n/500})$ and scalar at most $2^{-n^2/80000}$ eventually. The complementary sector with at least $n/1000$ natural A_4 orbits has the scalar in Theorem 30.1.*

Proof. An actual-orbit filtration gives, for a group K with no natural C_3 orbit and t natural A_4 orbits, $d_3(K) \leq 3 \deg(K)/20 + 2t/5$. First delete all $c \geq 1$ natural C_3 orbits. The remaining degree is $b = n - 3c$ and the original-degree cutoff gives

$$d_3(K) \leq (3/20 + 2/5000)b + (6/5000)c.$$

Its exact unrestricted external fibre is $\sum_j \binom{c}{j}_3 3^{jd_3(K)}$, at most $\rho_3(c+1)3^{cd_3(K)+c^2/4}$, where $\rho_3 = \prod_{i \geq 1} (1 - 3^{-i})^{-1}$. Thus the complete cell is

$$K_3(n, n-3c) = \frac{\Delta(n, n-3c)}{6^c c!} \rho_3(c+1) 3^{(3/20 + 2/5000)cb + (1/4 + 6/5000)c^2}.$$

At both parities its Gaussian loss is at least $(3c-1)b/8 + 9c^2/16 - O(c+1)$ and its coefficient cost is $O(c \log(n+2))$. Using $\log_2 3 < 8/5$, the remaining linear and quadratic coefficients exceed $9/1000$ and $27/1000$, respectively. Consequently $K_3(n, n-3c) \leq 2^{-cn/500}$ uniformly, including $b=0$; summing c gives a geometric row.

In the remaining no- C_3 sector, point one actual A_4 orbit and let $J \leq S_b$, $b = n - 4$, be its complete complement. The exact fibre is $1 + |\text{Epi}(J, C_3)| + |\text{Epi}(J, A_4)|$. Call J hot if some index-three normal subgroup F has $d_2(F) > (1/4 + 1/200)b$. For fixed $e > 0$, the graph injection $(F, \phi : F \rightarrow C_2^r) \mapsto \text{graph } \phi \leq S_{b+2r}$ and the coarse subgroup bound, with r nearest $2eb$, prove

$$|\{F \leq S_b : d_2(F) > (1/4 + e)b\}| \leq 2^{b^2/16 - e^2 b^2 + O_e(b^{3/2} + b \log(b+2))}.$$

Each F has at most $b!$ index-three overgroups, by choosing a coset generator in S_b . The complete hot pointed count, using the elementary fibre bound $5 \cdot 2^{(1/2 + (\log_2 3)/3)b}$ and its single factor $n!/(24b!)$, is at most $2^{-n^2/80000}$ after normalization.

For cold J and one surjection $\theta : J \rightarrow C_3$, let a_θ be the matching $\mathbb{F}_4$ multiplicity in $H_1(\ker \theta, \mathbb{F}_2)$. The literal cocycle count gives $4(4^{a_\theta} - 1)$ surjections to A_4 above θ , and $2a_\theta \leq (1/4 + 1/200)b$.

The residual still satisfies the original cutoff, so $d_3(J) \leq (3/20 + 2/5000)b + 8/5000$. Its complete cold fibre is at most

$$z(b) = 5 \cdot 2^{(1/4+1/200)b+(\log_2 3)((3/20+2/5000)b+8/5000)}.$$

The pointed coefficient is exactly $\Delta(n, n-4)z(n-4)/24$. Its source exponent is at most $0.49564b$, whereas deleting four points supplies Gaussian loss $b/2$ and polynomial coefficient cost. It is therefore at most $2^{-n/500}$ eventually. Choosing a deterministic actual A_4 orbit makes hot and cold physical inputs disjoint; enlarging each to all pointings pays each complete positive bound once. This proves the theorem. $\square$

32 Exhaustive counting and boundedness-first assembly

Let F_N be the normalized count of all fixed-point-free binary subgroups of S_{2N} , and retain the complete binary error E_N from Section 27. Actual orbit projections give the exact disjoint partition

$$F_N = C_{2N}^{\text{crit}} + E_N. \quad (32.1)$$

Proposition 32.1 (Native binary frontier transport). *Put*

$$B_{2N} = E_N, \quad B_{2N+1} = u_N E_N + v_N E_{N-1}.$$

There are nonnegative functions $\eta_B^{\text{amb}}(n)$ and $K_B^{\text{amb}}(n, m)$, supported on $m < n$, and constants $a'_B, C'_B > 0$ such that eventually

$$B_n \leq \eta_B^{\text{amb}}(n) + \sum_{m < n} K_B^{\text{amb}}(n, m) A_m, \quad \eta_B^{\text{amb}}(n) + \sum_{m < n} K_B^{\text{amb}}(n, m) \leq C'_B 2^{-a'_B n}. \quad (32.2)$$

Proof. Theorem 24.8 gives

$$E_N \leq \eta_B(N) + \sum_{M < N} K_B(N, M) F_M, \quad \eta_B(N) + \sum_{M < N} K_B(N, M) \leq C_B 2^{-a_B N}. \quad (32.3)$$

Every fixed-point-free binary subgroup is an ordinary subgroup, with the same normalization, so $F_M \leq A_{2M}$. In even ambient degree, use (32.3) and pad its even targets by zero to the full range $m < 2N$. In odd degree apply the same recurrence at N and $N-1$, multiply the two inequalities by u_N and v_N , and again pad both target rows by zero. The coefficients are nonnegative, and Section 27 gives

$$u_N \leq 4(N+1), \quad v_N \leq 2\Phi^{-2}(N+1)^3.$$

These fixed polynomial factors are absorbed by any smaller positive exponential rate. Since $n = 2N$ or $2N+1$, this proves (32.2). Thus the binary recurrence is transported before any boundedness of F , E , or A is used. $\square$

Theorem 32.2 (Exhaustive ordinary counting estimate). *Write $R_n = A_n - C_n^{\text{crit}}$ for the normalized complement of the complete critical family. There are nonnegative functions $\eta(n)$ and $\mathcal{K}(n, m)$, supported on $m < n$, and constants $a, C > 0$ such that eventually*

$$0 \leq R_n \leq \eta(n) + \sum_{m < n} \mathcal{K}(n, m) A_m, \quad \eta(n) + \sum_{m < n} \mathcal{K}(n, m) \leq C 2^{-an}. \quad (32.4)$$

The kernel is the sum of the outside-action row, the exact positive marker row, and the transported binary row of Proposition 32.1. Every forward edge has one complete ordinary target.

Proof. First take all subgroups with an actual transitive orbit outside the alphabet consisting of binary groups, natural C_3, A_4, S_3 , and fixed points. Theorem 20.1 bounds this complete family by an exponentially small ordinary forward row and a quadratically small scalar. Every pointing deletes its whole original orbit, so its exact weight is $n!/(b!a(U))$. A cold estimate uses it in the form $\Delta(n, b)/a(U)$ on A_b ; a hot estimate retains the factor $1/(b!a(U)p_n G_{\lfloor n/2 \rfloor})$ on its complete source sum. These are bounds for already assembled subgroups, including their other orbits and any external characters.

In the complementary alphabet let c, t be the numbers of natural C_3, A_4 orbits. If $t \geq n/1000$, apply the complete scalar of Theorem 30.1. If $t < n/1000$ and $c + t > 0$, apply Theorem 31.1. Its intrinsic action condition holds here: every relative ternary head of a two-group is zero; the normals $1, C_3, S_3$ of natural S_3 also have zero relative ternary head, since $[C_3, S_3] = C_3$ and S_3 has no ternary abelian quotient. Fixed points contribute zero. This proves the condition for every residual action in the alphabet, without an assumption about which normal kernel is selected later. Add the ordinary row and scalar of this theorem once. The macro and small packet sectors are disjoint by their original-degree cutoff.

It remains that every moved orbit is binary or natural S_3 . Write the marker counts as g, f , and $n = 2N + \varepsilon$. Under the exact collapse of Proposition 28.1, let q be the number of distinct marker signs and put $D = (g + f - \varepsilon)/2$. This is an integer, because the binary exterior has even degree, and is nonnegative. When $D \geq 1$, the complete row in Theorem 28.3 applies directly with the collection of all collapsed subgroups. It contributes $\sum_{m < n} K_{\text{mark}}(n, m)A_m$, and its row is exponentially small without any boundedness assumption on A_m .

When $D = 0$, in even degree there are no markers or fixed points. The complete binary partition gives $C_{2N}^{\text{crit}} + E_N$, each once. In odd degree there is exactly one fixed point or one natural S_3 orbit. With critical binary orbits these are exactly the two disjoint odd parts of C_n^{crit} . Otherwise their exact contributions are $\alpha_N E_N$ and $\beta_N p E_N$; the complete pair moment of Proposition 27.2 bounds their sum by $u_N E_N + v_N E_{N-1}$. Thus the entire zero-defect remainder is B_n . Proposition 32.1 replaces this term by one exponentially small scalar and one strictly forward ordinary row. No estimate for a restricted binary residual is substituted for the complete moment.

These sectors exhaust all actual subgroups and are pairwise disjoint before their positive enlargements. Within the first sector fix an order of the proved action families and assign a subgroup to the first one it meets, before testing temperatures. For each assigned family its hot part means that some accepted pointing is hot, and its cold part that none is hot. Each theorem's complete positive envelope is then paid once, over all its assigned inputs. Multiple orbit occurrences are already accounted for by pointing the original weight $\prod_U (a(U)^{m_U} m_U!)^{-1}$; auxiliary representations and alternative certificates contribute no new physical factorial. All subsequent sectors are restrictions of their proved positive envelopes and are likewise paid once.

Define η to be the sum of the nonbinary, macro, small-packet, and transported binary scalars. Define $\mathcal{K}$ to be the sum of the nonbinary and small-packet ordinary rows, the positive marker row, and the transported binary row. Each summand is nonnegative, strictly forward, and exponentially small in total mass. Taking the minimum of their positive rates gives (32.4). $\square$

Proof of Theorem 1.3. The critical ratios C_n^{crit} are bounded. Choose an onset at which the row in (32.4) is at most $1/2$, and let M exceed all finitely many initial A_m and twice a uniform bound for $C_n^{\text{crit}} + \eta(n)$. The exact partition $A_n = C_n^{\text{crit}} + R_n$ and induction in n give $A_n \leq M$ for every n . Substituting this established bound into (32.4) gives $R_n = O(2^{-cn})$ for some $c > 0$. The critical-family estimate now yields $|A_n - 1| = O(2^{-c'n})$. Taking the smaller positive rate proves the result at both parities. No boundedness premise was used in deriving any row or scalar;

boundedness is a consequence of their eventual contraction. $\square$

33 Evaluation of the benchmark

The results in this section estimate the explicit sequence L_n in (1.2). Theorem 1.3 transfers them to the subgroup count with an exponentially smaller relative error.

33.1 The binary Gaussian sum

Define

$$\kappa_0 = \phi^{-1} \sum_{j \in \mathbb{Z}} 2^{-j^2}, \quad \kappa_1 = \phi^{-1} \sum_{j \in \mathbb{Z}} 2^{-j(j-1)}. \quad (33.1)$$

Both series converge absolutely.

Lemma 33.1. *As $R \rightarrow \infty$,*

$$G_R = \kappa_{R \bmod 2} 2^{\lfloor R^2/4 \rfloor} (1 + O(R2^{-R/2})).$$

Proof. The product formula reads

$$\left[\begin{matrix} R \\ k \end{matrix} \right]_2 = 2^{k(R-k)} \frac{\prod_{i=R-k+1}^R (1 - 2^{-i})}{\prod_{i=1}^k (1 - 2^{-i})}.$$

For central k the fraction is $\phi^{-1}(1 + O(2^{-k} + 2^{-(R-k)}))$. When $R = 2m$, put $k = m + j$, giving exponent $m^2 - j^2$. When $R = 2m + 1$, put $k = m + j$, giving exponent $m(m+1) - j(j-1)$. On $|j| \leq R/4$ the summable Gaussian weights times $2^{|j|}$ give error $O(2^{-R/2})$ up to an absolute parity constant. The remaining tails are $2^{-\Omega(R^2)}$ by (2.1). The stated, slightly weaker, error follows. $\square$

33.2 One positive saddle

Let $\rho = \rho_R > 0$ be the unique solution of

$$R = \rho P'(\rho) = \frac{\rho}{2} + \frac{\rho^2}{3} + \frac{\rho^4}{96}, \quad b(\rho) = \frac{\rho}{2} + \frac{2\rho^2}{3} + \frac{\rho^4}{24}. \quad (33.2)$$

Existence and uniqueness follow from strict monotonicity on the positive axis. In particular $\rho \sim (96R)^{1/4}$ and $b(\rho) \sim 4R$.

Theorem 33.2 (Saddle formula). *For $n = 2R + \varepsilon$,*

$$L_n = n! G_R \frac{(1 + \rho/6)^\varepsilon e^{P(\rho)}}{\rho^R \sqrt{2\pi b(\rho)}} \left(1 + O(R^{-1})\right). \quad (33.3)$$

In particular $c_{R-1}/c_R = \rho(1 + O(R^{-1}))$.

Proof. Put $\lambda_1 = \rho/2$, $\lambda_2 = \rho^2/6$ and $\lambda_4 = \rho^4/384$, so that $R = \sum_k k\lambda_k$ and $b = \sum_k k^2\lambda_k$ satisfy $R \leq b \leq 4R$. Here and below $k \in \{1, 2, 4\}$. Define

$$A(\theta) = \sum_k \lambda_k (1 - \cos k\theta), \quad B(\theta) = \sum_k \lambda_k (\sin k\theta - k\theta).$$

Then $P(\rho e^{i\theta}) - P(\rho) - iR\theta = -A + iB$. For any $d \in [0, 1]$, set

$$F_d(\theta) = (1 - d) \cos B(\theta) + d \cos(\theta + B(\theta)), \quad H_d(\theta) = e^{-A(\theta)} F_d(\theta).$$

Taking real parts in Cauchy's formula on $y = \rho e^{i\theta}$ gives

$$\frac{\rho^R [y^R] (1 + y/6)^\varepsilon e^{P(y)}}{(1 + \rho/6)^\varepsilon e^{P(\rho)}} = \frac{1}{2\pi} \int_{-\pi}^{\pi} H_d(\theta) d\theta, \quad d = \begin{cases} 0, & \varepsilon = 0, \\ \rho/(6 + \rho), & \varepsilon = 1. \end{cases}$$

We estimate this integral uniformly for all $d \in [0, 1]$.

Use the fixed central interval $|\theta| \leq \alpha = \pi/4$ and write $Q = b\theta^2/2$. The inequalities $u^2/8 \leq 1 - \cos u \leq u^2/2$ for $|u| \leq \pi$, applied to each $k\theta$, give

$$R\theta^2/8 \leq A \leq Q, \quad 0 \leq Q - A = O(R\theta^4), \quad |B| = O(R|\theta|^3).$$

The last two bounds follow from the fourth-order cosine and third-order sine remainders; their constants are absolute. Since F_d is a convex combination of cosines,

$$|F_d| \leq 1, \quad 0 \leq 1 - F_d \leq B^2/2 + |\theta B| + \theta^2/2.$$

Together with $0 \leq e^{-A} - e^{-Q} \leq e^{-A}(Q - A)$, this yields

$$|H_d(\theta) - e^{-b\theta^2/2}| \leq C e^{-R\theta^2/8} (\theta^2 + R\theta^4 + R^2\theta^6).$$

Integrating this bound gives $O(R^{-3/2})$, by the change of variable $u = \sqrt{R}\theta$. This argument requires no smallness assumption on B .

On the entire complement $\alpha \leq |\theta| \leq \pi$, the linear term alone gives

$$A(\theta) \geq \frac{\rho}{2}(1 - \cos \alpha) \geq \frac{\rho}{16}, \quad |H_d(\theta)| \leq e^{-\rho/16}.$$

Because $\rho \asymp R^{1/4}$, these tails are smaller than every fixed inverse power of R . The Gaussian tails outside $[-\alpha, \alpha]$ are $O(e^{-cR})$ for an absolute $c > 0$. Thus, uniformly in d ,

$$\int_{-\pi}^{\pi} H_d(\theta) d\theta = \sqrt{\frac{2\pi}{b}} + O(R^{-3/2}) = \sqrt{\frac{2\pi}{b}} (1 + O(R^{-1})),$$

which proves (33.3). Finally, Cauchy's formula with amplitude y/ρ gives H_1 , at the same saddle ρ_R . Comparing it with H_0 gives

$$\frac{c_{R-1}}{\rho c_R} = \frac{\int_{-\pi}^{\pi} H_1(\theta) d\theta}{\int_{-\pi}^{\pi} H_0(\theta) d\theta} = 1 + O(R^{-1}). \quad \square$$

33.3 An elementary four-periodic expression

Set

$$\begin{aligned} C_0 &= e^{-4/3} \kappa_0 / 2^{1/2}, & C_2 &= e^{-4/3} \kappa_1 / 2^{3/4}, \\ C_1 &= \frac{e^{-4/3} \kappa_0 48^{3/8}}{6 \cdot 2^{7/16}}, & C_3 &= \frac{e^{-4/3} \kappa_1 48^{3/8}}{6 \cdot 2^{11/16}}. \end{aligned} \quad (33.4)$$

For $\varepsilon = n \bmod 2$ define

$$M_n = C_{n \bmod 4} n^{3\varepsilon/8} 2^{n^2/16} \left(\frac{n^7}{48 \cdot 2^\varepsilon e^7} \right)^{n/8} \exp \left(2\sqrt{n/3} + \frac{(48n)^{1/4}}{2} \right). \quad (33.5)$$

Theorem 33.3 (Elementary expansion). *The exact benchmark satisfies*

$$\frac{L_n}{M_n} = 1 + \frac{6\varepsilon - 4}{(48n)^{1/4}} + O(n^{-1/2}). \quad (33.6)$$

By Theorem 1.3, the same formula holds with s_n in place of L_n , and (33.3) holds with s_n on the left.

Proof. Put $x = (96R)^{1/4}$ and $v = x(x - \rho)$. Only the first displacement of the saddle is needed:

$$\rho = x - 8x^{-1} + O(x^{-2}), \quad v = 8 + O(x^{-1}). \quad (33.7)$$

Indeed, the equation is $\rho^4 + 32\rho^2 + 48\rho = x^4$. For large x its positive root lies in $[x/2, x]$; substituting $x - 8/x$ leaves residual $O(x)$, and the derivative throughout the intervening interval is of order x^3 . The mean value theorem proves (33.7).

Expand $\log_e \rho = \log_e x + \log_e(1 - v/x^2)$ to quadratic order. Since v is bounded, the cubic remainder, multiplied by $R = x^4/96$, is $O(x^{-2})$. Direct substitution gives

$$\begin{aligned} P(\rho) - R \log_e \rho &= \frac{R}{4} - R \log_e x + \frac{x^2}{6} + \frac{x}{2} + \frac{v^2}{48} - \frac{v}{3} - \frac{v}{2x} + O(x^{-2}) \\ &= \frac{R}{4} - R \log_e x + \frac{x^2}{6} + \frac{x}{2} - \frac{4}{3} - \frac{4}{x} + O(x^{-2}). \end{aligned}$$

The cancellation is explicit: $v^2/48 - v/3 = -4/3 + (v - 8)^2/48$. Also $b(\rho)/(4R) = 1 + O(x^{-2})$.

Define

$$T_R = \frac{1}{\sqrt{8\pi R}} \exp \left(-\frac{R}{4} \log_e R + \frac{1 - \log_e 96}{4} R + \frac{2\sqrt{6}}{3} \sqrt{R} + \frac{x}{2} - \frac{4}{3} \right).$$

The preceding estimates and the saddle formula yield

$$\log_e \frac{c_R}{T_R} = -\frac{4}{x} + O(x^{-2}). \quad (33.8)$$

The relative saddle error is $O(R^{-1}) = O(x^{-4})$ and is absorbed in this remainder. For the odd amplitude, the same first displacement gives

$$\log_e \frac{1 + \rho/6}{x/6} = \frac{6}{x} + O(x^{-2}). \quad (33.9)$$

To make the parity conversion explicit, set $y = (48n)^{1/4}$ and

$$\begin{aligned} \Lambda_n &= (n + \tfrac{1}{2}) \log_e n - n + \tfrac{1}{2} \log_e(2\pi) + \lfloor R^2/4 \rfloor \log_e 2 + \log_e \kappa_{R \bmod 2} \\ &\quad + \log_e T_R + \varepsilon \log_e(x/6) + (6\varepsilon - 4)/x. \end{aligned}$$

Stirling's formula, Lemma 33.1 and the two expansions above give $\log_e L_n = \Lambda_n + O(n^{-1/2})$. Using $R = (n - \varepsilon)/2$ and the exact constants in (33.4), elementary algebra gives

$$\begin{aligned} \Lambda_n - \log_e M_n - \frac{6\varepsilon - 4}{y} &= \frac{n - 3\varepsilon + 4}{8} \log_e \frac{n}{n - \varepsilon} - \frac{\varepsilon}{8} \\ &\quad + \frac{x^2 - y^2}{6} + \frac{x - y}{2} + (6\varepsilon - 4) \left(\frac{1}{x} - \frac{1}{y} \right). \end{aligned}$$

In even degree this is identically zero. In odd degree the first line is $O(n^{-1})$, while $y^4 - x^4 = 48$ bounds the remaining three differences by $O(n^{-1/2})$, $O(n^{-3/4})$ and $O(n^{-5/4})$, respectively. Thus

$$\log_e(L_n/M_n) = (6\varepsilon - 4)(48n)^{-1/4} + O(n^{-1/2}).$$

Exponentiation proves (33.6). Finally the exponential relative error of Theorem 1.3 is smaller than either polynomial error asserted here. $\square$

Remark 33.4. The relative errors refer to three different approximations. The complete counting theorem gives exponential error against the exact coefficient benchmark L_n . Replacing the coefficient by its single-saddle expression gives $O(n^{-1})$. Replacing the saddle by the elementary expression M_n gives an explicit term of order $n^{-1/4}$ and a remainder $O(n^{-1/2})$.

A Finite evidence and reproduction

The accompanying computations distinguish three assertions: a literal record satisfies its algebraic equations; a stated finite domain is covered by such records; and an infinite counting theorem follows from that finite domain together with proved structural and summation lemmas. These assertions require different arguments.

The publication package contains the following mathematical data.

Dataset	Mathematical role
Small binary actions	Literal actions through degree 16, normal axes, quotient maps, and the four local witness types.
Carrier data	Twelve noncritical actions, normal profiles, transition data, and specified quotient-replacement maps.
Nonbinary and rank data	Finite action and normal-rank domains used in the selected primitive and pair-top reductions.
Pair-top data	Concrete modules and constructive witnesses in the count-twelve and count-sixteen domains.
Four- and six-block data	Constructive classifications and actual quotient interfaces.
Zero-ternary frame data	Thirty complete binary module lattices and thirty-eight literal odd-subgroup/normalizer witnesses.
Critical and audit data	Quadratic realization constants, critical normalizers, and independent local chart controls.

The data contracts specify literal point actions, normal subgroups, source and target maps, and the original symmetric normalizers. A quotient order or isomorphism-type name alone does not specify a counting witness. For a proper subdirect replacement, the image group and both its projection maps are part of the record.

For the critical quadratic constants of Lemma 2.3, one can enumerate $\mathrm{GL}(2, 2)$ and $\mathrm{GL}(4, 2)$ and test the equation $q(gv) = q(v)$ on every vector. This establishes the orders 2 and 72 for the displayed forms. The torsor argument in the lemma is what extends these finite calculations to arbitrary source dimension. Likewise the exact rational inequality (21.5) is a finite arithmetic certificate whose implication for all degree-32 sections is proved by the preceding incidence argument.

The classification libraries are pinned to TransGrp 3.6.5 and PrimGrp 3.4.4 [Hul23, HRDR23]; their mathematical construction and scope are described in [Hul05, HR20, RD05]. The recorded interpreter is GAP 4.13.1 [The24]. Where a coverage proof uses an independent constructive

generation argument, the catalogue is a source of literal representatives, not its sole completeness argument.

All reproduction commands are local. The file `computations/README.md` documents the producers and independent checkers, and `certificates/manifests/finite_witnesses.json` binds their finite input and output files. Hashes identify bytes; they do not prove a mathematical theorem. The checks do not initialize or invoke a proof assistant.

The same-source C_3 example, the inverse-complement reciprocal, the proper-subdirect transport examples, and the binary duplicate factor $1/4$ are regression conditions on the counting maps. Passing these conditions alone does not establish an asymptotic theorem. The binary coverage principles are proved in Lemma 24.5; the universal mixture and recurrence arguments are proved in Theorems 23.15 and 24.8. The complete nonbinary exhaustion is proved in Theorem 20.1; the ordinary counting inequality and its boundedness-first induction are proved in Section 32. The supplementary affine-nine and narrow socket data provide independent local checks; the proof above uses the general frame and mixture arguments for those domains.

B The finite carrier reduction and its two rational cones

The finite local input consists of the eight master groups and all their literal normal kernels. The respective numbers of kernels are 27, 13, 28, 2911, 2911, 2911, 2292, 2915. Their generators and all central-involution extension edges are supplied in the carrier normal certificate. Lemma 24.5 proves completeness from these edges. For each kernel the checked tuple is (k, n, m, a_2, c, g) as in Lemma 23.1 and (23.4). In particular the maximum defining m is computed over the same complete normal list, and the original quotient is used for c, g .

For the five degree-sixteen groups, the structural proof in Lemma 23.10 gives exhaustive original-normal coverage without constructing these normal lists. Effective polygon inclusion reduces its rows to the displayed H16 rows and the two additional sixth-colour envelopes described in the proof above. The latter obey all displayed cross-column comparisons and their mutual comparison, so they use the same rational cone certificate below.

For the degree-sixteen groups the identities $\Phi(G) = G'$ permit $a_2 \leq m$: here $N^2[N, G] \leq N \cap G'$. Their rows are dominated by the following twelve scale-two rows; the other three blocks below are the scale-one rows of X, J, P . Domination is simultaneous on the pair supports and both marks $k, q = \max(m, a_2)$, including every cross column between degrees eight and sixteen.

<i>H16</i>	<i>k</i>	<i>n</i>	<i>m</i>	<i>a</i> ₂	<i>c</i>	<i>g</i>
0	0	0	0	0	1	6
1	1	1	1	1	2	5
2	1	2	1	1	4	4
3	2	3	2	2	3	3
4	2	4	2	2	4	2
5	3	6	3	3	6	0
6	4	6	4	4	5	0
7	4	7	4	4	4	0
8	4	8	4	4	3	0
9	5	11	3	3	1	0
10	5	11	4	4	0	0
11	6	12	3	3	0	0

X	k	n	m	a_2	c	g
0	0	0	0	0	1	3
1	1	1	1	0	2	2
2	1	2	1	1	2	1
3	1	3	1	1	1	1
4	1	4	2	2	2	0
5	2	3	2	1	3	0
6	2	4	2	1	2	0
7	2	5	2	2	1	0
8	3	6	2	2	0	0

J	k	n	m	a_2	c	g
0	0	0	0	0	1	3
1	1	1	1	0	1	2
2	1	2	1	1	2	1
3	1	3	1	1	1	1
4	1	3	1	1	3	0
5	1	4	1	1	2	0
6	1	5	1	2	1	0
7	2	4	1	1	2	0
8	2	5	1	1	1	0
9	2	6	1	2	0	0

P	k	n	m	a_2	c	g
0	0	0	0	0	1	4
1	1	1	1	0	1	3
2	1	2	1	1	2	2
3	1	3	1	1	2	1
4	1	4	1	1	1	1
5	1	5	2	2	2	0
6	2	4	2	1	3	0
7	2	5	2	1	2	0
8	2	6	2	2	1	0
9	3	7	2	2	0	0

The normalized representatives are

	k	n	m	a_2	c	g	s
H_9	5	11	3	3	1	0	2
X_{whole}	3	6	2	2	0	0	1
P_0	0	0	0	0	1	4	1
P_1	1	1	1	0	1	3	1
P_2	1	2	1	1	2	2	1
P_6	2	4	2	1	3	0	1
P_{whole}	3	7	2	2	0	0	1

For completeness, the following list gives a dominating representative for each row in each displayed block, in its displayed order. The last representative has exactly the same normalized pair row and marks as X_{whole} , so they are merged. All comparisons here are finite rational inequalities of the

explicit support functions. They can be checked without enumerating the feasible integer pairs: for $x, y \geq 0$, the support of the row (k, n, m, a_2, c, g) is at most

$$\min\{xk + ym, (x - y)_+k + (y - x)_+m + \min(x, y)n\}.$$

The first expression uses the two separate capacity bounds. For the second, use the joint bound on $\delta + \epsilon$ and the separate bound for the variable with the larger coefficient. Substitution of the displayed rational entries proves all 41^2 ordered pair comparisons, with their original scales; the two mark comparisons are coordinatewise. This specialization is used only for the nonnegative quotient slopes, not to discard a negative first mark in the recurrence.

$$H16 : (P_0, P_1, P_2, P_2, P_2, P_6, P_6, P_6, P_6, H_9, X_{\text{whole}}, X_{\text{whole}}).$$

$$X : (P_0, P_2, P_2, P_2, P_6, P_6, P_6, P_6, X_{\text{whole}}).$$

$$J : (P_0, P_1, P_2, P_2, P_2, P_2, P_6, P_6, P_6, X_{\text{whole}}).$$

$$P : (P_0, P_1, P_2, P_2, P_2, P_6, P_6, P_6, P_6, X_{\text{whole}}).$$

The reduction of the actual degree-sixteen tuples to the twelve rows also checks all degree-eight columns, and the two marks, in the committed transition certificate. Thus the preceding reduction does not presume that domination inside a smaller submatrix survives the addition of new colours. Scaling and merging give exactly (23.6).

It remains to prove the scalar inequality used in Theorem 23.13. Put $p = \sum_i p_i$ and

$$\mathcal{D} = R^2/4 + (139/328)Rp + (627/2624)p^2 - \mathcal{E}.$$

In the first cone let $(L, J, S) = (\ell, j - 2\ell, R - j)$, and in the second let $(A, C, S) = (j, 2\ell - j, R - 2\ell)$. These are nonnegative in their respective cones. Let z consist of $p_1, \dots, p_6$ followed by the three corresponding cone coordinates. Expansion gives $5248\mathcal{D} = z^\top M_\nu z$. The upper left 6×6 block, common to both M_ν , is

$$M_0 = \begin{pmatrix} 1049 & 1008 & -140 & 106 & -58 & 24 \\ 1008 & 1254 & -550 & -222 & -386 & -222 \\ -140 & -550 & 1254 & 598 & 434 & -386 \\ 106 & -222 & 598 & 762 & 598 & -58 \\ -58 & -386 & 434 & 598 & 598 & -58 \\ 24 & -222 & -386 & -58 & -58 & 270 \end{pmatrix}.$$

Write $M_\nu = \begin{pmatrix} M_0 & D_\nu \\ D_\nu^\top & E_\nu \end{pmatrix}$. The remaining blocks are

$$D_A = \begin{pmatrix} -400 & -528 & 1112 \\ -1056 & -856 & 1112 \\ 2224 & 1112 & 1112 \\ 912 & 456 & 1112 \\ 912 & 456 & 1112 \\ -400 & -200 & 1112 \end{pmatrix}, \quad E_A = \begin{pmatrix} 5248 & 1312 & -1312 \\ 1312 & 1312 & -1312 \\ -1312 & -1312 & 1312 \end{pmatrix}.$$

$$D_B = \begin{pmatrix} -200 & -200 & 1112 \\ -528 & -528 & 1112 \\ 1112 & 1112 & 1112 \\ 456 & 456 & 1112 \\ 456 & 456 & 1112 \\ -200 & -200 & 1112 \end{pmatrix}, \quad E_B = \begin{pmatrix} 1312 & 0 & -656 \\ 0 & 3936 & 1968 \\ -656 & 1968 & 1312 \end{pmatrix}.$$

The following rational matrices give positive-semidefinite parts.

$$P_A = \begin{pmatrix} 817 & 804 & -548 & -328 & -393 & -\frac{540}{11} & -896 & -688 & 688 \\ 804 & 1254 & -550 & -288 & -386 & -222 & -1129 & -856 & 856 \\ -548 & -550 & 1254 & 423 & 434 & -386 & 1836 & 1112 & -1112 \\ -328 & -288 & 423 & 577 & 237 & -\frac{3987}{44} & 563 & 323 & -323 \\ -393 & -386 & 434 & 237 & 490 & -58 & 646 & 363 & -363 \\ -\frac{540}{11} & -222 & -386 & -\frac{3987}{44} & -58 & 270 & -\frac{19363}{44} & -200 & 200 \\ -896 & -1129 & 1836 & 563 & 646 & -\frac{19363}{44} & 4645 & 1312 & -1312 \\ -688 & -856 & 1112 & 323 & 363 & -200 & 1312 & 1312 & -1312 \\ 688 & 856 & -1112 & -323 & -363 & 200 & -1312 & -1312 & 1312 \end{pmatrix}.$$

$$P_B = \begin{pmatrix} 729 & 721 & -469 & -236 & -318 & -\frac{602}{11} & -416 & -480 & 86 \\ 721 & 1254 & -550 & -296 & -386 & -222 & -565 & -585 & -2 \\ -469 & -550 & 1254 & 433 & 434 & -386 & 917 & 950 & -407 \\ -236 & -296 & 433 & 542 & 241 & -\frac{4061}{44} & 275 & 197 & -56 \\ -318 & -386 & 434 & 241 & 469 & -58 & 303 & 231 & -65 \\ -\frac{602}{11} & -222 & -386 & -\frac{4061}{44} & -58 & 270 & -\frac{2413}{11} & -\frac{10005}{44} & \frac{8581}{44} \\ -416 & -565 & 917 & 275 & 303 & -\frac{2413}{11} & 1254 & -134 & -813 \\ -480 & -585 & 950 & 197 & 231 & -\frac{10005}{44} & -134 & 2690 & 591 \\ 86 & -2 & -407 & -56 & -65 & \frac{8581}{44} & -813 & 591 & 891 \end{pmatrix}.$$

Subtraction gives $M_A - P_A \geq 0$ entrywise, with entries at most 2224, and $M_B - P_B \geq 0$ entrywise, with entries at most 1519. These assertions are direct comparisons of the displayed rational entries. Both matrices annihilate $v = (0, 17, 21, 0, 0, 44, 0, 0, 0)$; P_A also annihilates $w = (0, 0, 0, 0, 0, 0, 0, 1, 1)$. Delete coordinates six and nine from P_A . The successive leading principal determinants of the retained matrix are

$$\begin{aligned} &817, \quad 378102, \quad 335067392, \quad 132631407594, \\ &35034163435774, \quad 63581756789450739, \quad 2100371145348629273. \end{aligned}$$

They are positive. The omitted coordinates of (v, w) form $\text{diag}(44, 1)$, so subtracting their unique linear combination carries any vector into this retained subspace without changing its quadratic value. Thus P_A is positive semidefinite. Delete coordinate six from P_B . Its retained leading principal determinants are

$$\begin{aligned} &729, \quad 394325, \quad 370093856, \quad 140632346331, \\ &36346448139435, \quad 19760569015182064, \\ &7216647669027584752, \quad 1335792379687239644084. \end{aligned}$$

Again they are positive, and the omitted coordinate of v is 44, so the same kernel elimination proves that P_B is positive semidefinite. Hence $z^\top M_\nu z \geq 0$ for every nonnegative z . The two cones exhaust all $0 \leq j \leq R$, $0 \leq \ell \leq R/2$, including $j < \ell$, proving the required scalar inequality.

References

- [BBERJS22] Adolfo Ballester-Bolínches, Ramón Esteban-Romero, and Paz Jiménez-Seral. Maximal subgroups of small index of finite almost simple groups, 2022. Theorem A(4), version 2, p. 4.

- [GM15] Martino Garonzi and Attila Maróti. On the number of conjugacy classes of a permutation group. *Journal of Combinatorial Theory, Series A*, 133:251–260, 2015.
- [GPRV18] S. P. Glasby, Cheryl E. Praeger, Kyle Rosa, and Gabriel Verret. Bounding the composition length of primitive permutation groups and completely reducible linear groups. *Journal of the London Mathematical Society*, 98(3):557–572, 2018.
- [HM16] Zoltán Halasi and Attila Maróti. The minimal base size for a p -solvable linear group. *Proceedings of the American Mathematical Society*, 144(8):3231–3242, 2016.
- [HR20] Derek F. Holt and Gordon Royle. A census of small transitive groups and vertex-transitive graphs. *Journal of Symbolic Computation*, 101:51–60, 2020.
- [HRD13] Derek F. Holt and Colva M. Roney-Dougal. Minimal and random generation of permutation and matrix groups. *Journal of Algebra*, 387:195–214, 2013.
- [HRDR23] Alexander Hulpke, Colva M. Roney-Dougal, and Christopher Russell. *PrimGrp: GAP Primitive Permutation Groups Library*, 3.4.4 edition, 2023. Released 25 February 2023.
- [Hul05] Alexander Hulpke. Constructing transitive permutation groups. *Journal of Symbolic Computation*, 39(1):1–30, 2005.
- [Hul23] Alexander Hulpke. *TransGrp: Transitive Groups Library*, 3.6.5 edition, 2023. Released 15 December 2023.
- [KP89] L. G. Kovács and Cheryl E. Praeger. Finite permutation groups with large abelian quotients. *Pacific Journal of Mathematics*, 136(2):283–292, 1989.
- [LMM00] Andrea Lucchini, Federico Menegazzo, and Marta Morigi. Asymptotic results for transitive permutation groups. *Bulletin of the London Mathematical Society*, 32(2):191–195, 2000. The labelled transitive-subgroup bound is cited through the explicit statement in Roney-Dougal and Tracey (2025), version 1, introduction, p. 2.
- [Mar05] Attila Maróti. Bounding the number of conjugacy classes of a permutation group. *Journal of Group Theory*, 8(3):273–289, 2005.
- [RD05] Colva M. Roney-Dougal. The primitive permutation groups of degree less than 2500. *Journal of Algebra*, 292(1):154–183, 2005.
- [RDT25] Colva M. Roney-Dougal and Gareth Tracey. Subgroups of symmetric groups: enumeration and asymptotic properties, 2025. Version 1; all theorem locators in the input register refer to this version.
- [The24] The GAP Group. *GAP: Groups, Algorithms, Programming, a System for Computational Discrete Algebra*, 4.13.1 edition, 2024.
- [Tra18] Gareth M. Tracey. Minimal generation of transitive permutation groups. *Journal of Algebra*, 509:40–100, 2018.
- [Tra21] Gareth Tracey. Sharp upper bounds on the minimal number of elements required to generate a transitive permutation group, 2021. Version 1, Theorem 1.1.