

COVID-19 vaccination in Ontario: Exploring intra-provincial variations within Health Regions and socio-economic strata

Ariel Mundo Ortiz¹ Bouchra Nasri^{2,*}

¹ Centre de Recherches Mathématiques, University of Montreal, Montréal, Canada

² Department of Social and Preventive Medicine, École de Santé Publique, University of Montreal, Montréal, Canada

* Correspondence: [Bouchra Nasri <bouchra.nasri@umontreal.ca>](mailto:bouchra.nasri@umontreal.ca)

Abstract

The COVID-19 pandemic continues to be a worldwide public health concern. Although vaccines against this disease were rapidly developed, vaccination uptake has not been equal across all the segments of the population. In particular, it has been shown that there have been differences in vaccine uptake across different segments of the population. However, there are also differences in vaccination across geographical areas, which might be important to consider in the development of future public health policies against COVID-19. In this study, we examined the relationship between vaccination status (having received the first dose of a COVID-19 vaccine), and different socio-economic and geographical factors. Our results show differences in vaccination due to race/ethnicity, income, Health Regions (geographical areas used for health service access in Ontario), and their interactions. In particular, we show that individuals who identified as Arab/Middle Eastern, Black, or Latin American, had significantly lower odds of vaccination than White/Caucasian individuals (ORs=0.31, 0.32, 0.28, and $p=0.004$, $p<0.001$ and $p=0.004$, respectively), and that individuals with a household income below CAD 25,000 who identified as Arab/Middle Eastern (OR=3.05, $p=0.013$), Black (OR=3.19, $p=0.004$), Latin American (OR=2.80, $p=0.041$), or that belonged to other minority groups (OR=4.59, $p<0.001$) had higher odds of vaccination than individuals from the same racial/ethnic group in higher income brackets. Finally, we also identified lower odds of vaccination within certain minority groups in the West Health Region, which comprises the regions of Waterloo and Niagara, the counties of Wellington, Essex and Lambton, and the

cities of Hamilton, Haldimand, Brant, and Chatham-Kent. This study shows that there is an ongoing need to better understand and address differences in vaccination uptake across diverse segments of the population that have been largely impacted by the pandemic.

Keywords

Covid-19, vaccination, survey, socio-economic factors, visible minorities.

Background

The vaccines against COVID-19 have been considered a major achievement of modern medicine as their rapid development allowed the start of broad vaccination campaigns towards the end of 2020 in certain countries, such as the US and Canada¹⁻³. This made some believe that vaccines were destined to be a determinant factor in a rapid ending of the pandemic⁴. However, although it has been estimated that COVID-19 vaccines have prevented around 14 million of deaths worldwide⁵, their implementation has been far from being equal to that of the vaccines of smallpox and polio, which were implemented on a global scale and that were indeed crucial to control these diseases⁶. In fact, the rollout of COVID-19 vaccines has faced multiple challenges since its inception which ultimately have hampered their use to achieve the ultimate goal of global immunity.

This problematic of COVID-19 vaccines rollout is a multifaceted issue resulting from, among other things, the development of new variants due to inadequate public health measures⁷, inequality in vaccine access between high- and low-income countries^{8,9}, vaccine hesitancy¹⁰, and differences in vaccination uptake across different segments of the population¹¹. In particular, it is well established that differences in vaccination uptake have been present even in countries that have had ample access to vaccines since 2020 (such as the US, the UK, and Canada), where lower vaccine uptake has been observed within racial minorities (i.e., individuals that identify as Black, Asian, or Indigenous), and in individuals within low income brackets¹²⁻¹⁵. Reasons given for lower vaccine uptake in these cases have included medical mistrust due to systemic medical racism¹⁴, mistrust in vaccines¹², and the influence of conspiracy theories¹⁶⁻¹⁸. Moreover, in the case of Canada, lower vaccine uptake has been observed in young individuals, those with a low educational level, households with children, those without a regular healthcare provider, individuals that identify as part of a visible minorities or Indigenous, and those that are financially unstable¹⁹⁻²¹.

However, it is important to consider that vaccination uptake can also be influenced by geographical (spatial) factors. In this regard, differences in COVID-19 vaccination rates have been associated with varied regional attitudes towards vaccination¹¹, spatial differences in

vaccine access and supply, vaccination location availability, and lack of prioritization of areas where vulnerable groups reside^{2,22}. Other studies have also shown heterogeneity in vaccine uptake within small governmental administrative units such as counties²³⁻²⁶, and that and that accounting for geographical differences in vaccination can help predict patterns of booster uptake²⁷. Overall, the evidence provided by the literature demonstrates the existence of spatially-driven heterogeneities in vaccine uptake that be used by decision-makers in the development of public health policies that are focused on addressing these disparities within specific administrative or geographical areas.

However, such analyses have been carried mostly in territories outside of Canada, where available studies have been focused in certain cities (such as Toronto²⁸, or Montreal²⁹), or have explored differences at a province-wide level¹⁹. Thus, there is a need for studies that explore spatial differences in vaccination within the Canadian territory and that consequently, can help identify disparities that need to be addressed within specific areas in each province.

This need is particularly important in the case of Ontario, the most populated province in Canada. Between 2006 and 2019, Ontario provided healthcare access to its inhabitants using 14 intra-provincial divisions called the Local Health Integrated Networks (LHINs). However, this approach was complex, bureaucratic, and led to systemic inequalities³⁰. In late 2019, the 14 LHINs were phased out and the areas they covered were incorporated into 6 Health Regions (North East, North West, Central, Toronto, West, and East) in an effort to improve the healthcare system of the province³¹. Because the adoption of the Health Regions occurred at a relatively recent time, there is an ongoing need to analyze the impact of this measure and identify the existence of intra-regional differences that might exist, and which could be specially important in the context of the COVID-19 pandemic.

In this study, we analyzed differences in self-reported COVID-19 vaccination status in Ontario using socio-economic (e.g., income, racial/ethnic identification), and spatial information at the level of the Health Regions to identify the existence of differences that might need to be addressed to ensure that the healthcare system of Ontario is more inclusive and that responds to the needs of its most vulnerable population.

Methods

Data

We used data from the *Survey of COVID-19 related Behaviours and Attitudes*, a repeated cross sectional survey focused on the Canadian province of Ontario that was commissioned by the Fields Institute for Research in Mathematical Sciences (henceforth Fields) and the Mathematical Modelling of COVID-19 Task Force under ethical guidance from the University of Toronto, and which ran between September 30th, 2021 and January 17th, 2022. The survey collected socio-economic information from participants (Table 1), recorded their location (using the nearest municipality), and asked information on vaccination status by using the question

“Have you received the first dose of the COVID vaccine?”, with possible answers “yes” and “no”. The original dataset contained 39,029 entries (where each entry corresponded to a unique respondent).

This dataset was cleaned to remove outliers that were identified during preliminary analyses, and processing the geographical information in the survey (city where the survey was responded) in order to match each city to its correspondent Health Region.

Table 1: Socio-economic factors from the Fields COVID-19 survey

Variable	Levels
Age group	16-34,35-54,55 and over
Income bracket (CAD)	under 25,000, 25,000-59,999, 60,000 and above
Race/ethnicity	Arab/Middle Eastern, Black, East Asian/Pacific Islander, Indigenous, Latin American, Mixed, South Asian, White Caucasian, Other

The clean dataset contained responses from more than 200 different municipalities within Ontario (Figure 1). Because of the lack of a publicly available list of all municipalities within each Health Region, we used a dataset of long-term care homes and LHINs to match each city to LHIN, followed by matching each LHIN to a Health Region following the information provided on the Ontario Health Website, where the list of LHINs and corresponding Health Regions is available. In the case of municipalities that did not appear in the long-term care home dataset, we manually searched each city in the LHINs websites in order to provide geographical information. The original dataset, clean dataset, and details on the data cleaning process are described in detail in the GitHub repository for this paper, which can be found at https://github.com/aimundo/Fields_COVID-19/.

Following an assessment of the number of entries corresponding to each Health Region in the final dataset, only 107 observations (4.3% of the total) corresponded to cities located in the North West and North East Health Regions. The low representation of these Health Regions in the dataset is noticeable in Figure 1, which shows that responses from these areas came from a relatively low number of cities when compared to the most populated Health Regions, such as the Toronto or Central Regions. We omitted the North East and North East Health Regions from further analyses due to the low number of entries. Therefore, the total number of unique entries used for analysis was 3,551 which included the East, Central, Toronto, and West Health Regions and that covered the period between September 30th, 2021 and December 12th, 2021.




Figure 1: Geographic representation of the data collected by the *Survey of COVID-19 related Behaviours and Attitudes*, collected by the Fields Institute in Ontario. The municipalities (cities) from where survey participants provided answers (in the clean dataset) appear as points. The Health six Regions are color-coded. Internal boundaries within certain Health Regions indicate areas that belonged to the Local Integrated Health Networks (LIHNS), the geographic areas for healthcare in Ontario before the adoption of the Health Regions.

Statistical analyses

We used a logistic regression model to estimate the probability of vaccination depending on the socio-economic factors described in Table 1, the Health Regions from Ontario indicated in Section , and the interactions between Race and Health Region, and Race and income, as previous studies have shown that socio-economic factors and their interactions are significant predictors of intent of vaccination and vaccination status³²⁻³⁴.

The model was fitted first to the clean dataset to obtain uncorrected estimates. Additionally, because we identified differences between the proportions of all the socio-economic factors included in the analysis (Table 1) and the Census data for Ontario, we used an iterative proportional fitting procedure (*raking*)³⁵ to correct the data using Census socio-economic data and Health Region population totals, in order to obtain corrected estimates from the model. Details regarding the correction can be found in the Appendix. All analyses were conducted in R 4.2.2 using the packages `survey`³⁶, `tidyverse`³⁷, and `quarto`³⁸.

Results

Survey Results

Table 2 shows the descriptive statistics from the Fields COVID-19 survey data for vaccination status and each of the covariates analyzed. The total number of entries analyzed was 3,551. Overall, 26.9% of survey respondents (958) reported not having received the first dose of the vaccine, whereas 73.1% (2,593) reported having received it. Within each socio-economic factor, respondents who reported living in a household with an income under CAD 25,000 represented 37% of the total number of entries, those within the CAD 25,000-59,999 income bracket represented 25% of the total sample, and those with an income above CAD 60,000 represented 38 % of the sample; across all income brackets, the percentage of individuals that reported having received a first dose of the vaccine was consistent, above 69%.

Within the age groups of survey respondents, the age group between 16-34 years had the highest representation in the survey responses (1,521, 42.8% of all responses). Within this age bracket, 73% of respondents indicated having received the vaccine, whereas the lowest vaccination rate was in the bracket of those 55 years of age and above, with a total of 72%. The Health Region with highest representation in the survey was Toronto, accounting for 1,324 entries (37.2%), with a vaccination rate of 72%. Regarding race/ethnicity, individuals that identified as White/Caucasian represented 1313 (37%) of all entries and had the highest vaccination uptake with 82% of them indicating to have received the COVID-19 vaccine. On the other hand, the ethnic group with the lowest number of entries in the survey was Latin American, with a total of 180, or 5% of all entries. Vaccination rates across all minority groups were below the value reported by White/Caucasians, with the lowest vaccination rate (60%) being reported by individuals that identified as Indigenous.

Table 2: Descriptive Statistics of the Fields COVID-19 Survey (by Vaccination Status)

Variable	no, N = 958	yes, N = 2,591
Income		
60000_and_above	305 (23%)	1,048 (77%)
25000_59999	253 (28%)	636 (72%)
under_25000	400 (31%)	907 (69%)
Age Group		
16_34	409 (27%)	1,111 (73%)
35_54	252 (26%)	712 (74%)
55_and_over	297 (28%)	768 (72%)
Health Region		
Toronto	371 (28%)	952 (72%)
Central	224 (28%)	580 (72%)
East	135 (23%)	448 (77%)
West	228 (27%)	611 (73%)
Race		
white_caucasian	233 (18%)	1,079 (82%)
arab_middle_eastern	76 (36%)	138 (64%)
black	114 (38%)	184 (62%)
east_asian_pacific_islander	69 (23%)	234 (77%)
indigenous	76 (40%)	115 (60%)
latin_american	69 (38%)	111 (62%)
mixed	105 (34%)	205 (66%)
other	128 (35%)	239 (65%)
south_asian	88 (24%)	286 (76%)

¹ n (%)

Multivariate Regression

Table 3 shows the results of the logistic regression models (for the uncorrected and corrected data) on vaccination status using socio-economic factors (age group, income, race), geographical areas (Health Regions) and the interactions between income and race and Health Region and race. There were no statistically significant differences in vaccination rates within the age groups from the survey, but significant odds ratios were estimated for other covariates. Within household income brackets, individuals with an income under CAD 25,000 or between CAD 25,000-59,999 had significantly lower odds of vaccination than those with an income above CAD 60,000 (ORs=0.37 and 0.59, $p=0.011$ and <0.001 , respectively). Within Race/Ethnicity, individuals who identified as Arab/Middle Eastern, Black, or Latin American, had significantly lower odds of vaccination than those in the White/Caucasian group (ORs=0.31, 0.32, 0.28, and $p=0.004$, <0.001 and 0.004 , respectively); additionally, those individuals in the Other Race/Ethnicity group (a group that included Southeast Asian, Filipino, West Asian, and Minorities Not Identified Elsewhere) had even lower odds of vaccination than the other minority groups (OR=0.22, $p<0.001$). Regarding Health Regions, individuals that reported living in the West Health Region (which comprises the regions of Waterloo and Niagara, the counties of Wellington, Essex, and Lambton, and the cities of Hamilton, Haldimand, Brant, and Chatham-Kent) had significantly higher odds of vaccination than those in the Health Region of Toronto (OR=1.55, $p=0.029$).

Moreover, statistically-significant odd ratios were determined in the case of the interaction of income and race; specifically, for individuals with a household income below CAD 25,000 who identified as Arab/Middle Eastern (OR=3.05, $p=0.013$), Black (OR=3.19, $p=0.004$), Latin American (OR=2.80, $p=0.041$), or that belonged to other minority groups (OR=4.59, $p<0.001$). Within the CAD 25,000-59,999 income bracket, individuals who identified as belonging to other racial minority groups had significantly higher odds of vaccination (OR=6.93, $p<0.001$).

For the interaction of Health Region and race, significant odds of vaccination were identified for Black individuals in the Central Health Region, which comprises the region of York, counties of Dufferin and Simcoe and the district of Muskoka (OR=0.44, $p=0.046$), and in individuals that identified as part of other racial minorities or South Asian that lived in the West Health Region (ORs=0.41, $p=0.032$ and $p=0.037$, respectively).

Table 3: Multiple Regression Analysis-Predictors of Vaccination Status

Characteristic	Uncorrected			Corrected		
	OR	95% CI	p-value	OR	95% CI	p-value
Age Group						
16_34	—	—	—	—	—	—
35_54	0.93	0.77, 1.14	0.5	0.90	0.67, 1.21	0.5
55_and_over	0.74	0.61, 0.89	0.002	0.99	0.74, 1.32	>0.9
Month						
October	—	—	—	—	—	—
December	1.16	0.91, 1.49	0.2	1.16	0.78, 1.71	0.5
November	0.95	0.81, 1.12	0.6	0.92	0.71, 1.18	0.5
Income						

(continued)

Characteristic	Uncorrected			Corrected		
	OR	95% CI	p-value	OR	95% CI	p-value
60000_and_above	—	—	—	—	—	—
25000_59999	0.59	0.41, 0.84	0.003	0.59	0.39, 0.88	0.010
under_25000	0.41	0.29, 0.58	<0.001	0.37	0.25, 0.56	<0.001
Race						
white_caucasian	—	—	—	—	—	—
arab_middle_eastern	0.24	0.12, 0.49	<0.001	0.31	0.14, 0.68	0.003
black	0.30	0.17, 0.54	<0.001	0.32	0.17, 0.60	<0.001
east_asian_pacific_islander	0.73	0.36, 1.51	0.4	1.15	0.50, 2.66	0.7
indigenous	0.38	0.17, 0.82	0.014	0.45	0.20, 1.04	0.061
latin_american	0.28	0.13, 0.59	<0.001	0.27	0.11, 0.66	0.004
mixed	0.58	0.30, 1.11	0.10	0.63	0.24, 1.62	0.3
other	0.20	0.11, 0.35	<0.001	0.22	0.12, 0.41	<0.001
south_asian	0.79	0.44, 1.43	0.4	0.90	0.48, 1.67	0.7
Health Region						
Toronto	—	—	—	—	—	—
Central	1.30	0.84, 2.00	0.2	1.47	0.92, 2.36	0.11
East	1.54	1.01, 2.34	0.044	1.41	0.90, 2.22	0.14
West	1.35	0.94, 1.93	0.10	1.54	1.04, 2.29	0.031
Income * Race						
25000_59999 * arab_middle_eastern	2.19	0.95, 5.05	0.067	1.85	0.68, 4.99	0.2
under_25000 * arab_middle_eastern	2.95	1.39, 6.24	0.005	3.08	1.27, 7.47	0.013
25000_59999 * black	1.18	0.59, 2.36	0.6	1.31	0.58, 2.98	0.5
under_25000 * black	2.84	1.46, 5.53	0.002	3.15	1.43, 6.92	0.004
25000_59999 * east_asian_pacific_islander	0.97	0.44, 2.10	>0.9	0.42	0.17, 1.06	0.066
under_25000 * east_asian_pacific_islander	1.90	0.90, 4.01	0.094	1.15	0.47, 2.84	0.8
25000_59999 * indigenous	1.77	0.72, 4.32	0.2	1.33	0.46, 3.80	0.6
under_25000 * indigenous	1.60	0.71, 3.60	0.3	1.41	0.54, 3.72	0.5
25000_59999 * latin_american	0.89	0.38, 2.10	0.8	1.25	0.45, 3.45	0.7
under_25000 * latin_american	3.10	1.34, 7.21	0.009	2.81	1.04, 7.59	0.041
25000_59999 * mixed	0.87	0.39, 1.93	0.7	0.87	0.33, 2.33	0.8
under_25000 * mixed	1.27	0.65, 2.49	0.5	1.12	0.38, 3.33	0.8
25000_59999 * other	5.50	2.43, 12.5	<0.001	6.96	2.67, 18.2	<0.001
under_25000 * other	4.09	2.27, 7.37	<0.001	4.63	2.34, 9.13	<0.001
25000_59999 * south_asian	1.14	0.55, 2.38	0.7	1.22	0.52, 2.87	0.7
under_25000 * south_asian	1.58	0.82, 3.05	0.2	2.01	0.93, 4.34	0.075
Race * Health Region						
arab_middle_eastern * Central	0.76	0.33, 1.71	0.5	0.66	0.25, 1.69	0.4
black * Central	0.48	0.23, 1.02	0.055	0.44	0.19, 0.99	0.046
east_asian_pacific_islander * Central	0.84	0.38, 1.89	0.7	0.98	0.38, 2.53	>0.9
indigenous * Central	0.60	0.24, 1.50	0.3	0.63	0.22, 1.78	0.4
latin_american * Central	0.69	0.28, 1.71	0.4	0.66	0.22, 1.93	0.4
mixed * Central	0.63	0.30, 1.35	0.2	0.73	0.24, 2.23	0.6
other * Central	0.98	0.49, 1.95	>0.9	0.81	0.36, 1.81	0.6
south_asian * Central	0.71	0.34, 1.47	0.4	0.55	0.25, 1.21	0.14
arab_middle_eastern * East	0.57	0.20, 1.63	0.3	0.43	0.13, 1.43	0.2
black * East	0.84	0.37, 1.89	0.7	0.86	0.35, 2.11	0.7
east_asian_pacific_islander * East	0.82	0.30, 2.24	0.7	0.86	0.29, 2.54	0.8
indigenous * East	0.56	0.21, 1.45	0.2	0.70	0.23, 2.11	0.5
latin_american * East	0.80	0.28, 2.25	0.7	1.03	0.31, 3.40	>0.9
mixed * East	0.70	0.31, 1.60	0.4	0.92	0.28, 3.05	0.9
other * East	0.87	0.38, 2.02	0.7	1.08	0.40, 2.91	0.9
south_asian * East	0.50	0.20, 1.24	0.14	0.51	0.19, 1.42	0.2
arab_middle_eastern * West	1.18	0.49, 2.82	0.7	1.02	0.37, 2.77	>0.9
black * West	0.78	0.36, 1.67	0.5	0.77	0.32, 1.82	0.5
east_asian_pacific_islander * West	0.54	0.24, 1.21	0.14	0.52	0.20, 1.36	0.2
indigenous * West	0.45	0.19, 1.05	0.064	0.40	0.14, 1.12	0.082
latin_american * West	0.99	0.39, 2.48	>0.9	0.96	0.33, 2.79	>0.9
mixed * West	0.53	0.25, 1.10	0.088	0.37	0.12, 1.15	0.085
other * West	0.55	0.27, 1.12	0.10	0.41	0.18, 0.92	0.032
south_asian * West	0.50	0.24, 1.08	0.078	0.41	0.18, 0.95	0.037

¹ OR = Odds Ratio, CI = Confidence Interval

Discussion

There existence of healthcare disparities within Ontario is a topic of particular interest in due to the recent change in the healthcare system of this region, which eliminated the LHIN model and adopted larger Health Regions system in late 2019 in an attempt to address the disparities of the former approach^{30,31}. In this context, analyzing COVID-19 vaccination estimates within the province is important as they can provide an initial assessment of variations that might need to be addressed by decision-makers.

Our results indicate that across the most densely populated Health Regions of Ontario, almost three quarters of the surveyed individuals reported having received the first dose of the COVID-19 vaccine (Table 2). It is worth mentioning that province-wide vaccination rates for the period of interest are somewhat different from those of the survey, particularly in the case of those 55 years of age and older, which in the survey had a vaccination rate of 72%, against a rate of 88.4% reported for the closest age bracket (50 years of age and older) reported by Public Health Ontario at the start of the period covered by the data (between September 30th, 2021 and December 12th, 2021)³⁹. In this regard, differences are to be expected because the survey represents a random sample from the population, and therefore, the sampling process is likely to cause variations between the values from the survey and province-wide estimates.

We believe that these differences did not have a significant impact in our analyses, because our results indicate that there were no significant differences in vaccination odds among the age groups considered in the survey, in agreement with the overall vaccination rates reported for Canada, which have been relatively higher when compared to other high income countries⁴⁰, and with vaccination uptake rates across different age groups presented in other studies^{19,41}. Moreover, vaccination rates within each age in the dataset are in good agreement with province-wide estimates (e.g., a rate of 95% for those with 61 years of age, Supplementary Table A-6). Additionally, when overall vaccination rates for the province are disaggregated, it can be seen that regional differences were present during the period analyzed, which can be masked by province-wide estimates. For example, the Public Health Unit of Lambton (a region within the West Health Region) and the Public Health Unit of Haliburton, Kawartha, and the Pine Ridge District (an area covered by the East Health Region) reported lower vaccination rates (78%) for those 50 years of age and older at the beginning of the period of interest, whereas other regions had higher vaccination rates³⁹. To this day, differences in vaccination rates within the province are present, because according to Public Health Ontario, as of March of 2023 some regions have less than 75% vaccination rate⁴². Overall, these reasons indicate that the estimates obtained from the data are in good agreement with the trends from the population.

We identified significant intra-provincial differences in vaccination based on socio-economic and geographical factors. First, our results show differences in odds of vaccination in individuals with a household income below CAD 60,000 and in individuals belonging to visible minority groups. Those who identified as Arab/Middle Eastern, Black, Latin American, or that belonged to a minority group not included in the survey (Southeast Asian, Filipino, West

Asian, and minority groups not identified elsewhere) had vaccination odds that were less than a third of individuals that identified as White/Caucasian (Table 3). These results are consistent with other studies that have shown lower vaccination rates in individuals that identify as part of a racial minority, or that have a low household income^{19-21,43}.

In this study, we also decided to explore the interactions between income and race and race and Health Region, as it is known that many individuals within racial minority groups perform tend to occupy certain types of occupations that fall within income brackets that have been shown to be associated with differences in vaccination uptake. In other words, we decided to explore if there were differences in vaccination within racial groups in certain income brackets and in certain the Health Regions. In this regard, it is interesting to note that although overall self-reported vaccination rates were found to be statistically significantly lower in various racial minority groups when compared to White/Caucasian individuals (Table 3), the change in odds of vaccination within certain racial groups and income strata was actually positive, in contrast to the White/Caucasian group, for which vaccination odds decreased in lower income brackets (when compared to the CAD 60,000 and over bracket, Supplementary Figure A-3). More specifically, the change in odds of vaccination increased in individuals who identified as Arab/Middle Eastern, Black, Latin American, or belonging to other minority groups with a household income below CAD 25,000, which was also true for individuals in other racial minority groups with an income between CAD 25,000-59,999 (Table 3, Supplementary Figure A-3).

This result is likely due to the fact that individuals that belong racial minority groups tend to perform occupations that have been deemed as “essential” in the context of the pandemic^{44,45}, which include occupations such as grocery store workers, gas station workers, warehouse and distribution workers, and manufacturing workers, all being occupations for which an income within the significant brackets is to be expected. In the case of Ontario, essential workers had priority for COVID-19 vaccination⁴⁶, which would explain the higher odds of vaccination for these individuals in certain income brackets, in contrast to the lower odds of vaccination for the same type of individuals with higher household income. In other words, it is possible that the type of occupation played an important role in increasing the odds of vaccination in these racial minority groups.

Additionally, significant higher vaccination odds were identified in the West Health Region when compared to the Health Region of Toronto (Table 3). The West Health Region comprises the regions of Waterloo and Niagara, the counties of Wellington, Essex and Lambton, and the cities of Hamilton, Haldimand, Brant, and Chatham-Kent. In this case, a possible rationale for the results is the fact that in the survey, about 47% of the entries for this Health Region corresponded to White/Caucasian individuals, who reported an overall 83% vaccination rate (Supplementary Table A-7). However, the interaction effect of Health Region and race was also significant in the case of individuals identifying as South Asian or other minorities not included in the survey Table 3. In this case, the results of the interaction term in the model indicate that the odds of vaccination for those within the South Asian and Other minority groups in the

West Region decreased when compared to the other Health Regions (Supplementary Figure A-4).

According to Ontario Health, 13.2% of the population in the West Health Region identifies as a visible minority, whereas 2.5% identifies as Indigenous⁴⁷. In the case of this analysis, the estimated lower odds are likely to be explained from a socio-economic perspective. In fact, 50% of the answers from this region in the survey came from the former LHINs of Hamilton Niagara Haldimand Brant, and Erie St. Clair, both which are among the regions of Ontario with the highest proportion of their population (more than 20%) in the lowest income quintile⁴⁸ (Supplementary Table A-8). Therefore, this result partly reinforces the well-known existing association between low vaccination rates and income, but it additionally indicates that there were intra-regional differences in vaccination. Interestingly, a disproportionate number of COVID-19 cases and low vaccination rate (under 50%) have been previously reported in the South Asian community of Ontario⁴⁹; in this regard, our result provides additional context by showing that within the South Asian community, there were differences in vaccination uptake across Ontario. Moreover, because significant lower odds of vaccination were also identified other minority groups, this provides a rationale for future studies that explore how vaccination uptake varies across different minority groups within Ontario and other Canadian provinces.

There are some limitations to the present study. First, the data collection design, which allowed respondents to withdraw from the survey at any point, resulted in a high number of unique entries in the survey with multiple missing answers. Because we focused on entries that had complete observations in the covariates of interest for our analysis, it is possible that some information was not considered by excluding observations that had information in other variables (such as work from home, or number of persons in the household). However, we attempted to minimize this possibility by correcting the dataset using information from the Census. More granular corrections, which for example could be based on demographic information by municipality, could be used in the future to obtain a more accurate approximation to the population totals of the province. Moreover, our analysis did not consider the North West and North East Health Regions, due to the low number of entries from these areas in the survey (Figure 1). Although low representation from these areas is based on the fact that these regions only account for 5% of the total population of Ontario, these regions are the home to more than 100,000 individuals that identify as Indigenous⁴⁷, a minority group that has historically suffered from reduced access to health care and discrimination¹⁷. Therefore, there is a need for additional studies that focus on these low-populated Health Regions in Ontario where disparities in vaccination might be significant and understudied.

The results in this study are based on self-reported data, where there is a risk that biased values are reported. Despite this, because in the context of COVID-19 it has been shown that good agreement exists between self-reported and documented vaccination status⁵⁰, and therefore, the effect of self-reported bias is likely to not be significant in our analyses. Finally, it is likely that there have been differences in vaccination across the province as more doses of the vaccine were administered and as successive variants emerged. Because this study focused only on vaccination status regarding the first dose of the vaccine within a relatively short

time window, it can only provide a snapshot of the societal dynamics behind the pandemic. Nonetheless, the results presented here can serve as a starting point to motivate the collection of robust longitudinal data that can be used to quantify geographical and temporal differences within vulnerable segments of the population, and that can be used to inform the development of adequate public health policies within the province of Ontario or across other provinces that aim to minimize disparities in health access.

Conclusion

This study explored differences in COVID-19 vaccination across the province of Ontario between late 2021 and early 2022 by taking into consideration socio-economic factors, such as income and race, their interactions, and the Health Regions within the province. Our results show that, during the period analyzed, significant differences in vaccination existed across different visible minority groups, income brackets, and Health Regions, showing intra-provincial disparities in vaccine uptake. As the COVID-19 continues around the world, it is important that future public policies take into consideration how to adequately reach individuals within minority groups that live across geographical areas where less probabilities of being vaccinated are likely. At the moment, this is an ongoing issue that needs to be addressed to ensure a more homogeneous outcome from the pandemic.

References

1. Davis CJ, Golding M, McKay R. Efficacy information influences intention to take COVID-19 vaccine. *British Journal of Health Psychology*. 2022;27(2):300-319. doi:<https://doi.org/10.1111/bjhp.12546>
2. Bogoch II, Halani S. COVID-19 vaccines: A geographic, social and policy view of vaccination efforts in Ontario, Canada. *Cambridge Journal of Regions, Economy and Society*. Published online November 2022. doi:[10.1093/cjres/rsac043](https://doi.org/10.1093/cjres/rsac043)
3. Tanne JH. Covid-19: FDA panel votes to authorise pfizer BioNTech vaccine. *BMJ*. Published online December 2020:m4799. doi:[10.1136/bmj.m4799](https://doi.org/10.1136/bmj.m4799)
4. Microbe TL. COVID-19 vaccines: The pandemic will not end overnight. *The Lancet Microbe*. 2021;2(1):e1. doi:[10.1016/s2666-5247\(20\)30226-3](https://doi.org/10.1016/s2666-5247(20)30226-3)
5. Watson OJ, Barnsley G, Toor J, Hogan AB, Winskill P, Ghani AC. Global impact of the first year of COVID-19 vaccination: A mathematical modelling study. *The Lancet Infectious Diseases*. 2022;22(9):1293-1302. doi:[10.1016/s1473-3099\(22\)00320-6](https://doi.org/10.1016/s1473-3099(22)00320-6)
6. Kayser V, Ramzan I. Vaccines and vaccination: History and emerging issues. *Human Vaccines & Immunotherapeutics*. 2021;17(12):5255-5268. doi:[10.1080/21645515.2021.1977057](https://doi.org/10.1080/21645515.2021.1977057)

7. Li Q, Wang J, Tang Y, Lu H. Next-generation COVID-19 vaccines: Opportunities for vaccine development and challenges in tackling COVID-19. *Drug Discoveries & Therapeutics*. 2021;15(3):118-123. doi:[10.5582/ddt.2021.0105](https://doi.org/10.5582/ddt.2021.0105)
8. Gerretsen P, Kim J, Caravaggio F, et al. Individual determinants of COVID-19 vaccine hesitancy. Inbaraj LR, ed. *PLOS ONE*. 2021;16(11):e0258462. doi:[10.1371/journal.pone.0258462](https://doi.org/10.1371/journal.pone.0258462)
9. Yamey G, Garcia P, Hassan F, et al. It is not too late to achieve global covid-19 vaccine equity. *BMJ*. Published online March 2022:e070650. doi:[10.1136/bmj-2022-070650](https://doi.org/10.1136/bmj-2022-070650)
10. Nafilyan V, Dolby T, Razieh C, et al. Sociodemographic inequality in COVID-19 vaccination coverage among elderly adults in england: A national linked data study. *BMJ Open*. 2021;11(7):e053402. doi:[10.1136/bmjopen-2021-053402](https://doi.org/10.1136/bmjopen-2021-053402)
11. Malik AA, McFadden SM, Elharake J, Omer SB. Determinants of COVID-19 vaccine acceptance in the US. *EClinicalMedicine*. 2020;26:100495. doi:[10.1016/j.eclinm.2020.100495](https://doi.org/10.1016/j.eclinm.2020.100495)
12. Willis DE, Andersen JA, Bryant-Moore K, et al. COVID-19 vaccine hesitancy: Race/ethnicity, trust, and fear. *Clinical and Translational Science*. 2021;14(6):2200-2207. doi:[10.1111/cts.13077](https://doi.org/10.1111/cts.13077)
13. Skirrow H, Barnett S, Bell S, et al. Women's views on accepting COVID-19 vaccination during and after pregnancy, and for their babies: A multi-methods study in the UK. *BMC Pregnancy and Childbirth*. 2022;22(1). doi:[10.1186/s12884-021-04321-3](https://doi.org/10.1186/s12884-021-04321-3)
14. Stoler J, Enders AM, Klostad CA, Uscinski JE. The limits of medical trust in mitigating COVID-19 vaccine hesitancy among black americans. *Journal of General Internal Medicine*. 2021;36(11):3629-3631. doi:[10.1007/s11606-021-06743-3](https://doi.org/10.1007/s11606-021-06743-3)
15. Khubchandani J, Sharma S, Price JH, Wiblshauer MJ, Sharma M, Webb FJ. COVID-19 vaccination hesitancy in the united states: A rapid national assessment. *Journal of Community Health*. 2021;46(2):270-277. doi:[10.1007/s10900-020-00958-x](https://doi.org/10.1007/s10900-020-00958-x)
16. Bogart LM, Ojikutu BO, Tyagi K, et al. COVID-19 related medical mistrust, health impacts, and potential vaccine hesitancy among black americans living with HIV. *JAIDS Journal of Acquired Immune Deficiency Syndromes*. 2021;86(2):200-207. doi:[10.1097/qai.0000000000002570](https://doi.org/10.1097/qai.0000000000002570)
17. Mosby I, Swidrovich J. Medical experimentation and the roots of COVID-19 vaccine hesitancy among indigenous peoples in canada. *Canadian Medical Association Journal*. 2021;193(11):E381-E383. doi:[10.1503/cmaj.210112](https://doi.org/10.1503/cmaj.210112)
18. Freeman D, Loe BS, Chadwick A, et al. COVID-19 vaccine hesitancy in the UK: The oxford coronavirus explanations, attitudes, and narratives survey (oceans) II. *Psychological Medicine*. 2020;52(14):3127-3141. doi:[10.1017/s0033291720005188](https://doi.org/10.1017/s0033291720005188)
19. Guay M, Maquiling A, Chen R, et al. Measuring inequalities in COVID-19 vaccination uptake and intent: Results from the canadian community health survey 2021. *BMC Public Health*. 2022;22(1). doi:[10.1186/s12889-022-14090-z](https://doi.org/10.1186/s12889-022-14090-z)

- 369 20. Muhajarine N, Adeyinka DA, McCutcheon J, Green KL, Fahlman M, Kallio N. COVID-19 vaccine hesitancy and refusal and associated factors in an adult population in saskatchewan, canada: Evidence from predictive modelling. Gesser-Edelsburg A, ed. *PLOS ONE*. 2021;16(11):e0259513. doi:[10.1371/journal.pone.0259513](https://doi.org/10.1371/journal.pone.0259513)
- 370 21. Hussain B, Latif A, Timmons S, Nkhoma K, Nellums LB. Overcoming COVID-19 vaccine hesitancy among ethnic minorities: A systematic review of UK studies. *Vaccine*. 2022;40(25):3413-3432. doi:[10.1016/j.vaccine.2022.04.030](https://doi.org/10.1016/j.vaccine.2022.04.030)
- 372 22. Nguyen KH, Nguyen K, Corlin L, Allen JD, Chung M. Changes in COVID-19 vaccination receipt and intention to vaccinate by socioeconomic characteristics and geographic area, united states, january 6 – march 29, 2021. *Annals of Medicine*. 2021;53(1):1419-1428. doi:[10.1080/07853890.2021.1957998](https://doi.org/10.1080/07853890.2021.1957998)
- 374 23. Mollalo A, Tatar M. Spatial modeling of COVID-19 vaccine hesitancy in the united states. *International Journal of Environmental Research and Public Health*. 2021;18(18):9488. doi:[10.3390/ijerph18189488](https://doi.org/10.3390/ijerph18189488)
- 376 24. Yang TC, Matthews SA, Sun F. Multiscale dimensions of spatial process: COVID-19 fully vaccinated rates in u.s. counties. *American Journal of Preventive Medicine*. 2022;63(6):954-961. doi:[10.1016/j.amepre.2022.06.006](https://doi.org/10.1016/j.amepre.2022.06.006)
- 378 25. Tiu A, Susswein Z, Merritt A, Bansal S. Characterizing the spatiotemporal heterogeneity of the COVID-19 vaccination landscape. *American Journal of Epidemiology*. 2022;191(10):1792-1802. doi:[10.1093/aje/kwac080](https://doi.org/10.1093/aje/kwac080)
- 380 26. Bhuiyan MAN, Davis TC, Arnold CL, et al. Using the social vulnerability index to assess COVID-19 vaccine uptake in louisiana. *GeoJournal*. Published online December 2022. doi:[10.1007/s10708-022-10802-5](https://doi.org/10.1007/s10708-022-10802-5)
- 382 27. Wood AJ, MacKintosh AM, Stead M, Kao RR. Predicting future spatial patterns in COVID-19 booster vaccine uptake. Published online September 2022. doi:[10.1101/2022.08.30.22279415](https://doi.org/10.1101/2022.08.30.22279415)
- 384 28. Choi KH, Denice PA, Ramaj S. Vaccine and COVID-19 trajectories. *Socius: Sociological Research for a Dynamic World*. 2021;7:237802312110529. doi:[10.1177/23780231211052946](https://doi.org/10.1177/23780231211052946)
- 386 29. McKinnon B, Quach C, Dubé Ève, Nguyen CT, Zinszer K. Social inequalities in COVID-19 vaccine acceptance and uptake for children and adolescents in montreal, canada. *Vaccine*. 2021;39(49):7140-7145. doi:[10.1016/j.vaccine.2021.10.077](https://doi.org/10.1016/j.vaccine.2021.10.077)
- 388 30. Tsasis P, Evans JM, Owen S. Reframing the challenges to integrated care: A complex-adaptive systems perspective. *International Journal of Integrated Care*. 2012;12(5). doi:[10.5334/ijic.843](https://doi.org/10.5334/ijic.843)
- 390 31. Dong L, Sahu R, Black R. Governance in the transformational journey toward integrated healthcare: The case of ontario. *Journal of Information Technology Teaching Cases*. Published online December 2022:204388692211473. doi:[10.1177/20438869221147313](https://doi.org/10.1177/20438869221147313)
- 392

32. Nguyen KH, Anneser E, Toppo A, Allen JD, Parott JS, Corlin L. Disparities in national and state estimates of COVID-19 vaccination receipt and intent to vaccinate by race/ethnicity, income, and age group among adults ≥ 18 years, united states. *Vaccine*. 2022;40(1):107-113. doi:[10.1016/j.vaccine.2021.11.040](https://doi.org/10.1016/j.vaccine.2021.11.040)
33. Shih SF, Wagner AL, Masters NB, Prosser LA, Lu Y, Zikmund-Fisher BJ. Vaccine hesitancy and rejection of a vaccine for the novel coronavirus in the united states. *Frontiers in Immunology*. 2021;12. doi:[10.3389/fimmu.2021.558270](https://doi.org/10.3389/fimmu.2021.558270)
34. Cénat JM, Noorishad PG, Farahi SMMM, et al. Prevalence and factors related to COVID-19 vaccine hesitancy and unwillingness in canada: A systematic review and meta-analysis. *Journal of Medical Virology*. 2022;95(1). doi:[10.1002/jmv.28156](https://doi.org/10.1002/jmv.28156)
35. Deming WE, Stephan FF. On a least squares adjustment of a sampled frequency table when the expected marginal totals are known. *The Annals of Mathematical Statistics*. 1940;11(4):427-444. doi:[10.1214/aoms/1177731829](https://doi.org/10.1214/aoms/1177731829)
36. Lumley T. *Complex Surveys*. John Wiley & Sons; 2011.
37. Wickham H, Averick M, Bryan J, et al. Welcome to the tidyverse. *Journal of Open Source Software*. 2019;4(43):1686. doi:[10.21105/joss.01686](https://doi.org/10.21105/joss.01686)
38. Allaire J. *Quarto: R Interface to 'Quarto' Markdown Publishing System.*; 2022. <https://CRAN.R-project.org/package=quarto>
39. Ontario COVID-19 Data Tool. Accessed February 27, 2023. <https://www.publichealthontario.ca/en/data-and-analysis/infectious-disease/covid-19-data-surveillance/covid-19-data-tool?tab=vaccine>
40. Dubé E, Gagnon D, MacDonald N. Between persuasion and compulsion: The case of COVID-19 vaccination in canada. *Vaccine*. 2022;40(29):3923-3926. doi:[10.1016/j.vaccine.2022.05.053](https://doi.org/10.1016/j.vaccine.2022.05.053)
41. MacDonald NE, Comeau J, Dubé Ève, et al. Royal society of canada COVID-19 report: Enhancing COVID-19 vaccine acceptance in canada. Blais JM, ed. *FACETS*. 2021;6:1184-1246. doi:[10.1139/facets-2021-0037](https://doi.org/10.1139/facets-2021-0037)
42. Ontario COVID-19 Data Tool. Accessed March 24, 2023. <https://www.publichealthontario.ca/en/data-and-analysis/infectious-disease/covid-19-data-surveillance/covid-19-data-tool?tab=maps>
43. Carter MA, Biro S, Maier A, Shingler C, Guan TH. COVID-19 vaccine uptake in southeastern ontario, canada: Monitoring and addressing health inequities. *Journal of Public Health Management and Practice*. 2022;28(6):615-623. doi:[10.1097/phh.0000000000001565](https://doi.org/10.1097/phh.0000000000001565)
44. Hawkins D. Differential occupational risk for COVID-19 and other infection exposure according to race and ethnicity. *American Journal of Industrial Medicine*. 2020;63(9):817-820. doi:[10.1002/ajim.23145](https://doi.org/10.1002/ajim.23145)

- 419 45. Côté D, Durant S, MacEachen E, et al. A rapid scoping review of COVID-19 and vulnerable workers: Intersecting occupational and public health issues. *American Journal of Industrial Medicine*. 2021;64(7):551-566. doi:[10.1002/ajim.23256](https://doi.org/10.1002/ajim.23256)
- 420
- 421 46. Mishra S, Stall NM, Ma H, et al. *A Vaccination Strategy for Ontario COVID-19 Hotspots and Essential Workers*. Ontario COVID-19 Science Advisory Table; 2021. doi:[10.47326/ocsat.2021.02.26.1.0](https://doi.org/10.47326/ocsat.2021.02.26.1.0)
- 422
- 423 47. *Annual Business Plan 2022/23*. Ontario Health; https://www.ontariohealth.ca/sites/ontariohealth/files/2022-05/OHBusinessPlan22_23.pdf; 2022.
- 424
- 425 48. Buajitti E, Watson T, Kornas K, Bornbaum C, Henry D, Rosella LC. Ontario atlas of adult mortality, 1992-2015: Trends in Local Health Integration Networks. Published online 2018. <https://tspace.library.utoronto.ca/handle/1807/82836>
- 426
- 427 49. Anand SS, Arnold C, Bangdiwala SI, et al. Seropositivity and risk factors for SARS-CoV-2 infection in a south asian community in ontario: A cross-sectional analysis of a prospective cohort study. *CMAJ Open*. 2022;10(3):E599-E609. doi:[10.9778/cmajo.20220031](https://doi.org/10.9778/cmajo.20220031)
- 428
- 429 50. Stephenson M, Olson SM, Self WH, et al. Ascertainment of vaccination status by self-report versus source documentation: Impact on measuring COVID-19 vaccine effectiveness. *Influenza and Other Respiratory Viruses*. 2022;16(6):1101-1111. doi:[10.1111/irv.13023](https://doi.org/10.1111/irv.13023)
- 430