- Scoping Review Protocol: Statistical Models for Longitudinal Data
- in Health and Biomedical Research: Current State, Challenges,

and Opportunities

Ariel I. Mundo Ortiz

5 2022-09-07

6 Table of contents

7	1	Notes	2
8	2	Registration	3
9	3	Author Contributions	3
10	4	Amendements	3
11	5	Support	3
12		5.1 Sources	3
13	6	Introduction	3
14		6.1 Rationale	3
15	7	Objectives	5
16	8	Review Question	5

17	9	Met	m thods	6
18		9.1	Types of Studies	6
19		9.2	Eligibility Criteria	6
20			9.2.1 For the Application of Modern Statistical Models on Longitudinal Biomedical/Health	
21			Data (Aims 1a and 1b)	6
22			9.2.2 For Methods on Longitudinal Data (Aim 2)	6
23		9.3	Information Sources	7
24		9.4	Search Strategy	7
25			9.4.1 For the Application of Modern Models on Longitudinal Biomedical/Health Data $$	7
26			9.4.2 For Methods on Longitudinal Data	9
27		9.5	Data Collection and Analysis	10
28			9.5.1 Selection Process and Data Management	10
29			9.5.2 Data Collection Process	11
30		9.6	Data Items	11
31		9.7	Risk of Bias in Individual Studies	12
32		9.8	Data Synthesis	12
33		9.9	Meta-Biases	12
34	10	Refe	erences	12

35 1 Notes

As of Sept 7, 2022 this document follows the structure recommended by PRISMA-P

 $^{37} \quad \text{https://prisma-statement.org/documents/PRISMA-P-checklist.pdf}$

38 2 Registration

- 39 This section will be populated with the registration number and registry name once the protocol is submitted
- 40 for peer review.

41 3 Author Contributions

- AM: Writing, query design, data extraction and analysis . . .
- 43 Other authors to add later

4 Amendements

- 45 Protocol amendments resulting from peer review will be indicated in this section indicating the date of each
- 46 amendment.

5 Support

This section will indicate the sources of financial or other support for the review

5.1 Sources

50 6 Introduction

51 6.1 Rationale

- 52 Longitudinal studies are frequently used in the health sciences (biomedical research, epidemiology, public
- bealth, among others) as they allow to examine how the temporal effect of a treatment or an intervention,
- 54 in contrast to a cross-sectional study, which only allows to examine the effect of the intervention at a single
- 55 time point. When compared their cross-sectional counterparts, longitudinal studies allow for increased
- statistical power and more cost efficient strategies^{1,2}. However, the statistical analysis of longitudinal data
- 57 requires to take into consideration factors such as data missingness, correlation, and non-linear trends,

which do not occur on cross-sectional data^{3,4}. In other words, there is an "analytic cost" associated with the increased complexity of longitudinal data².

This additional layer of complexity has led to a problem of model misspecification in the statistical analysis of the data (i.e., the use of a statistical model that is not coherent with the data), which has been reported to occur in many fields, including the health sciences⁵. For example, in a landmark study Liu et al. showed that in a subset of papers in the biomedical sciences, the most popular model used to analyze longitudinal data was the analysis of variance (ANOVA, an approach that fails to take into account the correlation between measures over time), and that only 18% of the studies analyzed used models intended for longitudinal analysis while checking that the assumptions of the model were satisfied by the data⁶.

Historically, the repeated measures ANOVA (rm-ANOVA, a statistical model for longitudinal data) has been the preferred method in the health sciences to analyze longitudinal data, despite the fact that the multiple assumptions required by this model are frequently not satisfied by the data collected in longitudinal studies⁴. On the other hand, the last 30 years have seen incredible progress in the field of Statistics with the development of statistical models for longitudinal data that relax the assumptions of rm-ANOVA. Linear mixed models, generalized additive mixed models, and generalized estimating equations are among these modern statistical models developed for longitudinal data⁷⁻¹¹. From these statistical methods, linear mixed models and generalized estimating equations are the two classes of models that have been frequently applied to analyze longitudinal data in the health sciences during the last decade¹²⁻¹⁴.

However, modern statistical methods that are suited to analyze longitudinal data have been the exception rather than the norm in the health sciences. In 2001, a study reported that only 30% of the clinical trials analyzed used linear mixed models to analyze their results, and that the preferred method of analysis continued to be rm-ANOVA¹⁵ (in comparison, McCullagh and Nelder's seminal book on the generalized linear model (GLM) was published in 1989¹⁶, and there was ongoing work on the extension of the GLM framework to the mixed model case by 1993¹⁷). Apart from the aforementioned study, there are not recent papers that examine the use of modern statistical methods for longitudinal data in the health sciences. Such information is critical to understand if the use of these methods has increased or decreased in the field over the last 20 years, and the reasons behind such changes.

Additionally, the reproducibility crisis is an ongoing issue in the health sciences^{18,19}, a major component of it being the misuse and lack of reproducibility of statistical analyses^{20,21}. Despite the fact that the landscape of statistical software has vastly increased in the last decade with many statistical computational tools now available to researchers, reproducibility standards vary between each computational tool²². Furthermore, there is still high variability in the amount of statistical reporting across journals²³. Understanding what statistical computational tools are used nowadays by researchers in the health sciences can provide an assessment of the advances in the field towards research reproducibility, while identifying limitations that might still be in place.

7 Objectives

94 This study aims to:

101

102

- Identify the different statistical models for longitudinal data that are used in the health sciences in order to measure the current extent in the adoption of modern statistical methods by the field (Aim 1a)
- Summarize the computational tools used by researchers in the health sciences to statistically analyze longitudinal data to understand the current status of the field with regards to reproducibility. (Aim 1b)
 - List statistical methods for longitudinal data developed within the last decade in order to showcase newer methods that may be applicable for longitudinal data in a biomedical/health context. (Aim 2)

103 8 Review Question

- What are the statistical methods used in biomedical/health sciences research?
- Has the use of modern statistical methods increased in the field during the last 20 years?
- What computational tools are most commonly used by researchers to analyze longitudinal data, and how in turn this affects reproducibility?
- What are most recent statistical methods developed for longitudinal data, and how can they be applied in the health sciences?

$_{110}$ 9 Methods

111 9.1 Types of Studies

For all the study aims, studies included in the analysis correspond to peer-reviewed publications in English.

113 9.2 Eligibility Criteria

9.2.1 For the Application of Modern Statistical Models on Longitudinal Biomedical/Health

Data (Aims 1a and 1b)

9.2.1.1 Inclusion Criteria

- Articles that:
- Are written in English
- Belong to the biomedical/health sciences fields
- Describe the collection and analysis of continuous or discrete longitudinal data
- Indicate the statistical model used to analyze the data
- Report the results of their statistical analyses

9.2.1.2 Exclusion Criteria

- Cross-sectional studies
- Tutorials that present the application of existing statistical methods to biomedical/health data
- Reviews, meta-analyses, or systematic reviews on existing statistical methods for longitudinal data
- Studies that use only descriptive statistics to summarize/analyze the data
- Studies that collect and analyze categorical data

9.2.2 For Methods on Longitudinal Data (Aim 2)

9.2.2.1 Inclusion Criteria

- Articles that:
- Are written in English
- Present new methodologies or significant improvements to existing methods for longitudinal data

9.2.2.2 Exclusion Criteria

- Systematic reviews, meta-analyses, or reviews of statistical methods for longitudinal data
- Tutorials that present the application of existing statistical methods to biomedical/health longitudinal data

9.3 Information Sources

139 Studies will be retrieved from PubMed and Web of Science.

9.4 Search Strategy

9.4.1 For the Application of Modern Models on Longitudinal Biomedical/Health Data

142 9.4.1.1 PubMed

- Note: I removed the query I had with specific names of specific statistical models (as the results were too broad and non-specific).
- 9.4.1.1.1 Query 2:
- (biomedical OR health) AND ((repeated measures) OR (longitudinal study) OR longitudinal data) AND
- ((statistical analyses) OR (statistical analysis)) NOT ((review) OR (meta analysis)) when you put NOT
- that might exclude papers with Review and meta analysis as words in the paper
- 149 Hits: 12,617
- 150 Response to comment: I followed your advice and re-wrote the query, but now I was sure to exclude meta-
- analysis and review papers by type of publication, and papers that are classified in PubMed as devoted to
- 152 Statistical methodologies (not about application of methods to longitudinal data):

153 9.4.1.1.2 Query 3 (modified Query 2):

- biomedical OR health) AND ((repeated measures) OR (longitudinal study) OR longitudinal data)

 AND ((statistical analyses) OR (statistical analysis)) NOT (Review[Publication Type] OR Meta

 analy*[Publication Type]) NOT ("Statistics as Topic/methods"[Majr] OR "Statistics as Topic/statistics

 and numerical data"[Majr] OR "Models, Statistical"[Mesh] OR "Research Design"[Mesh])
- 158 Hits: 10,948
- 159 Comments: This query is better than Query 2.
- Papers from this query appear to be good. The query catches many papers from psychology and psychiatry,
- but the ones I checked did said used linear mixed models or regression in their analyses. A few of them still
- deal with methodologies, but seems to be much more less than in the previous query.

9.4.1.2 Web of Science

164 9.4.1.2.1 Query 1:

- WC=(biom* OR health OR allergy OR cell biology OR cardio* OR hematology OR immunology OR life sciences biomedicine other topics OR medical informatics OR neuro* OR oncology OR pharmacology OR radiology, nuclear medicine & medical imaging OR research & experimental medicine OR substance abuse OR optics) AND AK=(longitudinal study OR repeated measures study) NOT ALL=(review OR meta analysis) NOT AK=(model* AND study design) NOT KP=(model)
- 170 Hits: 4.716
- when you put NOT that might exclude papers with Review and meta analysis as words in the paper
- Remove the NOT and check the suggestions for the PubMed section

9.4.1.2.2 Query 2 (Updated Query 1):

- WC=(biom* OR health OR allergy OR cell biology OR cardio* OR hematology OR immunology OR life sciences biomedicine other topics OR medical informatics OR neuro* OR oncology OR pharmacology OR radiology, nuclear medicine & medical imaging OR research & experimental medicine OR substance abuse OR optics) AND AK=(longitudinal study OR repeated measures study) NOT AK=(model* AND study design) NOT KP=(model) NOT TI=(review OR meta analy*)
- 179 Comments: Updated this query based on your comments.

180 Hits: 4,595

I updated this query based on your comments. Now I used NOT but focused on the title (hence, TI) so articles with "review" or "meta analysis" are excluded. An additional filter is that Web of Science allows to select the type of publications so I chose "Article", and this would remove review papers, editorials or other materials that are not research papers.

9.4.2 For Methods on Longitudinal Data

you need to describe this part in your objectives

I re-wrote the query here based on your comments. After reading your comments and the queries

I created, I realized that the filtering was not correct. I wrote a new query that I believe better represents
the terms we want to look at.

190 9.4.2.0.1 Query 1:

("Models, Statistical" [Mesh] OR "Biostatistics/methods" [Mesh]) AND ("Longitudinal Studies" [Mesh])

NOT (Review [Publication Type] OR Meta Analys* [Publication Type] OR "editorial" [Publication Type])

NOT ("survival" [Title/abstract]) NOT ("tutorial" [title/abstract] OR "orientation" [title/abstract]) NOT

(Humans [Mesh] OR Adolescent [Mesh] OR Animals [Mesh])

195 Hits: 142

196 Comments:

The rationale for this query is to find papers that have been labeled as dealing with models in Biostatistics or Statistics, that deal with longitudinal data, but excluding reviews, editorials, meta analysis, tutorials (that show how to implement an existing model, but not the development of a new model). Additionally, I added the "humans", "adolescent", and "animal" labels to exclude, because there are **many** papers that have all the previous labels but that are devoted to comparing methods, or about studies with animal or clinical data (without those last filters for humans, adolescents, and animals the hits are 14,702).

Again, papers that describe the development of new methods for longitudinal data should be relatively few when compared to papers that deal with application, and that is why to me the result of the query (142 hits) makes sense. I did take a look at the papers of this query and all of them seem to be about models, which is what we want.

207 9.4.2.1 Web of Science

208 **9.4.2.1.1** Query 1:

- AK=((longitudinal OR repeated measures sout OR longitudinal data, already included in longitudinal) AND
- 210 (model OR design OR method)) NOT ALL=(review OR meta analysis) NOT ALL=(survival analysis)
- 211 Hits: 3,071
- 212 Comments: This query seems to be good.
- This query returns papers that deal with methods for longitudinal analysis. Two additional options can be
- selected: 1) include only articles (which reduces the number of hits to 2,936 as book chapters and editorials
- are omitted) and 2) select from the 01/01/2000 until today (which could be reasonable as the increment of
- models has occurred during the last two decades. This option reduces the number to papers to 2,849).
- we will see when the queries is revised what is the best strategy

218 9.4.2.1.2 Query 2 (updated Query 1):

- 219 I updated the query based on your comments. The query is more specific now, please see below.
- 220 AK=((longitudinal OR repeated measures) AND (model OR design OR method)) NOT TI=(review OR
- 221 meta analysis) NOT AK=(survival analysis) AND SU=(stat* OR MATH*) NOT AB=(tutorial)
- 222 Hits: 1,978
- 223 Comments:
- This query looks for papers where the authors used the keywords longitudinal or repeated measures, and
- 225 model, or design or method; excludes papers that have "review" or "meta analysis" in the title, excludes
- those that have keywords of "survival analysis", and searches papers where the main subject has been
- tagged as statistics or math. Finally, papers that have the word "tutorial" in the abstract are excluded as
- well.

9.5 Data Collection and Analysis

9.5.1 Selection Process and Data Management

- Two reviewers will independently analyze the database search results and pre-screen articles based on ti-
- tle and abstract content following the aforementioned inclusion/exclusion criteria. Manuscripts from the

database(s) search will be stored in the Covidence platform, where duplicated entries will be removed. For articles where pre-screening inclusion (or exclusion) is unclear based on title and abstract analysis, full-text review will be used to make a decision following review by a third independent reviewer. Manuscripts included after title and abstract pre-screening will be further screening by two reviewers that will independently examine the full text of each article.

9.5.2 Data Collection Process

Pilot forms (electronic spreadsheets) will be tested using a representative sample of the studies to be reviewed (~100 studies). Information in the forms will be independently included by each reviewer. The forms will be updated (if needed), after the pilot test by consensus between the reviewers.

Information obtained from each study (statistical method used, software, etc.) will be tabulated independently by the reviewers in an electronic spreadsheet.

9.6 Data Items

- 245 Aims 1a and 1b:
- Statistical method used
- Sub-area of application (oncology, psychology, public health, etc)
- Computational tool used
- Congruence between statistical method used and the data
- Year of publication
- 251 Aim 2:
- Statistical method reported
- Assumptions of the model
- Computational tools available for its implementation
- Year of publication

9.7 Risk of Bias in Individual Studies

257 N/A

9.8 Data Synthesis

- 259 The data from the results of each included study will be extracted into electronic spreadsheets. Summary
- measures for Aims 1a and 1b include plots (pie, bar, etc) to show the relative use of each statistical method
- 261 reported, computational tool, and congruence between statistical method and the data. Each plot will be
- 262 segmented by year to show trends over time.
- For Aim 2, a table will be created where statistical method, year of publication, assumptions of the model,
- 264 and applicability to health data is reported.

9.9 Meta-Biases

266 N/A

$_{^{267}}$ 10 References

- Edwards LJ. Modern statistical techniques for the analysis of longitudinal data in biomedical research. Pediatric Pulmonology. 2000;30(4):330-344. doi:https://doi.org/10.1002/1099-0496(200010)
 30:4%3C330::AID-PPUL10%3E3.0.CO;2-D
- Zeger SL, Liang K-Y. An overview of methods for the analysis of longitudinal data. <u>Statistics in Medicine</u>. 1992;11(14-15):1825-1839. doi:https://doi.org/10.1002/sim.4780111406
- 272 3. Caruana EJ, Roman M, Hernández-Sánchez J, Solli P. Longitudinal studies. <u>Journal of Thoracic</u>

 Disease. 2015;7(11):E537-40.
- Mundo AI, Tipton JR, Muldoon TJ. Generalized additive models to analyze nonlinear trends in biomedical longitudinal data using r: Beyond repeated measures ANOVA and linear mixed models.
 Statistics in Medicine. Published online July 2022.
- Thiese MS, Arnold ZC, Walker SD. The misuse and abuse of statistics in biomedical research.

 Biochem Med (Zagreb). 2015;25(1):5-11.

- Liu C, Cripe TP, Kim M-O. Statistical issues in longitudinal data analysis for treatment efficacy studies in the biomedical sciences. <u>Molecular Therapy</u>. 2010;18(9):1724-1730. doi:https://doi.org/10.1038/mt.2010.127
- 7. Linear mixed-effects models: Basic concepts and examples. In: Mixed-Effects Models in s and

 s-PLUS. Springer New York; 2000:3-56. doi:10.1007/0-387-22747-4_1
- 282 8. Jiang J, Nguyen T. <u>Linear and Generalized Linear Mixed Models and Their Applications</u>. 2nd ed. Springer; 2021.
- 284 9. Hastie TJ. Statistical Models in S. (Chambers JM, Hastie TJ, eds.). Routledge; 2017.
- Rosa GJM, Gianola D, Padovani CR. Bayesian longitudinal data analysis with mixed models and thick-tailed distributions using MCMC. <u>Journal of Applied Statistics</u>. 2004;31(7):855-873.
- Ballinger GA. Using generalized estimating equations for longitudinal data analysis. Organizational

 Research Methods. 2004;7(2):127-150.
- ²⁹⁰ 12. Wang M. Generalized estimating equations in longitudinal data analysis: A review and recent developments. <u>Advances in Statistics</u>. 2014;2014:1-11.
- Tian Q, Qin L, Zhu W, Xiong S, Wu B. Analysis of factors contributing to postoperative body weight change in patients with gastric cancer: Based on generalized estimation equation. PeerJ. 2020;8(e9390):e9390.
- ²⁹⁴ 14. Şevik M, Doğan M. Epidemiological and molecular studies on lumpy skin disease outbreaks in turkey during 2014-2015. Transboundary and Emerging Diseases. 2017;64(4):1268-1279.
- 296 15. Gueorguieva R, Krystal JH. Move Over ANOVA: Progress in Analyzing Repeated-Measures Data and Reflection in Papers Published in the Archives of General Psychiatry. <u>Archives of General Psychiatry</u>. 2004;61(3):310-317. doi:10.1001/archpsyc.61.3.310
- ²⁹⁸ 16. McCullagh P, Nelder JA. Generalized Linear Models. Routledge; 2019.

299

- 300 17. Breslow NE, Clayton DG. Approximate inference in generalized linear mixed models. <u>Journal of the</u>

 American Statistical Association. 1993;88(421):9-25. doi:10.1080/01621459.1993.10594284
- Jarvis MF, Williams M. Irreproducibility in preclinical biomedical research: Perceptions, uncertainties, and knowledge gaps. <u>Trends in Pharmacological Sciences</u>. 2016;37(4):290-302. doi:https://doi.org/10.1016/j.tips.2015.12.001

- ³⁰⁴ 19. Turkiewicz A, Luta G, Hughes HV, Ranstam J. Statistical mistakes and how to avoid them lessons learned from the reproducibility crisis. <u>Osteoarthritis and Cartilage</u>. 2018;26(11):1409-1411. doi:10.1016/j.joca.2018.07.017
- Gosselin R-D. Statistical analysis must improve to address the reproducibility crisis: The ACcess to transparent statistics (ACTS) call to action. <u>Bioessays</u>. 2020;42(1):e1900189.
- ³⁰⁸ 21. Lang TA, Altman DG. Basic statistical reporting for articles published in biomedical journals: The "statistical analyses and methods in the published literature" or the SAMPL guidelines. <u>Int J Nurs</u>

 Stud. 2015;52(1):5-9.
- Gentleman R, Lang DT. Statistical analyses and reproducible research. <u>Journal of Computational and Graphical Statistics</u>. 2007;16(1):1-23. Accessed August 16, 2022. http://www.jstor.org/stable/27594227
- 23. Indrayan A. Reporting of basic statistical methods in biomedical journals: Improved SAMPL guidelines. <u>Indian Pediatrics</u>. 2020;57(1):43-48. doi:10.1007/s13312-020-1702-4