

# **CS-GY 6763: Lecture 4**

## **High Dimensional Geometry**

---

NYU, Prof. Ainesh Bakshi

# Unifying Theme of the Course

**How do we deal with data (vectors) in high-dimensions?**

- High-dimensional similarity search.

**How do we deal with data (vectors) in high-dimensions?**

- High-dimensional similarity search.
- Iterative methods for optimizing functions in high-dimensions.

## How do we deal with data (vectors) in high-dimensions?


- High-dimensional similarity search.
- Iterative methods for optimizing functions in high-dimensions.
- SVD + low-rank approximation to find and visualize low-dimensional structure.

## How do we deal with data (vectors) in high-dimensions?

- High-dimensional similarity search.
- Iterative methods for optimizing functions in high-dimensions.
- SVD + low-rank approximation to find and visualize low-dimensional structure.
- Convert large graphs to high-dimensional vector data to uncover interesting things.


# High Dimensional is Not Like Low Dimensional

Often visualize data and algorithms in 1,2, or 3 dimensions.



# High Dimensional is Not Like Low Dimensional

Often visualize data and algorithms in 1,2, or 3 dimensions.



**This lecture:** Prove that high-dimensional space looks **very different** from low-dimensional space. These images are rarely very informative!

# Sketching and Dimensionality Reduction

**Next lecture:** Ignore our own advice.




# Sketching and Dimensionality Reduction

**Next lecture:** Ignore our own advice.

Learn about **sketching, aka dimensionality reduction** techniques that seek to approximate high-dimensional vectors with much lower dimensional vectors.

- Johnson-Lindenstrauss lemma for  $\ell_2$  space.
- MinHash for binary vectors (next class) .




# Sketching and Dimensionality Reduction

**Next lecture:** Ignore our own advice.

Learn about **sketching, aka dimensionality reduction** techniques that seek to approximate high-dimensional vectors with much lower dimensional vectors.

- Johnson-Lindenstrauss lemma for  $\ell_2$  space.
- MinHash for binary vectors (next class) .



This lecture should help you understand the potential and limitations of these methods.

# Orthogonal Vectors



Recall the inner product between two  $d$  dimensional vectors:

$$\langle \mathbf{x}, \mathbf{y} \rangle = \mathbf{x}^T \mathbf{y} = \mathbf{y}^T \mathbf{x} = \sum_{j=1}^d x[j]y[j]$$

# Orthogonal Vectors

Recall the inner product between two  $d$  dimensional vectors:



$$\langle \mathbf{x}, \mathbf{y} \rangle = \mathbf{x}^T \mathbf{y} = \mathbf{y}^T \mathbf{x} = \sum_{j=1}^d x[j]y[j]$$



# Orthogonal Vectors

Recall the inner product between two  $d$  dimensional vectors:



$$\langle \mathbf{x}, \mathbf{y} \rangle = \mathbf{x}^T \mathbf{y} = \mathbf{y}^T \mathbf{x} = \sum_{j=1}^d x[j]y[j]$$



$$\langle \mathbf{x}, \mathbf{y} \rangle = \cos(\theta) \cdot \|\mathbf{x}\|_2 \cdot \|\mathbf{y}\|_2$$



# Orthogonal Vectors

What is the largest set of **mutually orthogonal** unit vectors  $\mathbf{x}_1, \dots, \mathbf{x}_t$  in  $d$ -dimensional space? I.e. with inner product  $|\mathbf{x}_i^T \mathbf{x}_j| = 0$  for all  $i, j$ .



# Orthogonal Vectors

What is the largest set **nearly orthogonal** unit vectors  $\mathbf{x}_1, \dots, \mathbf{x}_t$  in 2 or 3 dimensions? I.e., with inner product  $|\mathbf{x}_i^T \mathbf{x}_j| \leq \epsilon$  for all  $i, j$ .  
Consider the case when  $\epsilon$  is a constant. E.g.  $\epsilon = 1/10$ .



# Orthogonal Vectors

What is the largest set **nearly orthogonal** unit vectors  $\mathbf{x}_1, \dots, \mathbf{x}_t$  in  $d$  dimensions? I.e., with inner product  $|\mathbf{x}_i^T \mathbf{x}_j| \leq \epsilon$  for all  $i, j$ .  
Consider the case when  $\epsilon$  is a constant. E.g.  $\epsilon = 1/10$ .

1.  $d$

2.  $\Theta(d)$

3.  $\Theta(d^2)$

4.  $2^{\Theta(d)}$



# Orthogonal Vectors

**Formal Claim:** In  $d$ -dimensional space, there are  $2^{\Theta(\epsilon^2 d)}$  unit vectors with all pairwise inner products  $\leq \epsilon$ , where  $\epsilon \gg 1/\sqrt{d}$ .

# Orthogonal Vectors

**Formal Claim:** In  $d$ -dimensional space, there are  $2^{\Theta(\epsilon^2 d)}$  unit vectors with all pairwise inner products  $\leq \epsilon$ , where  $\epsilon \gg 1/\sqrt{d}$ .

**Proof strategy:** Use the **Probabilistic Method**! For  $t = 2^{\Theta(\epsilon^2 d)}$ , define a random process which generates random vectors  $\mathbf{x}_1, \dots, \mathbf{x}_t$  that are unlikely to have large inner product.

# Orthogonal Vectors

**Formal Claim:** In  $d$ -dimensional space, there are  $2^{\Theta(\epsilon^2 d)}$  unit vectors with all pairwise inner products  $\leq \epsilon$ , where  $\epsilon \gg 1/\sqrt{d}$ .

**Proof strategy:** Use the **Probabilistic Method**! For  $t = 2^{\Theta(\epsilon^2 d)}$ , define a random process which generates random vectors  $\mathbf{x}_1, \dots, \mathbf{x}_t$  that are unlikely to have large inner product.

1. Claim that, with non-zero probability,  $|\mathbf{x}_i^T \mathbf{x}_j| \leq \epsilon$  for all  $i, j$ .
2. Conclude that there must exist some set of  $t$  unit vectors with all pairwise inner-products bounded by  $\epsilon$ .

# Probabilistic Method

**Claim:** There is an exponential number (i.e.,  $2^{\Theta(d)}$ ) of nearly orthogonal unit vectors in  $d$  dimensional space.

**Proof:** Let  $\mathbf{x}_1, \dots, \mathbf{x}_t$  all have independent random entries, each set to  $\pm \frac{1}{\sqrt{d}}$  with equal probability. Ex.

$$\mathbf{x}_1 = \left( \frac{1}{\sqrt{d}}, \frac{1}{\sqrt{d}}, -\frac{1}{\sqrt{d}}, \frac{1}{\sqrt{d}}, \dots, \frac{1}{\sqrt{d}} \right)$$

# Probabilistic Method

**Claim:** There is an exponential number (i.e.,  $2^{\Theta(d)}$ ) of nearly orthogonal unit vectors in  $d$  dimensional space.

**Proof:** Let  $\mathbf{x}_1, \dots, \mathbf{x}_t$  have independent random entries, each  $\pm \frac{1}{\sqrt{d}}$  w.p.  $1/2$ .

- $\|\mathbf{x}_i\|_2^2 =$

# Probabilistic Method

**Claim:** There is an exponential number (i.e.,  $2^{\Theta(d)}$ ) of nearly orthogonal unit vectors in  $d$  dimensional space.

**Proof:** Let  $\mathbf{x}_1, \dots, \mathbf{x}_t$  have independent random entries, each  $\pm \frac{1}{\sqrt{d}}$  w.p.  $1/2$ .

- $\|\mathbf{x}_i\|_2^2 = \sum_{k=1}^d \frac{1}{d} = 1$

# Probabilistic Method

**Claim:** There is an exponential number (i.e.,  $2^{\Theta(d)}$ ) of nearly orthogonal unit vectors in  $d$  dimensional space.

**Proof:** Let  $\mathbf{x}_1, \dots, \mathbf{x}_t$  have independent random entries, each  $\pm \frac{1}{\sqrt{d}}$  w.p.  $1/2$ .

- $\|\mathbf{x}_i\|_2^2 = \sum_{k=1}^d \frac{1}{d} = 1$
- $\mathbb{E}[\mathbf{x}_i^T \mathbf{x}_j] =$

# Probabilistic Method

**Claim:** There is an exponential number (i.e.,  $2^{\Theta(d)}$ ) of nearly orthogonal unit vectors in  $d$  dimensional space.

**Proof:** Let  $\mathbf{x}_1, \dots, \mathbf{x}_t$  have independent random entries, each  $\pm \frac{1}{\sqrt{d}}$  w.p.  $1/2$ .

- $\|\mathbf{x}_i\|_2^2 = \sum_{k=1}^d \frac{1}{d} = 1$
- $\mathbb{E}[\mathbf{x}_i^T \mathbf{x}_j] = \sum_{k=1}^d \mathbb{E}[\mathbf{x}_i[k] \mathbf{x}_j[k]] = \sum_{k=1}^d \mathbb{E}[\mathbf{x}_i[k]] \mathbb{E}[\mathbf{x}_j[k]] = 0$



# Probabilistic Method

**Claim:** There is an exponential number (i.e.,  $2^{\Theta(d)}$ ) of nearly orthogonal unit vectors in  $d$  dimensional space.

**Proof:** Let  $\mathbf{x}_1, \dots, \mathbf{x}_t$  have independent random entries, each  $\pm \frac{1}{\sqrt{d}}$  w.p.  $1/2$ .

- $\|\mathbf{x}_i\|_2^2 = \sum_{k=1}^d \frac{1}{d} = 1$
- $\mathbb{E}[\mathbf{x}_i^T \mathbf{x}_j] = \sum_{k=1}^d \mathbb{E}[\mathbf{x}_i[k] \mathbf{x}_j[k]] = \sum_{k=1}^d \mathbb{E}[\mathbf{x}_i[k]] \mathbb{E}[\mathbf{x}_j[k]] = 0$
- $\mathbf{x}_i[k] \mathbf{x}_j[k] = \begin{cases} +1/d & \text{w.p. } 1/2 \\ -1/d & \text{w.p. } 1/2 \end{cases}$

# Probabilistic Method

**Claim:** There is an exponential number (i.e.,  $2^{\Theta(d)}$ ) of nearly orthogonal unit vectors in  $d$  dimensional space.

**Proof:** Let  $\mathbf{x}_1, \dots, \mathbf{x}_t$  have independent random entries, each  $\pm \frac{1}{\sqrt{d}}$  w.p.  $1/2$ .

- $\|\mathbf{x}_i\|_2^2 = \sum_{k=1}^d \frac{1}{d} = 1$
- $\mathbb{E}[\mathbf{x}_i^T \mathbf{x}_j] = \sum_{k=1}^d \mathbb{E}[\mathbf{x}_i[k] \mathbf{x}_j[k]] = \sum_{k=1}^d \mathbb{E}[\mathbf{x}_i[k]] \mathbb{E}[\mathbf{x}_j[k]] = 0$
- $\mathbf{x}_i[k] \mathbf{x}_j[k] = \begin{cases} +1/d & \text{w.p. } 1/2 \\ -1/d & \text{w.p. } 1/2 \end{cases}$
- $\text{Var}[\mathbf{x}_i^T \mathbf{x}_j] =$

# Probabilistic Method

**Claim:** There is an exponential number (i.e.,  $2^{\Theta(d)}$ ) of nearly orthogonal unit vectors in  $d$  dimensional space.

**Proof:** Let  $\mathbf{x}_1, \dots, \mathbf{x}_t$  have independent random entries, each  $\pm \frac{1}{\sqrt{d}}$  w.p.  $1/2$ .

- $\|\mathbf{x}_i\|_2^2 = \sum_{k=1}^d \frac{1}{d} = 1$
- $\mathbb{E}[\mathbf{x}_i^T \mathbf{x}_j] = \sum_{k=1}^d \mathbb{E}[\mathbf{x}_i[k] \mathbf{x}_j[k]] = \sum_{k=1}^d \mathbb{E}[\mathbf{x}_i[k]] \mathbb{E}[\mathbf{x}_j[k]] = 0$
- $\mathbf{x}_i[k] \mathbf{x}_j[k] = \begin{cases} +1/d & \text{w.p. } 1/2 \\ -1/d & \text{w.p. } 1/2 \end{cases}$
- $\text{Var}[\mathbf{x}_i^T \mathbf{x}_j] = \sum_{k=1}^d \text{Var}[\mathbf{x}_i[k] \mathbf{x}_j[k]] = \sum_{k=1}^d \frac{1}{d^2} = \frac{1}{d}$

## Informal Proof

Let  $Z = \mathbf{x}_i^T \mathbf{x}_j = \sum_{k=1}^d C_k$  where each  $C_k$  is random  $\pm \frac{1}{d}$ .

## Informal Proof

Let  $Z = \mathbf{x}_i^T \mathbf{x}_j = \sum_{k=1}^d C_k$  where each  $C_k$  is random  $\pm \frac{1}{d}$ .

$Z$  is a sum of many i.i.d. random variables, so looks approximately Gaussian. Roughly, we expect that:

$$\Pr[|Z - \mathbb{E}Z| \geq \alpha \cdot \sigma] \leq O(e^{-\alpha^2})$$

## Informal Proof

Let  $Z = \mathbf{x}_i^T \mathbf{x}_j = \sum_{k=1}^d C_k$  where each  $C_k$  is random  $\pm \frac{1}{d}$ .

$Z$  is a sum of many i.i.d. random variables, so looks approximately Gaussian. Roughly, we expect that:

$$\Pr[|Z - \mathbb{E}Z| \geq \alpha \cdot \sigma] \leq O(e^{-\alpha^2})$$

We want  $\alpha \cdot \sigma = \epsilon$ , where  $\sigma = 1/\sqrt{d}$ . Therefore, applying Chernoff with  $\alpha = \epsilon\sqrt{d}$ , we have:

$$\Pr[|Z| \geq \epsilon] \leq O(e^{-\epsilon^2 d}).$$

## Informal Proof

Let  $Z = \mathbf{x}_i^T \mathbf{x}_j = \sum_{k=1}^d C_k$  where each  $C_k$  is random  $\pm \frac{1}{d}$ .

$Z$  is a sum of many i.i.d. random variables, so looks approximately Gaussian. Roughly, we expect that:

$$\Pr[|Z - \mathbb{E}Z| \geq \alpha \cdot \sigma] \leq O(e^{-\alpha^2})$$

We want  $\alpha \cdot \sigma = \epsilon$ , where  $\sigma = 1/\sqrt{d}$ . Therefore, applying Chernoff with  $\alpha = \epsilon\sqrt{d}$ , we have:

$$\Pr[|Z| \geq \epsilon] \leq O(e^{-\epsilon^2 d}).$$

By a union bound, for any pairs  $x_i, x_j$  from  $x_1, x_2, x_3, \dots, x_t$ ,

$$\Pr[|\mathbf{x}_i^T \mathbf{x}_j| \geq \epsilon] \leq \binom{t}{2} \cdot O(e^{-\epsilon^2 d}) \leq 0.1$$

## Informal Proof

Let  $Z = \mathbf{x}_i^T \mathbf{x}_j = \sum_{k=1}^d C_k$  where each  $C_k$  is random  $\pm \frac{1}{d}$ .

$Z$  is a sum of many i.i.d. random variables, so looks approximately Gaussian. Roughly, we expect that:

$$\Pr[|Z - \mathbb{E}Z| \geq \alpha \cdot \sigma] \leq O(e^{-\alpha^2})$$

We want  $\alpha \cdot \sigma = \epsilon$ , where  $\sigma = 1/\sqrt{d}$ . Therefore, applying Chernoff with  $\alpha = \epsilon\sqrt{d}$ , we have:

$$\Pr[|Z| \geq \epsilon] \leq O(e^{-\epsilon^2 d}).$$

By a union bound, for any pairs  $x_i, x_j$  from  $x_1, x_2, x_3, \dots, x_t$ ,

$$\Pr[|\mathbf{x}_i^T \mathbf{x}_j| \geq \epsilon] \leq \binom{t}{2} \cdot O(e^{-\epsilon^2 d}) \leq 0.1$$

Suffices to set  $t = e^{O(\epsilon^2 d)}$ .



Use an exponential concentration inequality!

## Theorem (Chernoff Bound)

*Let  $X_1, X_2, \dots, X_d$  be independent  $\{0, 1\}$ -valued random variables and let  $S = \sum_{i=1}^d X_i$ . We have for any  $\epsilon < 1$  :*

$$\Pr[|S - \mathbb{E}[S]| \geq \epsilon \mathbb{E}[S]] \leq 2e^{\frac{-\epsilon^2 \mathbb{E}[S]}{3}}.$$

Does not immediately apply because we have random variables that are  $\pm 1/d$ , not 0, 1.

Use an exponential concentration inequality!

## Theorem (Chernoff Bound)

*Let  $X_1, X_2, \dots, X_d$  be independent  $\{0, 1\}$ -valued random variables and let  $S = \sum_{i=1}^d X_i$ . We have for any  $\epsilon < 1$  :*

$$\Pr[|S - \mathbb{E}[S]| \geq \epsilon \mathbb{E}[S]] \leq 2e^{\frac{-\epsilon^2 \mathbb{E}[S]}{3}}.$$

Does not immediately apply because we have random variables that are  $\pm 1/d$ , not 0, 1.

Use an exponential concentration inequality!

## Theorem (Chernoff Bound)

*Let  $X_1, X_2, \dots, X_d$  be independent  $\{0, 1\}$ -valued random variables and let  $S = \sum_{i=1}^d X_i$ . We have for any  $\epsilon < 1$  :*

$$\Pr[|S - \mathbb{E}[S]| \geq \epsilon \mathbb{E}[S]] \leq 2e^{\frac{-\epsilon^2 \mathbb{E}[S]}{3}}.$$

Does not immediately apply because we have random variables that are  $\pm 1/d$ , not 0, 1.

**Common trick: shift and scale to transform to the binary case.**

$$\begin{aligned}\mathbf{x}_i^T \mathbf{x}_j = Z &= \sum_{k=1}^d C_k = \frac{2}{d} \sum_{k=1}^d \frac{d}{2} \cdot C_k && \text{Let } \left( \frac{d}{2} C_k = B_k - \frac{1}{2} \right) \\ &= \frac{2}{d} \cdot \left( \sum_{k=1}^d B_k - 1/2 \right) \\ &= \frac{2}{d} \cdot \left( -\frac{d}{2} + \sum_{k=1}^d B_k \right)\end{aligned}$$

$$\begin{aligned}\mathbf{x}_i^T \mathbf{x}_j &= Z = \sum_{k=1}^d C_k = \frac{2}{d} \sum_{k=1}^d \frac{d}{2} \cdot C_k && \text{Let } \left( \frac{d}{2} C_k = B_k - \frac{1}{2} \right) \\ &= \frac{2}{d} \cdot \left( \sum_{k=1}^d B_k - 1/2 \right) \\ &= \frac{2}{d} \cdot \left( -\frac{d}{2} + \sum_{k=1}^d B_k \right)\end{aligned}$$

Observe  $B_k$  is uniform in  $\{0, 1\}$ .

$$\mathbf{x}_i^T \mathbf{x}_j = Z = \frac{2}{d} \cdot \left( -\frac{d}{2} + \sum_{k=1}^d B_k \right), \quad B_k \text{ uniform in } \{0, 1\}.$$

$$\begin{aligned} \Pr[|Z| > \epsilon] &= \Pr \left[ \left| \sum_{k=1}^d B_k - \frac{d}{2} \right| > \frac{\epsilon d}{2} \right] \\ &= \Pr \left[ \left| \sum_{k=1}^d B_k - \mathbb{E} \left[ \sum_{k=1}^d B_k \right] \right| > \epsilon \cdot \mathbb{E} \left[ \sum_{k=1}^d B_k \right] \right] \end{aligned}$$

# Chernoff Bound

## Theorem (Chernoff Bound)

*Let  $X_1, X_2, \dots, X_d$  be independent  $\{0, 1\}$ -valued random variables and let  $S = \sum_{i=1}^d X_i$ . We have for any  $\epsilon < 1$  :*

$$\Pr[|S - \mathbb{E}[S]| \geq \epsilon \mathbb{E}[S]] \leq 2e^{\frac{-\epsilon^2 \mathbb{E}[S]}{3}}.$$

Apply with  $X_1, \dots, X_d = B_1, \dots, B_d$ :

$$\Pr[|S - \mathbb{E}[S]| \geq \epsilon \mathbb{E}[S]] \leq 2e^{\frac{-\epsilon^2 \mathbb{E}[S]}{3}} = 2e^{\frac{-\epsilon^2 d}{6}}$$

## Conclusion from Chernoff bound:

For any  $i, j$  pair,  $\Pr[|\mathbf{x}_i^T \mathbf{x}_j| < \epsilon] \geq 1 - 2e^{-\epsilon^2 d/6}$ .

By a union bound:

For all  $i, j$  pairs simultaneously,  $\Pr[|\mathbf{x}_i^T \mathbf{x}_j| < \epsilon] \geq 1 - \binom{t}{2} \cdot 2e^{-\epsilon^2 d/6}$ .



## Conclusion from Chernoff bound:

For any  $i, j$  pair,  $\Pr[|\mathbf{x}_i^T \mathbf{x}_j| < \epsilon] \geq 1 - 2e^{-\epsilon^2 d/6}$ .

By a union bound:

For all  $i, j$  pairs simultaneously,  $\Pr[|\mathbf{x}_i^T \mathbf{x}_j| < \epsilon] \geq 1 - \binom{t}{2} \cdot 2e^{-\epsilon^2 d/6}$ .

Suffices to set  $t = e^{O(\epsilon^2 d)}$  to get for all pairs  $x_i, x_j$  in the set  $x_1, x_2, \dots, x_t$ ,  $\Pr[|\mathbf{x}_i^T \mathbf{x}_j| < \epsilon] \geq 0.9$ .

## Conclusion from Chernoff bound:

For any  $i, j$  pair,  $\Pr[|\mathbf{x}_i^T \mathbf{x}_j| < \epsilon] \geq 1 - 2e^{-\epsilon^2 d/6}$ .

By a union bound:

For all  $i, j$  pairs simultaneously,  $\Pr[|\mathbf{x}_i^T \mathbf{x}_j| < \epsilon] \geq 1 - \binom{t}{2} \cdot 2e^{-\epsilon^2 d/6}$ .

Suffices to set  $t = e^{O(\epsilon^2 d)}$  to get for all pairs  $x_i, x_j$  in the set  $x_1, x_2, \dots, x_t$ ,  $\Pr[|\mathbf{x}_i^T \mathbf{x}_j| < \epsilon] \geq 0.9$ .

Therefore, there exists some set of  $t = e^{O(\epsilon^2 d)}$  unit vectors with all pairwise inner products  $\leq \epsilon$ .

# Orthogonal Vectors

**Final result:** In  $d$ -dimensional space, there are  $2^{\Theta(\epsilon^2 d)}$  unit vectors with all pairwise inner products  $\leq \epsilon$ .


**Corollary of proof:** Random vectors tend to be far apart (and roughly equidistant) in high-dimensions.

# Orthogonal Vectors

**Final result:** In  $d$ -dimensional space, there are  $2^{\Theta(\epsilon^2 d)}$  unit vectors with all pairwise inner products  $\leq \epsilon$ .


**Corollary of proof:** Random vectors tend to be far apart (and roughly equidistant) in high-dimensions.

$$\|x_i - x_j\|_2^2 = \|x_i\|_2^2 + \|x_j\|_2^2 - 2\langle x_i, x_j \rangle \approx 2$$




# Curse of Dimensionality

**Curse of dimensionality:** Suppose we want to use e.g.  $k$ -nearest neighbors to learn a function or classify points in  $\mathbb{R}^d$ . If our data distribution is truly random, we typically need an exponential amount of data.



# Curse of Dimensionality


**Curse of dimensionality:** Suppose we want to use e.g.  $k$ -nearest neighbors to learn a function or classify points in  $\mathbb{R}^d$ . If our data distribution is truly random, we typically need an exponential amount of data.



**The existence of lower dimensional structure in our data is often the only reason we can hope to learn.**

# Curse of Dimensionality

## Low-dimensional structure.



For example, data lies on low-dimensional subspace, or does so after transformation. Or function can be represented by a restricted class of functions, like neural net with specific architecture.

## Unit Ball in High Dimensions

Let  $\mathcal{B}_d$  be the unit ball in  $d$  dimensions:

$$\mathcal{B}_d = \{\mathbf{x} \in \mathbb{R}^d : \|\mathbf{x}\|_2 \leq 1\}.$$



## Unit Ball in High Dimensions

Let  $\mathcal{B}_d$  be the unit ball in  $d$  dimensions:

$$\mathcal{B}_d = \{\mathbf{x} \in \mathbb{R}^d : \|\mathbf{x}\|_2 \leq 1\}.$$

Volume of radius  $R$  ball is  $\frac{\pi^{d/2}}{(d/2)!} \cdot R^d$ .


# Unit Ball in High Dimensions

Let  $\mathcal{B}_d$  be the unit ball in  $d$  dimensions:

$$\mathcal{B}_d = \{\mathbf{x} \in \mathbb{R}^d : \|\mathbf{x}\|_2 \leq 1\}.$$

Volume of radius  $R$  ball is  $\frac{\pi^{d/2}}{(d/2)!} \cdot R^d$ .

What percentage of volume of  $\mathcal{B}_d$  falls within  $\epsilon$  of its surface?




# Unit Ball in High Dimensions

Let  $\mathcal{B}_d$  be the unit ball in  $d$  dimensions:

$$\mathcal{B}_d = \{\mathbf{x} \in \mathbb{R}^d : \|\mathbf{x}\|_2 \leq 1\}.$$

Volume of radius  $R$  ball is  $\frac{\pi^{d/2}}{(d/2)!} \cdot R^d$ .

What percentage of volume of  $\mathcal{B}_d$  falls within  $\epsilon$  of its surface?



$$\frac{\text{Volume of Diff}}{\text{Total Volume}} = \frac{C \cdot 1^d - C \cdot (1 - \epsilon)^d}{C \cdot 1^d}$$


# Unit Ball in High Dimensions

Let  $\mathcal{B}_d$  be the unit ball in  $d$  dimensions:

$$\mathcal{B}_d = \{\mathbf{x} \in \mathbb{R}^d : \|\mathbf{x}\|_2 \leq 1\}.$$

Volume of radius  $R$  ball is  $\frac{\pi^{d/2}}{(d/2)!} \cdot R^d$ .

What percentage of volume of  $\mathcal{B}_d$  falls within  $\epsilon$  of its surface?



$$\begin{aligned} \frac{\text{Volume of Diff}}{\text{Total Volume}} &= \frac{C \cdot 1^d - C \cdot (1 - \epsilon)^d}{C \cdot 1^d} \\ &= 1 - (1 - \epsilon)^d \end{aligned}$$


# Unit Ball in High Dimensions

Let  $\mathcal{B}_d$  be the unit ball in  $d$  dimensions:

$$\mathcal{B}_d = \{\mathbf{x} \in \mathbb{R}^d : \|\mathbf{x}\|_2 \leq 1\}.$$

Volume of radius  $R$  ball is  $\frac{\pi^{d/2}}{(d/2)!} \cdot R^d$ .

What percentage of volume of  $\mathcal{B}_d$  falls within  $\epsilon$  of its surface?



$$\begin{aligned}\frac{\text{Volume of Diff}}{\text{Total Volume}} &= \frac{C \cdot 1^d - C \cdot (1 - \epsilon)^d}{C \cdot 1^d} \\ &= 1 - (1 - \epsilon)^d \\ &= 1 - \left( (1 - \epsilon)^{1/\epsilon} \right)^{\epsilon d}\end{aligned}$$


# Unit Ball in High Dimensions

Let  $\mathcal{B}_d$  be the unit ball in  $d$  dimensions:

$$\mathcal{B}_d = \{\mathbf{x} \in \mathbb{R}^d : \|\mathbf{x}\|_2 \leq 1\}.$$

Volume of radius  $R$  ball is  $\frac{\pi^{d/2}}{(d/2)!} \cdot R^d$ .

What percentage of volume of  $\mathcal{B}_d$  falls within  $\epsilon$  of its surface?



$$\begin{aligned}\frac{\text{Volume of Diff}}{\text{Total Volume}} &= \frac{C \cdot 1^d - C \cdot (1 - \epsilon)^d}{C \cdot 1^d} \\ &= 1 - (1 - \epsilon)^d \\ &= 1 - \left((1 - \epsilon)^{1/\epsilon}\right)^{\epsilon d} \\ &\approx 1 - (1/e)^{\epsilon d} = 1 - e^{-\epsilon d}.\end{aligned}$$

# Isoperimetric Inequality


All but a vanishing small  $2^{-\Theta(\epsilon d)}$  fraction of a unit ball's volume is within  $\epsilon$  of its surface.

**Isoperimetric Inequality:** the ball has the minimum surface area/volume ratio of any shape.

# Isoperimetric Inequality

All but a vanishing small  $2^{-\Theta(\epsilon d)}$  fraction of a unit ball's volume is within  $\epsilon$  of its surface.

**Isoperimetric Inequality:** the ball has the minimum surface area/volume ratio of any shape.






# Isoperimetric Inequality

All but a vanishing small  $2^{-\Theta(\epsilon^d)}$  fraction of a unit ball's volume is within  $\epsilon$  of its surface.


**Isoperimetric Inequality:** the ball has the minimum surface area/volume ratio of any shape.



- If we randomly sample points from any high-dimensional shape, nearly all will fall near its surface.
- 'All points are outliers.'

# Slices of the Unit Ball

What percentage of the volume of  $\mathcal{B}_d$  falls within  $\epsilon$  of its equator?




$$S = \{\mathbf{x} \in \mathcal{B}_d : |x_1| \leq \epsilon\}$$

# Slices of the Unit Ball

What percentage of the volume of  $\mathcal{B}_d$  falls within  $\epsilon$  of its equator?

**Answer:** all but a  $2^{-\Theta(\epsilon^2 d)}$  fraction.




By symmetry, this is true for any equator:

$$S_{\mathbf{t}} = \{\mathbf{x} \in \mathcal{B}_d : \mathbf{x}^T \mathbf{t} \leq \epsilon\}.$$

# Bizarre Shape of Unit Ball

1.  $(1 - 2^{-\Theta(\epsilon^d)})$  fraction of volume lies  $\epsilon$  close to surface.
2.  $(1 - 2^{-\Theta(\epsilon^2 d)})$  fraction of volume lies  $\epsilon$  close to any equator.



**High-dimensional ball looks nothing like 2D ball!**

## Concentration at Equator

**Claim:** All but a  $2^{-\Theta(\epsilon^2 d)}$  fraction of the volume of the ball falls within  $\epsilon$  of its equator.

**Equivalent:** Fix the equator to be horizontal. If we draw a point  $\mathbf{x}$  randomly from the unit ball,  $|x_1| \leq \epsilon$  with probability  $\geq 1 - 2^{-\Theta(\epsilon^2 d)}$ .

## Concentration at Equator

**Claim:** All but a  $2^{-\Theta(\epsilon^2 d)}$  fraction of the volume of the ball falls within  $\epsilon$  of its equator.

**Equivalent:** Fix the equator to be horizontal. If we draw a point  $\mathbf{x}$  randomly from the unit ball,  $|x_1| \leq \epsilon$  with probability  $\geq 1 - 2^{-\Theta(\epsilon^2 d)}$ .

**Proof Strategy:**

- Instead of drawing random vectors from the unit ball, we will draw a random vector from the unit sphere (the surface of the ball).

## Concentration at Equator

**Claim:** All but a  $2^{-\Theta(\epsilon^2 d)}$  fraction of the volume of the ball falls within  $\epsilon$  of its equator.

**Equivalent:** Fix the equator to be horizontal. If we draw a point  $\mathbf{x}$  randomly from the unit ball,  $|x_1| \leq \epsilon$  with probability  $\geq 1 - 2^{-\Theta(\epsilon^2 d)}$ .

**Proof Strategy:**

- Instead of drawing random vectors from the unit ball, we will draw a random vector from the unit sphere (the surface of the ball).
- Let  $\mathbf{w} = \frac{\mathbf{x}}{\|\mathbf{x}\|_2}$ , where  $\mathbf{x}$  is a random vector from the unit ball.


# Concentration at Equator

**Claim:** All but a  $2^{-\Theta(\epsilon^2 d)}$  fraction of the volume of the ball falls within  $\epsilon$  of its equator.

**Equivalent:** Fix the equator to be horizontal. If we draw a point  $\mathbf{x}$  randomly from the unit ball,  $|x_1| \leq \epsilon$  with probability  $\geq 1 - 2^{-\Theta(\epsilon^2 d)}$ .

**Proof Strategy:**

- Instead of drawing random vectors from the unit ball, we will draw a random vector from the unit sphere (the surface of the ball).
- Let  $\mathbf{w} = \frac{\mathbf{x}}{\|\mathbf{x}\|_2}$ , where  $\mathbf{x}$  is a random vector from the unit ball.
- $\Pr[|x_1| \leq \epsilon] \geq \Pr[|w_1| \leq \epsilon] \quad (\text{why?})$





## Concentration at Equator

**Claim:**  $|w_1| \leq \epsilon$  with probability  $\geq 1 - 2^{-\Theta(\epsilon^2 d)}$ . This then proves our statement from the previous slide.

## Concentration at Equator



**Claim:**  $|w_1| \leq \epsilon$  with probability  $\geq 1 - 2^{-\Theta(\epsilon^2 d)}$ . This then proves our statement from the previous slide.

How can we generate  $w$ , which is a random vector taken from the unit sphere (the surface of the ball)?

# Concentration at Equator

**Claim:**  $|w_1| \leq \epsilon$  with probability  $\geq 1 - 2^{-\Theta(\epsilon^2 d)}$ . This then proves our statement from the previous slide.

How can we generate  $w$ , which is a random vector taken from the unit sphere (the surface of the ball)?



## Important Fact in High Dimensional Geometry

**Rotational Invariance of Gaussian distribution:** Let  $\mathbf{g}$  be a random Gaussian vector, with each entry drawn from  $\mathcal{N}(0, 1)$ . Then  $\mathbf{w} = \mathbf{g}/\|\mathbf{g}\|_2$  is distributed uniformly on the unit sphere.

**Why?**

## Important Fact in High Dimensional Geometry

**Rotational Invariance of Gaussian distribution:** Let  $\mathbf{g}$  be a random Gaussian vector, with each entry drawn from  $\mathcal{N}(0, 1)$ . Then  $\mathbf{w} = \mathbf{g}/\|\mathbf{g}\|_2$  is distributed uniformly on the unit sphere.

**Why?** The PDF of a high-dimensional Gaussian does not depend on the direction of  $\mathbf{g}$ , only on its length.

## Important Fact in High Dimensional Geometry

**Rotational Invariance of Gaussian distribution:** Let  $\mathbf{g}$  be a random Gaussian vector, with each entry drawn from  $\mathcal{N}(0, 1)$ . Then  $\mathbf{w} = \mathbf{g}/\|\mathbf{g}\|_2$  is distributed uniformly on the unit sphere.

**Why?** The PDF of a high-dimensional Gaussian does not depend on the direction of  $\mathbf{g}$ , only on its length.

Consider the probability density function of a high dimensional Gaussian:

$$p(\mathbf{g}) = p(g[1]) \cdot \dots \cdot p(g[d]) = \prod_{i=1}^d c e^{-\mathbf{g}[i]^2/2}$$

## Important Fact in High Dimensional Geometry

**Rotational Invariance of Gaussian distribution:** Let  $\mathbf{g}$  be a random Gaussian vector, with each entry drawn from  $\mathcal{N}(0, 1)$ . Then  $\mathbf{w} = \mathbf{g}/\|\mathbf{g}\|_2$  is distributed uniformly on the unit sphere.

**Why?** The PDF of a high-dimensional Gaussian does not depend on the direction of  $\mathbf{g}$ , only on its length.

Consider the probability density function of a high dimensional Gaussian:

$$\begin{aligned} p(\mathbf{g}) &= p(g[1]) \cdot \dots \cdot p(g[d]) = \prod_{i=1}^d c e^{-g[i]^2/2} \\ &= c^d e^{\sum_{i=1}^d -g[i]^2/2} \end{aligned}$$

## Important Fact in High Dimensional Geometry

**Rotational Invariance of Gaussian distribution:** Let  $\mathbf{g}$  be a random Gaussian vector, with each entry drawn from  $\mathcal{N}(0, 1)$ . Then  $\mathbf{w} = \mathbf{g}/\|\mathbf{g}\|_2$  is distributed uniformly on the unit sphere.

**Why?** The PDF of a high-dimensional Gaussian does not depend on the direction of  $\mathbf{g}$ , only on its length.


Consider the probability density function of a high dimensional Gaussian:

$$\begin{aligned} p(\mathbf{g}) &= p(g[1]) \cdot \dots \cdot p(g[d]) = \prod_{i=1}^d c e^{-\mathbf{g}[i]^2/2} \\ &= c^d e^{\sum_{i=1}^d -g[i]^2/2} \\ &= c^d e^{-\|\mathbf{g}\|_2^2/2} \end{aligned}$$




# Important Fact in High Dimensional Geometry

**Rotational Invariance of Gaussian distribution:** Let  $\mathbf{g}$  be a random Gaussian vector, with each entry drawn from  $\mathcal{N}(0, 1)$ . Then  $\mathbf{w} = \mathbf{g}/\|\mathbf{g}\|_2$  is distributed uniformly on the unit sphere.




## Proof Strategy

Draw  $\mathbf{g} \sim \mathcal{N}(\mathbf{0}, \mathbf{I})$ . Show that first entry of  $\mathbf{w} = \mathbf{g} / \|\mathbf{g}\|_2 \leq \epsilon$  with very high probability.



# Proof Strategy


Draw  $\mathbf{g} \sim \mathcal{N}(\mathbf{0}, \mathbf{I})$ . Show that first entry of  $\mathbf{w} = \mathbf{g} / \|\mathbf{g}\|_2 \leq \epsilon$  with very high probability.



1. Prove that with high probability, the first entry of  $\mathbf{g} / \sqrt{d}$  is small.

# Proof Strategy

Draw  $\mathbf{g} \sim \mathcal{N}(\mathbf{0}, \mathbf{I})$ . Show that first entry of  $\mathbf{w} = \mathbf{g}/\|\mathbf{g}\|_2 \leq \epsilon$  with very high probability.



1. Prove that with high probability, the first entry of  $\mathbf{g}/\sqrt{d}$  is small.
2. Prove that  $\mathbf{g}/\sqrt{d}$  is very very close to  $\mathbf{g}/\|\mathbf{g}\|_2^2$ , so this vector also has small first entry.

## Concentration at Equator

Let  $\mathbf{g}$  be a random Gaussian vector and  $\mathbf{w} = \mathbf{g}/\|\mathbf{g}\|_2$ .

$$\mathbb{E}[\|\mathbf{g}\|_2^2] = \sum_{i=1}^d \mathbb{E}[g_i^2] = \sum_{i=1}^d \text{Var}[g_i] = d.$$

## Concentration at Equator

Let  $\mathbf{g}$  be a random Gaussian vector and  $\mathbf{w} = \mathbf{g}/\|\mathbf{g}\|_2$ .

$$\mathbb{E}[\|\mathbf{g}\|_2^2] = \sum_{i=1}^d \mathbb{E}[g_i^2] = \sum_{i=1}^d \text{Var}[g_i] = d.$$

**Exercise for home:** Prove that

$$\Pr \left[ \|\mathbf{g}\|_2^2 \leq \frac{1}{2} \mathbb{E}[\|\mathbf{g}\|_2^2] \right] \leq 2^{-\Theta(d)}$$

This should intuitively make sense. Can you tell me why?

## Concentration at Equator

For  $1 - 2^{-\Theta(d)}$  fraction of vectors  $\mathbf{g}$ ,  $\|\mathbf{g}\|_2 \geq \sqrt{d/2}$ . Condition on the event that we get a random vector in this set.

## Concentration at Equator

For  $1 - 2^{-\Theta(d)}$  fraction of vectors  $\mathbf{g}$ ,  $\|\mathbf{g}\|_2 \geq \sqrt{d/2}$ . Condition on the event that we get a random vector in this set.

Recall that  $\mathbf{w} = \frac{\mathbf{g}}{\|\mathbf{g}\|_2}$ . Given this event:

$$\Pr[|w_1| \leq \epsilon] = \Pr[|g_1|/\|\mathbf{g}\|_2 \leq \epsilon]$$



## Concentration at Equator

For  $1 - 2^{-\Theta(d)}$  fraction of vectors  $\mathbf{g}$ ,  $\|\mathbf{g}\|_2 \geq \sqrt{d/2}$ . Condition on the event that we get a random vector in this set.

Recall that  $\mathbf{w} = \frac{\mathbf{g}}{\|\mathbf{g}\|_2}$ . Given this event:

$$\begin{aligned}\Pr[|w_1| \leq \epsilon] &= \Pr[|g_1|/\|\mathbf{g}\|_2 \leq \epsilon] \\ &\geq \Pr\left[|g_1|/\sqrt{d/2} \leq \epsilon\right]\end{aligned}$$

## Concentration at Equator

For  $1 - 2^{-\Theta(d)}$  fraction of vectors  $\mathbf{g}$ ,  $\|\mathbf{g}\|_2 \geq \sqrt{d/2}$ . Condition on the event that we get a random vector in this set.

Recall that  $\mathbf{w} = \frac{\mathbf{g}}{\|\mathbf{g}\|_2}$ . Given this event:

$$\begin{aligned}\Pr[|w_1| \leq \epsilon] &= \Pr[|g_1|/\|\mathbf{g}\|_2 \leq \epsilon] \\ &\geq \Pr\left[|g_1|/\sqrt{d/2} \leq \epsilon\right] \\ &= \Pr\left[|g_1| \leq \epsilon \cdot \sqrt{d/2}\right]\end{aligned}$$

## Concentration at Equator

For  $1 - 2^{-\Theta(d)}$  fraction of vectors  $\mathbf{g}$ ,  $\|\mathbf{g}\|_2 \geq \sqrt{d/2}$ . Condition on the event that we get a random vector in this set.

Recall that  $\mathbf{w} = \frac{\mathbf{g}}{\|\mathbf{g}\|_2}$ . Given this event:

$$\begin{aligned}\Pr[|w_1| \leq \epsilon] &= \Pr[|g_1|/\|\mathbf{g}\|_2 \leq \epsilon] \\ &\geq \Pr\left[|g_1|/\sqrt{d/2} \leq \epsilon\right] \\ &= \Pr\left[|g_1| \leq \epsilon \cdot \sqrt{d/2}\right] \\ &\geq 1 - 2^{-\theta\left((\epsilon \cdot \sqrt{d/2})^2\right)} \quad (\text{Integrate the PDF})\end{aligned}$$

## Concentration at Equator

For  $1 - 2^{-\Theta(d)}$  fraction of vectors  $\mathbf{g}$ ,  $\|\mathbf{g}\|_2 \geq \sqrt{d/2}$ . Condition on the event that we get a random vector in this set.

Recall that  $\mathbf{w} = \frac{\mathbf{g}}{\|\mathbf{g}\|_2}$ . Given this event:

$$\begin{aligned}\Pr[|w_1| \leq \epsilon] &= \Pr[|g_1|/\|\mathbf{g}\|_2 \leq \epsilon] \\ &\geq \Pr\left[|g_1|/\sqrt{d/2} \leq \epsilon\right] \\ &= \Pr\left[|g_1| \leq \epsilon \cdot \sqrt{d/2}\right] \\ &\geq 1 - 2^{-\theta\left((\epsilon \cdot \sqrt{d/2})^2\right)} \quad (\text{Integrate the PDF})\end{aligned}$$

## Concentration at Equator

For  $1 - 2^{-\Theta(d)}$  fraction of vectors  $\mathbf{g}$ ,  $\|\mathbf{g}\|_2 \geq \sqrt{d/2}$ . Condition on the event that we get a random vector in this set.

Recall that  $\mathbf{w} = \frac{\mathbf{g}}{\|\mathbf{g}\|_2}$ . Given this event:


$$\begin{aligned}\Pr[|w_1| \leq \epsilon] &= \Pr[|g_1|/\|\mathbf{g}\|_2 \leq \epsilon] \\ &\geq \Pr\left[|g_1|/\sqrt{d/2} \leq \epsilon\right] \\ &= \Pr\left[|g_1| \leq \epsilon \cdot \sqrt{d/2}\right] \\ &\geq 1 - 2^{-\theta\left((\epsilon \cdot \sqrt{d/2})^2\right)} \quad (\text{Integrate the PDF})\end{aligned}$$

By union bound, overall we have:

$$\Pr[|w_1| \leq \epsilon] \geq 1 - 2^{-\Theta(\epsilon^2 d)} - 2^{-\theta(d)}$$

# Bizarre Shape of Unit Ball

1.  $(1 - 2^{-\Theta(\epsilon^d)})$  fraction of volume lies  $\epsilon$  close to surface.
2.  $(1 - 2^{-\Theta(\epsilon^2 d)})$  fraction of volume lies  $\epsilon$  close to any equator.




**High-dimensional ball looks nothing like 2D ball!**

# High Dimensional Cube

Let  $\mathcal{C}_d$  be the  $d$ -dimensional cube:


$$\mathcal{C}_d = \{\mathbf{x} \in \mathbb{R}^d : |\mathbf{x}(i)| \leq 1 \ \forall i\}.$$



# High Dimensional Cube

Let  $\mathcal{C}_d$  be the  $d$ -dimensional cube:

$$\mathcal{C}_d = \{\mathbf{x} \in \mathbb{R}^d : |\mathbf{x}(i)| \leq 1 \ \forall i\}.$$




In two dimensions, the cube is pretty similar to the ball.



# High Dimensional Cube

Let  $\mathcal{C}_d$  be the  $d$ -dimensional cube:

$$\mathcal{C}_d = \{\mathbf{x} \in \mathbb{R}^d : |\mathbf{x}(i)| \leq 1 \ \forall i\}.$$



In two dimensions, the cube is pretty similar to the ball.

But volume of  $\mathcal{C}_d$  is  $2^d$  while volume of unit ball is  $\frac{\sqrt{\pi}^d}{(d/2)!}$ .

**This is a huge gap!** Cube has  $O(d)^{O(d)}$  more volume.

# High Dimensional Cube

Some other ways to see these shapes are very different:

- $\max_{\mathbf{x} \in \mathcal{B}_d} \|\mathbf{x}\|_2^2 =$

# High Dimensional Cube

Some other ways to see these shapes are very different:

- $\max_{\mathbf{x} \in \mathcal{B}_d} \|\mathbf{x}\|_2^2 = 1$
- $\max_{\mathbf{x} \in \mathcal{C}_d} \|\mathbf{x}\|_2^2 =$

# High Dimensional Cube

Some other ways to see these shapes are very different:

- $\max_{\mathbf{x} \in \mathcal{B}_d} \|\mathbf{x}\|_2^2 = 1$

- $\max_{\mathbf{x} \in \mathcal{C}_d} \|\mathbf{x}\|_2^2 = d$

- $\mathbb{E}_{\mathbf{x} \sim \mathcal{B}_d} \|\mathbf{x}\|_2^2 =$

# High Dimensional Cube

Some other ways to see these shapes are very different:

- $\max_{\mathbf{x} \in \mathcal{B}_d} \|\mathbf{x}\|_2^2 = 1$
- $\max_{\mathbf{x} \in \mathcal{C}_d} \|\mathbf{x}\|_2^2 = d$
- $\mathbb{E}_{\mathbf{x} \sim \mathcal{B}_d} \|\mathbf{x}\|_2^2 = \frac{d}{d+2} \approx 1$
- $\mathbb{E}_{\mathbf{x} \sim \mathcal{C}_d} \|\mathbf{x}\|_2^2 =$


# High Dimensional Cube

Some other ways to see these shapes are very different:

- $\max_{\mathbf{x} \in \mathcal{B}_d} \|\mathbf{x}\|_2^2 = 1$
- $\max_{\mathbf{x} \in \mathcal{C}_d} \|\mathbf{x}\|_2^2 = d$
- $\mathbb{E}_{\mathbf{x} \sim \mathcal{B}_d} \|\mathbf{x}\|_2^2 = \frac{d}{d+2} \approx 1$
- $\mathbb{E}_{\mathbf{x} \sim \mathcal{C}_d} \|\mathbf{x}\|_2^2 = \sum_{i=1}^d \mathbb{E}[x_i^2] = d \cdot \frac{1}{3}$


# High Dimensional Cube

Almost all of the volume of the unit cube falls in its corners, and these corners lie far outside the unit ball.



# On Breaking Intuition



Consider the following setup:







# On Breaking Intuition

Consider the following setup:



# On Breaking Intuition



Consider the following setup:



Place  $2^d$  unit balls in box with side length 2. Look at sphere they enclose. It has radius  $\sqrt{d} - 1$ .

# On Breaking Intuition

Consider the following setup:



Place  $2^d$  unit balls in box with side length 2. Look at sphere they enclose. It has radius  $\sqrt{d} - 1$ .

So for  $d \geq 10$ , the central ball sticks out of the box...