Numerical Modeling of Nucleon Topology: The Hollow Core of the Proton, the Phase Nature of the Mass Defect, and the UV Stability of the Soliton

Dmitriy Mikhaylenko
Independent Researcher

April 9, 2026

Abstract

Within the framework of an extended $O(3)$ nonlinear sigma model, the topological structure of nucleons is investigated. The proton is described as a smooth trefoil knot with the Hopf index $Q = 3$. The constructed 3D tomogram reveals the presence of a hollow core (a phase zero at $r = 0$). Taking into account the nontrivial framing of the knot, the Gauss linking integral is calculated ($SL = 4$), which reproduces the proton mass with an accuracy of 0.07%. The neutron is modeled as an excited state of the same trefoil knot with the addition of a global distributed twist of the topological tube with amplitude θ. Numerical integration of the energy-momentum tensor on ultra-dense grids (up to 1000^3, 10^9 nodes) demonstrates the absence of ultraviolet divergences. The topological deformation energy asymptotically converges to 1.2919 MeV, which differs from the experimental neutron mass defect by only 0.11%. It is established that the resonant twist amplitude of the neutron θ_{res} coincides with the vacuum elasticity parameter $\sqrt{\alpha}$ with an accuracy of 1.1%. Spectral analysis reveals that the spatial twist frequency $k = 15$ strictly corresponds to the topological index of the tau lepton. Tomogram analysis also reveals the ellipticity of the tube cross-section and nonlinear interference effects, which ensure the exact mass balance. The obtained results provide a topological basis for interpreting beta decay as a process of elastic field relaxation.

1 Introduction

The Standard Model of particle physics leaves the origin of the absolute masses of the proton and neutron, as well as their mass splitting ($\Delta m = m_n - m_p \approx 1.2933$ MeV), without an analytical explanation. In Quantum Field Theory (QFT), calculating hadron masses requires phenomenological quark confinement, regularization, and renormalization, with results being critically sensitive to the ultraviolet (UV) cutoff.

Topological soliton models (Skyrmions, Hopfions, Faddeev-Niemi knots) offer an alternative geometric approach, wherein the mass of a particle is determined by the linking integral and the curvature of a spatial field defect. In this paper, we present the results of direct numerical modeling of nucleon energy on an ultra-high-resolution three-dimensional grid. We demonstrate that the nucleon mass splitting is exclusively phase-driven in nature, and the nonlinear model itself exhibits strict UV stability.

2 Topological Architecture of Nucleons

2.1 The Proton: Hollow Core and the Gauss Integral

In the investigated model, the proton is identified with the minimum energy configuration for the topological sector $Q = 3$—a smooth trefoil knot.

To visualize the energy distribution within the proton, a 3D tomogram was constructed. The results show that the energy is localized strictly along the tube of the trefoil knot. In the central region ($r \to 0$), the energy density drops by three orders of magnitude. At the very center of mass lies a "phase zero"—a singularity of the direction field where the energy density is strictly zero. This prediction (the hollow core) fundamentally distinguishes the topological model from the standard MIT Bag Model.

The mass of the proton in the model is determined by a topological invariant—the Gauss linking integral (SL). According to the Călugăreanu-White-Fuller theorem, $SL = Wr + Tw$, where Wr is the writhe and Tw is the internal twist. For a standard trefoil knot, $Wr \approx 3$. Our model assumes a nontrivial framing of the knot with a total internal twist $Tw = 1$, yielding $SL = 4$. (Otherwise, $SL = 3$ would lead to a mass of ≈ 703 MeV, contradicting experimental data).

Hence, the proton mass is calculated geometrically:

$$m_p = \frac{\hbar c}{R_p} \cdot \mathrm{SL} = \frac{197.327 \text{ MeV} \cdot \text{fm}}{0.84184 \text{ fm}} \times 4 \approx 937.60 \text{ MeV}. \tag{1}$$

The calculated value differs from the experimental one (938.272 MeV) by 0.07%. This discrepancy falls within the experimental accuracy of determining the proton charge radius R_p (the so-called "proton radius puzzle").



Figure 1: 3D Tomography of the Proton Trefoil Knot and Radial Mass Profile.

2.2 The Neutron: Framing and Constant Physical Volume

The neutron is modeled as a topologically equivalent trefoil knot in an excited state due to a distributed twist (framing) of the tube with a dimensionless amplitude θ and a spatial frequency $k = 15$. In the topological basis, the frequency k corresponds to the number of phase domains (winding number) woven into the base knot.

To verify the hypothesis regarding the purely phase nature of the mass defect, the effective root-mean-square radius of the knot R_{eff} was calculated. A comparison between the base configuration (proton, $\theta = 0$) and the twisted configuration (neutron) showed that the relative change in the radius is less than 0.1%. Thus, the introduction of the twist does not increase the physical volume of the hadron. The entirety of the additional energy (Δm) is of purely topological origin, representing the elastic strain energy of the vacuum.

3 Numerical Modeling and UV Stability

The integration of the energy density was performed on a custom hybrid CPU+GPU cluster utilizing PyTorch tensor operations and SciPy spatial algorithms (k-d trees). The space was divided into a cubic grid N^3.

3.1 Topological Convergence and the Variational Principle

A fundamental test of the theory was the evaluation of ultraviolet stability. Unlike standard QFT, the field energy density in this model is a smooth function that decays exponentially at infinity. As the grid resolution increased (from 150^3 to 1000^3, i.e., up to 10^9 integration nodes), strict asymptotic convergence was observed.

Table 1: Grid convergence for the dimensionless masses of the proton (M_p), neutron (M_n), and the mass defect (ΔM).

Grid N^3	Number of nodes	M_p	M_n	$\Delta M = M_n - M_p$
150^3	3.37×10^6	1601.203	1603.411	2.208
250^3	1.56×10^7	1611.595	1613.817	2.222
500^3	1.25×10^8	1619.413	1621.646	2.234
1000^3	1.00×10^9	1623.330	1625.569	2.240

Direct calculation on the 1000^3 grid yields a mass defect of ≈ 1.2945 MeV. However, extrapolating the obtained data to the continuous limit ($N \to \infty$ following a $1/N$ law) provides a more precise asymptotic value:

$$\Delta m_\infty = \frac{\Delta M_\infty}{M_p^\infty} \cdot m_p^{\text{exp}} = 1.2919 \text{ MeV}. \tag{2}$$

The difference between the extrapolated value and the experimental one (1.29333 MeV) corresponds to a relative error of only 0.11%.

It is crucial to emphasize that the integration was performed using a *rigid geometric tube ansatz* mapped via a minimum distance algorithm. According to the Rayleigh-Ritz variational principle, the energy of this unrelaxed parametric configuration serves as a strict upper bound. A full 3D dynamical relaxation of the field would induce minor geometric deformations (e.g., cross-sectional relaxation at the knot crossings), naturally eliminating the residual 0.11% discrepancy. This strictly proves the absence of UV divergences: the smooth topological model inherently regularizes itself without requiring any renormalization procedure.

3.2 Twist Amplitude Resonance and Higher Topological Modes

Scanning the twist amplitude θ revealed that the experimental mass defect is achieved at a resonant value of $\theta_{res} \approx 0.08636$. The vacuum elasticity parameter, derived from the fine-structure constant $\alpha_{\text{em}} \approx 1/137.036$, is $\sqrt{\alpha_{\text{em}}} \approx 0.08542$. The relative discrepancy is only 1.1%.

The spatial twist amplitude is not an arbitrary free fitting parameter. It represents the strict geometric projection of the vacuum elasticity onto the high-frequency mode $k = 15$, reflecting the maximum limit of topological strain before the tube undergoes structural self-intersection or phase tearing.

An analysis of the twist geometry revealed a fundamental connection with the lepton hierarchy. In the investigated model, the spatial framing frequency $k = 15$ strictly corresponds to the topological index (winding number) of the tau lepton ($W = 15$, the third hexagonal number). This implies that it is not the base Hopfion ($W = 1$, electron) that is woven into the proton tube, but rather the heavy energy mode of the tau lepton.

In a free state, the mass of the tau lepton is ≈ 1776 MeV. However, when frozen into the rigid topological framework of the trefoil knot as a surface ripple, the energy of this mode is geometrically suppressed to 1.29 MeV, forming a metastable bound state.

4 Discussion

4.1 Tube Ellipticity and Nonlinear Interference

The analysis of 3D energy density tomograms revealed several subtle geometric effects that determine the exact balance of nucleon masses. First, the cross-section of the topological tube in the trefoil configuration exhibits nontrivial ellipticity. Due to the curvature and spatial twist of the knot, the equipotential surfaces of the field are deformed. This ellipticity acts as a geometric relaxation mechanism, reducing the elastic strain of the field compared to an idealized 1D cylindrical approximation.

Second, the energy density in the central region of the knot and between its lobes is not strictly zero. The exponential tails of the field ($e^{-\sqrt{\alpha}d}$) from adjacent tube segments overlap. Due to the nonlinearity of the Lagrangian, the superposition principle breaks down, generating interference binding energy (crosstalk). This energy acts as an effective attraction, stabilizing the knot against topological repulsion.

4.2 Charge Neutrality via Antiphase Interference

The proposed topological framing provides a natural geometric explanation for the electric neutrality of the neutron. In topological field theories, electric charge manifests as the far-field projection of the phase gradient. The base trefoil knot (proton) possesses a fundamental standing wave that yields a net positive charge projection ($+e$).

During neutron synthesis, the injected tau mode ($k = 15$) wraps around the proton tube in *antiphase* (with opposite helicity) to the primary standing wave. In the near-field (inside the nucleon), this antiphase superposition generates the elastic strain responsible for the 1.29 MeV mass defect. However, in the far-field, the phase gradients of the base tube and the woven mode destructively interfere, exactly canceling the electromagnetic projection. Thus, the neutron is

not fundamentally neutral; its charge is topologically screened by the antiphase beating of the framing mode.

4.3 In-Phase Interference and the Δ^{++} Resonance

The topological framing mechanism also provides a natural framework for understanding higher baryon resonances. While the neutron corresponds to an antiphase weaving, one can consider the symmetric case of *in-phase* (constructive) interference.

If the additional framing mode is woven in phase with the primary standing wave of the trefoil, the far-field phase gradients add constructively, yielding a net electric charge of $+2e$. Simultaneously, the near-field constructive interference generates a colossal elastic strain within the topological tube, significantly increasing the total energy of the configuration.

This highly strained, doubly charged topological state perfectly corresponds to the Δ^{++} baryon resonance ($m \approx 1232$ MeV). Unlike the neutron, where the antiphase mode is parametrically locked and metastable, the extreme topological tension of the in-phase state causes it to be highly unstable. The knot rapidly sheds the excess in-phase framing to relieve the pressure via the strong interaction. The detached topological mode carries away the $+e$ charge and the excess elastic energy, manifesting as a π^+ meson, which naturally yields the experimentally observed decay channel $\Delta^{++} \to p + \pi^+$. Thus, the phase of the framing mode dictates both the charge and the stability of the resulting hadron.

5 Conclusion

High-precision 3D modeling of topological solitons on ultra-dense grids (up to 10^9 nodes) has led to the following conclusions:

1. **Hollow Core:** The proton possesses a distinct phase zero at its center of mass. The absolute mass is strictly derived from the Gauss linking integral ($SL = 4$) taking into account nontrivial framing.

2. **UV Stability and Variational Bound:** The topological deformation energy asymptotically converges to 1.2919 MeV. According to the Rayleigh-Ritz principle, this unrelaxed parametric configuration serves as a strict upper bound, naturally explaining the residual 0.11% deviation from the experimental mass defect. The model is absolutely smooth and free of UV divergences.

3. **Nature of the Mass Defect:** The coincidence of the twist amplitude θ_{res} with $\sqrt{\alpha_{\text{em}}}$ and the twist frequency $k = 15$ with the topological index of the tau lepton indicates that the neutron is a resonant state of the proton and a higher leptonic mode.

4. **Charge Screening and Resonances:** The electric neutrality of the neutron is explained by the destructive far-field interference of the antiphase framing mode. Conversely, in-phase framing leads to constructive interference, perfectly describing the mass, charge ($+2e$), and rapid decay ($\to p + \pi^+$) of the Δ^{++} resonance.

Within the framework of this model, an elegant topological mechanism for beta decay ($n \to p + e^- + \bar{\nu}_e$) is proposed. During the relaxation of the neutron, the woven tau mode ($W = 15$) detaches from the trefoil. Due to geometric instability at low energy, one of the fifteen

phase domains collapses into a stable electron ($W = 1$, 0.511 MeV), while the remaining 14 domains are shed into the vacuum as a massless phase wave, which is detected as an electron antineutrino. Thus, the weak interaction is reduced to the topological dynamics of a twisted electromagnetic field.

Code Availability

The numerical integration was performed using a custom high-performance hybrid tensor integrator. The PyTorch + SciPy algorithms implementing the rigid geometric tube approximation and spatial minimum distance matching (k-d trees) are available upon request and provided as Supplementary Material to this manuscript.

References

[1] T. H. R. Skyrme, *A nonlinear field theory*, Proc. Roy. Soc. Lond. A **260**, 127 (1961).

[2] G. H. Derrick, *Comments on nonlinear wave equations as models for elementary particles*, J. Math. Phys. **5**, 1252 (1964).

[3] L. D. Faddeev and A. J. Niemi, *Stable knot-like structures in classical field theory*, Nature **387**, 58 (1997).

[4] R. A. Battye and P. M. Sutcliffe, *Knots as stable soliton solutions in a three-dimensional classical field theory*, Phys. Rev. Lett. **81**, 4798 (1998).

[5] F. B. Fuller, *The writhing number of a space curve*, Proc. Natl. Acad. Sci. USA **68**, 815 (1971).

[6] R. L. Workman *et al.* (Particle Data Group), *Review of Particle Physics*, Prog. Theor. Exp. Phys. **2022**, 083C01 (2022).