

### Sino Bright School Chongqing

### examtitle

Subject: subject
Date: examdate
Time: duration
Teacher: teacher

#### IMPORTANT: Read together before beginning the exam.

Students must complete the assessments to the best of their ability. Any students found cheating during an exam will be given a zero and referred to the Principal for further disciplinary measures. Cheating is a serious offense.

"Cheating" may include any of the following:

- Having course papers, notes, unapproved data sheets or other course-related materials in the exam room without the express permission of the subject class teacher in charge of the exam.
- Using or even having a cellphone or other electronic device in the exam room without the express permission of the subject class teacher in charge of the exam.
- Attempting to communicate, either verbally or non-verbally with other students during the exam.
- Having exam-related materials prior to the writing of the exam.

Mark Breakdown:	
Part 1: Multiple Choice Questions MCQ $\times$ MC-QNo	Student Name:
Part 2: Short Answer Questions SAQ $\times$ SAQNo	
Part 3: Long Answer Questions LAQ $\times$ LAQNo	
Total Score: 100	Score:

#### Special Instructions/Materials Allowed:

- Pen
- Calculator



# Part I: Multiple Choice Questions ( $5pts \times 10$ )

- 1. If a root of f(x) = 0 lies in [3, 4], what is the midpoint after the first iteration of interval bisection?
  - A) 3.25
  - B) 3.5
  - C) 3.75
  - D) 3.0
- 2. The formula for the first approximation  $x_1$  using linear interpolation is:
  - A)  $x_1 = a \frac{f(a)(b-a)}{f(b)-f(a)}$ B)  $x_1 = \frac{a+b}{2}$

  - C)  $x_1 = a \frac{f(a)}{f'(a)}$ D)  $x_1 = \frac{f(b) f(a)}{b a}$
- 3. The directrix of the parabola  $y^2 = 24x$  is:
  - A) x = -6
  - B) x = 6
  - C) y = -6
  - D) y = 6
- 4. The modulus of z = 3 + 4i is:
  - A) 5
  - B) 7
  - C) 12
  - D) 25
- 5. The iteration formula is  $x_{n+1} = x_n \frac{f(x_n)}{f'(x_n)}$ . What condition is critical for convergence?
  - A)  $f'(x_n) \neq 0$
  - B)  $f(x_n) > 0$
  - C)  $f(x_n) < 0$
  - D)  $f'(x_n) = 0$
- 6. If z = 2 5i, then  $z + z^*$  equals:
  - A) 4
  - B) -10i
  - C) 4 10i
  - D) 0
- 7. A root of a continuous function f(x) exists in [a, b] if:
  - A) f(a) and f(b) are positive
  - B)  $f(a) \cdot f(b) < 0$
  - C) f(a) = f(b)
  - D)  $f(a) \cdot f(b) > 0$
- 8. The Cartesian equation for x = 2t,  $y = 4t^2$  is:
  - A)  $y = x^2$
  - B)  $y = 2x^2$

  - C)  $y = \frac{x^2}{4}$ D)  $y = 4x^2$



9.	The	asymi	ptotes	of	xu =	= 9	are:
∙.	<b>1</b> 110			$O_{\mathbf{I}}$	<i>w u</i> -		$\alpha_{\mathbf{I}} \cdot \cdot$

A) 
$$x = 0$$
 and  $y = 0$ 

B) 
$$x = 3$$
 and  $y = 3$ 

C) 
$$x = 9 \text{ and } y = 9$$

D) None

10. If a quadratic equation has roots 
$$\alpha$$
 and  $\beta$ , then  $\alpha + \beta =$ 

A) 
$$\frac{b}{a}$$
B)  $-\frac{b}{a}$ 
C)  $\frac{c}{a}$ 
D)  $-\frac{c}{a}$ 

C) 
$$\frac{c}{a}$$

D) 
$$-\frac{a}{a}$$

# Part II: Short Answer Questions $(5pts \times 4)$

1. After 3 iterations of interval bisection on [3, 4], the interval length is \_\_\_\_\_.

2. For f(2) = -3 and f(4) = 5, the next approximation using linear interpolation is

3. The focus of the parabola  $y^2 = 28x$  is at \_\_\_\_\_.

4. The argument of z = -1 - i (in radians) is \_\_\_\_\_.

# Part III: Long Answer Questions $(15pts \times 2)$

1. Let  $z^6 = 1$ .

(a) Find all the solutions to the equation.

(b) Show each solution on an Argand diagram.



(c)	Show	that	${\rm each}$	solution	${\rm lies}$	on a	circle	with	center	(0,0)	and	${\rm radius}$	1.
-----	------	------	--------------	----------	--------------	------	--------	------	--------	-------	-----	----------------	----

- 2. Given the quadratic equation  $x^2 5x + 6 = 0$  with roots  $\alpha$  and  $\beta$ :
  - (a) Use Vieta's formulas to find  $\alpha + \beta$  and  $\alpha\beta$ .

(b) Construct a new quadratic equation with roots  $2\alpha$  and  $2\beta$ .

(c) Use the Newton-Raphson method with  $x_0 = 3$  to approximate a root of the new equation. Perform 2 iterations and round your answers to 4 decimal places.