

Probability Theory Homework №2

Ivan Zhytkevych

September 16, 2021

Exercise 3.15

Prove that for arbitrary events A and B:

$$P(AB) = 1 - P(\bar{A}) - P(\bar{B}) + P(\bar{A}\bar{B})$$

Solution:

$$\begin{aligned} P(AB) &= P(A) + P(B) - P(A \cup B) = \\ &= 1 - P(\bar{A}) + (1 - P(\bar{B})) - (1 - P(\overline{A \cup B})) = \\ &= 1 - P(\bar{A}) - P(\bar{B}) + P(\bar{A}\bar{B}) \end{aligned}$$

Exercise 3.16

Four dice are tossed. Find the probability that they will lose the same number of points.

Solution:

Set of all dice configurations:

$$\Omega = \{ (a_1, a_2, a_3, a_4) : a_i = \overline{1, 6} \}$$

$$|\Omega| = C_{6+4-1}^4$$

The configurations we're interested in:

$$S = \{ (a_1, a_2, a_3, a_4) : a_1 = a_2 = a_3 = a_4, a_i = \overline{1, 6} \}$$

$$|S| = 6$$

All configurations are equiprobable:

$$P(S) = \frac{|S|}{|\Omega|} = \frac{6}{126} = 0.047$$

Exercise 3.17

Class consists of r students. Find the probability that at least two students were born in the same month (assume all months are equiprobable for birth).

Solution:

Make a set of all birth configurations:

$$\Omega = \{(a_1, a_2, \dots, a_{12}) : a_i - \text{students born in } i\text{-th month}, \sum a_i = r\}$$

To find probability that at least two students were born in the same month let's find probability of opposite event: number of configurations where none of the students were born in the same month. If r gets bigger than the number of month our probability inevitably turns into 0 because there's no such configurations where each month has less than 2 birthday boy or girl. So we found the probability we need for one case – $r > 12$:

$$r > 12 : P = 1$$

To find the probability for $r < 12$ we follow this path of thinking:

The probability that at least two students have the same birthday month is opposite to probability that none of them has same birthday month. This probability is easier to find:

$$A = \{(a_1, a_2, \dots, a_{12}) : a_i < 2, \sum a_i = r\}$$

$$|A| = A_{12}^r$$

$$\bar{P} = \frac{|A|}{C_{12+r}^r}$$

12^r is number of all configurations (each student has 12 months to pick out of).

$$P = 1 - \frac{A_{12}^r}{C_{12+r}^r}$$

Exercise 3.18

How many people should be in the room for the probability that at least two of those present were born on the same day was more than half?

Solution:

n – number of people in the room.

Using the same approach as in the previous problem: n should be less than number of days in the year in order to calculate the probability we want. Let the number of days in the year be 365, leap year is not taken into consideration.

$$\Omega = \{(a_1, a_2, \dots, a_{365}) : a_i = \overline{1, 365}, \sum a_i = n\}$$

The probability is calculated the same way:

$$P = 1 - \frac{A_{365}^n}{C_{365+n}^n} = 1 - \frac{365! \cdot n! \cdot 364!}{(365 - n)! \cdot (364 + n)!}$$

Exercise 3.19

Numbers $1, 2, \dots, n$ are in random order. Find probability that numbers: a) 1 and 2; b) 1, 2 and 3 are placed next to in the predefined order.

Solution:

$$N = \{1, 2, \dots, n\}$$

Set of all such number configurations:

$$A = \{(a_1, a_2, \dots, a_n) : a_i \in N \setminus \{a_{i-1}, \dots, a_1\}\}$$

$$|A| = n!$$

1. **1 and 2.** D is our event that number 2 is next to 1. The cardinality of this set can be calculated in the following way: number of positions to pick for 1 is $n - 1$ since number 2 is placed next to. Time to arrange other $n - 2$ number in $(n - 2)!$ ways.

$$D = \{t = (a_1, a_2, \dots, a_n) : t \in A, a_i = 1, a_{i+1} = 2\}$$

$$|D| = (n - 1) \cdot (n - 2)!$$

The probability is:

$$P(D) = \frac{(n - 1)(n - 2)!}{n!} = \frac{(n - 1)!}{n!}$$

2. **1, 2 and 3.** Use the same approach. T is our event that 1, 2 and 3 are placed next to each other. The number of such positions is $n - 2$.

$$T = \{t = (a_1, a_2, \dots, a_n) : t \in A, a_i = 1, a_{i+1} = 2, a_{i+2} = 3\}$$

Exercise 3.20

Exam consists of 10 questions. Each question should be answered "yes" or "no". Find a probability that student will give correct answers to at least 70% of all questions picking answer by chance. Solve this problem for a test of 30 and 50 questions.

Solution:

Exercise 3.21

A list of $4N$ tournament participants is divided into four parts. Find a probability that four strongest participants will be in distinct groups.

Solution:

Exercise 3.22

Playing poker, a player has five cards from a deck of 52 cards. Find the probability of the following combinations:

- a **royal flush** ten, jack, queen, king and ace of the same suit
- b **straight flush** five cards in a row of the same suit but not royal flush
- c **four of a kind** four cards of the same order
- d **full house** two and three cards of the same order respectively
- e **flush** five cards of the same suit but neither royal flush nor straight flush
- f **straight** five card in a row, not all are of the same suit

Solution:

Exercise 3.23

Computer center has 3 processors that received n tasks. For each task the processor is randomly picked. Find a probability that exactly one processor will be idle.

Solution: