## 1 Integrals Yielding the Natural Logarithmic Function

Date: Term 1, Fall '23

Notes: A.L. Maagma

**DEFINITION.** Since the natural logarithm is undefined for negative numbers, you will often encounter expressions of the form  $\ln |u|$ . Hence, for derivatives involving absolute values, if u is a differentiable function of x such that  $u \neq 0$ , then

$$\frac{d}{dx}[\ln|u|] = \frac{u'}{u}$$

**EXAMPLE 1.0.0.** Evaluate the equation  $\int \frac{1}{u} du$ .

$$= \ln |u| + C$$

**DEFINITION.** For any rational number  $n \neq -1$ ,

$$\int u^n \, du = \frac{u^{n+1}}{n+1} + C$$

However, for any rational number n = -1,

$$\int u^n \, du = \ln|u| + C$$

**EXAMPLE 1.0.1.** Evaluate the equation  $\int \frac{2}{x} dx$ .

**EXAMPLE 1.0.2.** Evaluate the equation  $\int \frac{dx}{4x-1}$ .

**EXAMPLE 1.0.3.** Find the area of the region on the x-axis and the line x = 3, bounded by the graph of

$$y = \frac{x}{x^2 + 1}$$

**EXAMPLE 1.0.4.** Evaluate the equation  $\int \frac{x+1}{x^2+2x} dx$ .

**EXAMPLE 1.0.5.** Evaluate the equation  $\int \frac{\sec^2 x}{\tan x} dx$ .

**EXAMPLE 1.0.6.** Evaluate the equation  $\int \frac{\ln x}{x} dx$ .

**NOTE.** The integrals to which this formula or rules can be applied may appear in disguised form. For instance, when a rational function has a numerator of degree greater than or equal to that of the denominator, division may reveal a form to which you can apply the rule.

**EXAMPLE 1.0.7.** Evaluate the equation  $\int \frac{x^2+x+1}{x^2+1} dx$ .

**EXAMPLE 1.0.8.** Evaluate the equation  $\int_0^2 \frac{x^2+2}{x+1} dx$ .

## 2 Integrals Involving Logarithmic Functions

**EXAMPLE 2.0.1.** Evaluate the equation  $\int \frac{\log_{10} x}{x}$ .

**EXAMPLE 2.0.2.** Evaluate the equation  $\int \frac{(\log_3 x)^2}{x}$ .

## 3 Integral of Trigonometric Functions

**EXAMPLE 3.0.1.** Evaluate the equation  $\int \tan x \, dx$ .

$$= -\int \frac{-\sin x}{\cos x} dx$$

$$= -\int \frac{du}{u}$$

$$= -\ln |u| + C$$

$$= -\ln |\cos x| + C$$

$$= \ln |(\cos x)^{-1}| + C$$

$$= \ln |\sec x| + C$$

**EXAMPLE 3.0.2.** Evaluate the equation  $\int \cot x \, dx$ .

$$= \int \frac{\cos x}{\sin x} dx \qquad u = \sin x$$

$$= \int \frac{du}{u} \qquad du = \cos x dx$$

$$= \ln|u| + C$$

$$= \ln|\sin x| + C$$