

CHEMOTHERAPY ROTA

W. F. Lamborn
Five South
2nd Avenue

Reference ...JCO 1994; 12: 671-8.....

FLAG

| | YES / NO |
|---------------|----------|
| Trial Patient | |
| Course No: | |
| Start Date: | |

The cytarabine infusion MUST be 4 hours after the fludarabine.

Granisetron 2mg daily orally days 2-6
Dexamethasone 2mg tds orally (last dose not later than 5pm)
days 2-6

Prednisolone 0.5% eye drops 2-hourly in both eyes days 2-11
Filgrastim 300mcg sc od, **days 1-7**

QUEEN ELIZABETH CENTRE FOR THE TREATMENT OF CANCER

CHEMOTHERAPY ROTA

| | | | | | |
|----------------|----------|--------------|----------------|---------------------------|--------------------|
| Document Code: | H-ROTA49 | Written by: | N Duncan | Authorised By: | Consultant |
| Issue No: | 1.0 | Issue date: | September 2004 | Clinical Nurse Specialist | <i>[Signature]</i> |
| Page: | 2 of 2 | Valid until: | Next review | Pharmacist | <i>[Signature]</i> |

Patient label:

FLAG

| DAY NO. DATE | DRUG or ELECTROLYTE | CALCULATION | DOSE | I.V. FLUID | VOLUME (ml) | FLOW RATE | SPECIAL DIRECTIONS | DRUG ADMINISTRATION sig. | DRUG ADMINISTRATION sig. | TIME |
|-----------------|------------------------|---------------------|------|------------|----------------|--------------|----------------------------------|--------------------------------|--------------------------------|------|
| 3 | FLUDARABINE | 30mg/m ² | | N/SALINE | 1000 | 4 hours | | | | |
| | FLUDARABINE | | | N/SALINE | 100 | 30 mins | | | | |
| | | | | N/SALINE | 1000 | 3.5 hrs | | | | |
| | CYTARABINE | 2g/m ² | | N/SALINE | 1000 | 4 hours | 4 hrs after start of fludarabine | | | |
| 4 | FLUDARABINE | 30mg/m ² | | N/SALINE | 1000 | 4 hours | | | | |
| | FLUDARABINE | | | N/SALINE | 100 | 30 mins | | | | |
| | | | | N/SALINE | 1000 | 3.5 hrs | | | | |
| | CYTARABINE | 2g/m ² | | N/SALINE | 1000 | 4 hours | 4 hrs after start of fludarabine | | | |
| 5 | FLUDARABINE | 30mg/m ² | | N/SALINE | 100 | 30 mins | | | | |
| | FLUDARABINE | | | N/SALINE | 1000 | 3.5 hrs | | | | |
| | CYTARABINE | 2g/m ² | | N/SALINE | 1000 | 4 hours | 4 hrs after start of fludarabine | | | |
| 6 | FLUDARABINE | 30mg/m ² | | N/SALINE | 100 | 30 mins | | | | |
| | | | | N/SALINE | 1000 | 3.5 hrs | | | | |
| | CYTARABINE | 2g/m ² | | N/SALINE | 1000 | 4 hours | 4 hrs after start of fludarabine | | | |
| | FLUDARABINE | | | N/SALINE | 100 | 30 mins | | | | |
| | | | | N/SALINE | 1000 | 3.5 hrs | | | | |
| | CYTARABINE | 2g/m ² | | N/SALINE | 1000 | 4 hours | 4 hrs after start of fludarabine | | | |

Prescriber Sig. Date: Chemo Nurse Date: Pharmacist Date: