

D VTD - twice weekly bortezomib (SC daratumumab, bortezomib, thalidomide and dexamethasone) CYCLE 1 ONLY

MRN:

Ward/Unit:

Name:

Consultant:

DOB:

Address:

NHS No:

Hb

WBC

Plt

Neuts

Na+

K+

Urea

Cr

GFR

Ca

Mg

Alb

AlKP

ALT

Bili

TSH

T4

Cortisol

Height

Weight

BSA

Date

Allergies:

Nature of allergy:

Recorded by

Date

Day No. Date	DRUG or ELECTROLYTE	CALCULATION	DOSE	IV FLUIDS	VOL. MLS.	ROUTE/FLOW RATE	SPECIAL DIRECTIONS/ ADMINISTRATION DETAILS	DRUG ADMINISTRATION	TIME	Pharmacy
22	Dexamethasone		40mg*			PO	1 hour before daratumumab	Sig.		
22	Paracetamol		1g			PO	1 hour before daratumumab			
22	Chlorphenamine		8mg			PO	1 hour before daratumumab			
22	DARATUMUMAB (and 30,000 units of humanised hyaluronidase)		1800mg			SC Over 3-5 Mins				1800mg

Prescriber sig:name:.....Date:Nurse clinical sig:.....name:.....Date:Pharmacist clinical sig:name:.....Date:

D VTD - twice weekly bortezomib (SC daratumumab, bortezomib, thalidomide and dexamethasone) CYCLE 1 ONLY

MRN:	Ward/Unit:
Name:	
DOB:	Consultant:
Address:	
NHS No:	

Proceed Rules: Bloods 96 hours before day 1 ONLY. No FBCs are required during the cycle

Drug	Neuts	Platelets	Renal	Hepatic (bilirubin $\mu\text{mol/L}$)	Neuropathy
Bortezomib	$>1 \times 10^9/\text{L}$	$>75 \times 10^9/\text{L}$	Dialysis may reduce lev-els, therefore it should be administered after dialysis.	$>50 =$ consider $0.7\text{mg}/\text{m}^2$ dose	G1 with pain or G2 = $1\text{mg}/\text{m}^2$ G2 with pain or G3 = Withhold treatment until toxicity resolves and re-initiate at $0.7\text{mg}/\text{m}^2$ G4 = stop treatment
Daratumumab	$>1 \times 10^9/\text{L}$	$>75 \times 10^9/\text{L}$	NA	NA	NA
Thalidomide	$>1 \times 10^9/\text{L}$	$>75 \times 10^9/\text{L}$	NA	NA	G2 and above withhold treatment and re-initiate at a lower dose once toxicity resolves.

Medications to be prescribed on PICS

Anti-emetics	Supportive medication
<ul style="list-style-type: none"> Metoclopramide 10mg TDS prn 	<ul style="list-style-type: none"> Dexamethasone 40mg on days 2, 9, 16 & 23. * Can be reduced to 20mg if required. Aciclovir 400mg BD Co-trimoxazole 480mg BD TTW Allopurinol 300mg OD for the FIRST cycle only Lansoprazole 30mg OD Thalidomide 50mg ON (start at 50mg and increase as clinically indicated) Apixaban 2.5mg BD Senna 15mg ON PRN Levofloxacin 500mg OD

Other Information

- A Prescription Initiation Form (PIF) must be completed for the first cycle treatment of thalidomide.
- A Prescription Authorisation Form (PAF) must be completed for every supply of thalidomide.