



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

The Effects of Lockdown During the Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) Pandemic on Neurotrauma-Related Hospital Admissions

Javier M. Figueroa, James Boddu, Michael Kader, Katherine Berry, Vignesh Kumar, Veronica Ayala, Steven Vanni, Jonathan Jagid

BACKGROUND: The response to the global severe acute respiratory syndrome coronavirus 2 pandemic culminated in mandatory isolation throughout the world, with nationwide confinement orders issued to decrease viral spread. These drastic measures were successful in “flattening the curve” and maintaining the previous rate of coronavirus disease 2019 infections and deaths. To date, the effects of the coronavirus disease 2019 pandemic on neurotrauma has not been reported.

METHODS: We retrospectively analyzed hospital admissions from Ryder Trauma Center at Jackson Memorial Hospital, during the months of March and April from 2016 to 2020. Specifically, we identified all patients who had cranial neurotrauma consisting of traumatic brain injury and/or skull fractures, as well as spinal neurotrauma consisting of vertebral fractures and/or spinal cord injury. We then performed chart review to determine mechanism of injury and if emergent surgical intervention was required.

RESULTS: Compared with previous years, we saw a significant decline in the number of neurotraumas during the pandemic, with a 62% decline after the lockdown began. The number of emergent neurotrauma surgical cases also significantly decreased by 84% in the month of April. Interestingly, although the number of vehicular traumas decreased by 77%, there was a significant 100% increase in the number of gunshot wounds.

CONCLUSIONS: Population seclusion had a direct effect on the frequency of neurotrauma, whereas the change in relative proportion of certain mechanisms may be

associated with the psychosocial effects of social distancing and quarantine.

INTRODUCTION

As of June 1, 2020, the global incidence of coronavirus disease 2019 (COVID-19) was 6.05 million confirmed cases, with 371,000 related deaths. The United States had a major proportion of infections, with 1.7 million confirmed cases and 102,000 related deaths.¹ Specific to our institution, Miami-Dade county had 18,139 confirmed cases with 702 associated deaths. The first case of COVID-19 in Miami-Dade county was confirmed on March 12, 2020, nearly 50 days after the initial case in the United States and 8 days after the initial case in Florida.² A subsequent statewide closing of restaurants and bars on March 17, 2020, was then implemented to decrease viral spread. However, after a significant increase in infection rate over the next few weeks, the Governor of Florida issued an executive “stay-at-home” order on April 1, 2020.³ Although the rate of viral spread improved, we also saw a decrease in both the number of accidents causing traumatic injuries and the number of emergent surgical procedures, secondary to a decline in both foot and automobile traffic.⁴

METHODS

After obtaining approval for this retrospective study from the University of Miami institutional review board, we queried the registry at Ryder Trauma Center to obtain a list of patients from 2016 to 2020 who sustained neurotrauma during the time frame of March 1 to April 30, 2020. Neurotrauma was defined as patients with traumatic brain injury, skull fractures, spinal cord injury, and

Key words

- Admissions
- COVID-19
- Lockdown
- Neurotrauma
- Pandemic

Abbreviations and Acronyms

COVID-19: Coronavirus disease 2019

GSW: Gunshot wound

PHBC: Pedestrian hit by car

Department of Neurosurgery, University of Miami Miller School of Medicine, Miami, Florida, USA

To whom correspondence should be addressed: Javier Figueroa, M.D., Ph.D.

[E-mail: Javier.Figueroa@jhsmiami.org]


Citation: *World Neurosurg.* (2020).

<https://doi.org/10.1016/j.wneu.2020.08.083>

Journal homepage: www.journals.elsevier.com/world-neurosurgery

Available online: www.sciencedirect.com

1878-8750/\$ - see front matter © 2020 Published by Elsevier Inc.




vertebral fractures. Chart review was then performed to obtain variables such as age, sex, mechanism of injury, type of injury, and need for emergent surgery. Mechanisms of injury included assaults, bicycle accidents, ground-level falls (sitting, standing), falls from height (ladder, roof, multiple stories), gunshot wounds (GSWs), motorcycle collisions, motor vehicle collisions, pedestrian hit by car (PHBC), and other (boating, jet-ski, diving). Emergent surgery included procedures such as craniotomy, craniectomy, elevation of depressed skull fracture, skull-base repair for leakage of cerebrospinal fluid, laminectomy for spinal decompression, and instrumented fusions for spinal instability. Data sources included hospital charts and imaging studies.

Data from the previous years 2016–2019 were used to establish baselines for statistical analysis. The absolute number of neurotrauma consults, surgeries, and mechanisms of injury in 2020 were compared with previous monthly averages and analyzed by Poisson regression analysis. The relative differences in proportion of mechanisms of injury in 2020 were compared before monthly averages by χ^2 proportion analysis. All statistical analysis was accomplished on GraphPad Prism 8 scientific software (GraphPad Software, San Diego, California, USA).

RESULTS

Overall, we found a significant difference in the average number of monthly neurotrauma consults from 2016 to 2019, with 83.5 ± 4.7 in March and 68.0 ± 8.8 in April ($P = 0.048$, Student *t* test). However, in March 2020 we saw a 20% decrease in total neurotrauma consults, which was significantly lower ($P = 0.036$, Poisson analysis) than previous years (Figure 1). This declining trend continued in April 2020, with the number of neurotrauma consults decreasing significantly by 62% ($P = 0.0001$, Poisson analysis), after a statewide “stay-at-home” order was issued on April 1, 2020 (Figure 1). Further analysis demonstrated that the number of operative cases for neurotrauma also decreased significantly by 84% ($P = 0.0004$, Poisson analysis) in April 2020 (Figure 2).

We then evaluated the mechanisms of injury for monthly neurotrauma consults and found no significant difference from 2016 to 2020. However, with only 28% of consults in 2020 coming in April, results may have been masked by normal proportions in March. When assessed individually, we found several differences in mechanisms of injury between March and April 2020 (Figure 3). Although absolute numbers of nearly all mechanisms of injury




declined compared with previous-year averages, the relative proportion of each mechanism did not. As expected, the proportion of motorcycle collisions, motor vehicle collisions, and bicycle accidents decreased by 4%, 10%, and 3% respectively. In addition, the proportion of ground-level falls resulting in neurotrauma decreased by 6%, whereas the proportion of falls from height increased by 6%. There was also a nonsignificant 2% increase in the proportion of assaults. Surprisingly, there was a 6% increase in the proportion of PHBC and a 12% significant increase ($P = 0.034$, χ^2 proportion analysis) in the proportion of GSWs. There were no traumatic injuries caused by “other” mechanisms during the pandemic.

DISCUSSION

In the United States, traumatic unintentional injuries are the leading cause of death in people younger than 45 years old and the third-leading cause of death among all age groups combined.⁵ An estimated 1.7 million people sustain traumatic brain injury annually, with approximately 52,000 deaths.⁶ In addition,

although not typically life-threatening, an estimated 18,000 people sustain spinal cord injury annually.⁷ Importantly, the effects of the severe acute respiratory syndrome coronavirus 2 pandemic on the incidence of neurotrauma has yet to be reported. While several hospitals found a decreasing trend in general trauma admissions from February to April 2020, none of these studies thoroughly evaluated the effects on emergent operative cases or changes in mechanisms of injury.⁸⁻¹⁸

Here, we found that the average number of neurotrauma consults differed significantly between March and April, likely secondary to South Florida being a destination for spring break, causing an influx of vacationers during that time frame. During the pandemic, however, travel restrictions in combination with less foot and vehicle traffic led to a decrease in all mechanisms of injury, except for GSW. Upon further investigation, we found that the relative proportion of mechanisms of injury also changed after the lockdown in April 2020. With fewer citizens commuting on the streets, the proportion of vehicular trauma decreased, as expected. Decreases in these types of traumas have been reported across the country, however, not in correlation



with specific events such as initiation of lockdown protocols.^{8,9,12,13} The proportion of ground-level falls also decreased; however, this may have been secondary to patients unwilling to take the risk of going to the emergency department after minor accidents for concern of contracting the virus.¹⁸ Importantly, some businesses were deemed “essential” and allowed to continue operating, of which some construction companies took advantage.⁷ This may explain the increase in the proportion of falls from height, in addition to people doing home repairs while “stay-at-home” orders were in place. With respect to increases in the proportions of assaults and GSWs during the pandemic, they may be secondary to the psychosocial effects of mandatory isolation.^{19,20} Family and friends were forced to be in close proximity to one another, which had the potential to ignite conflicts and violence leading to assault.²¹ Finally, with prolonged confinement comes an increase in the risk of suicide, which may explain the increase in presumed self-inflicted PHBC and GSW.²²

Although the results are compelling, there are several limitations that could be affecting the results of this study. For instance, ambulances may have avoided our hospital, which had a high COVID-19 census, and primary care physicians may have treated minor traumas rather than referring patients to the emergency department. These confounding variables are difficult to address

during the pandemic and must be taken into account when referencing this observational study.

CONCLUSIONS

The severe acute respiratory syndrome coronavirus 2 pandemic significantly impacted the incidence of neurotraumas and associated emergent neurosurgical interventions, as lockdown orders and apprehension of infection decreased the likelihood of certain mechanisms of injury such as vehicular trauma. Conversely, the psychosocial impact of sheltering at home increased the likelihood of interpersonal and self-inflicted trauma. These observations will be helpful if another wave of the epidemic arises.

CRediT AUTHORSHIP CONTRIBUTION STATEMENT

Javier M. Figueroa: Conceptualization, Investigation, Formal analysis, Writing - review & editing. **James Boddu:** Investigation, Formal analysis, Writing - review & editing. **Michael Kader:** Formal analysis, Writing - review & editing. **Katherine Berry:** Formal analysis, Writing - review & editing. **Vignesh Kumar:** Writing - review & editing. **Veronica Ayala:** Writing - review & editing. **Steven Vanni:** Conceptualization, Writing - review & editing. **Jonathan Jagid:** Conceptualization, Writing - review & editing.

05 REFERENCES

1. Fauci AS, Lane HC, Redfield RR. Covid-19—navigating the uncharted. *N Engl J Med*. 2020;382:1268-1269.
2. World Health Organization coronavirus disease (COVID-19) situation report—133. Available at: <https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200601-covid-19-sitrep-133>. Accessed ■■■■.
3. Coronavirus locations: COVID-19 map by county and state. Available at: <https://usafacts.org/visualizations/coronavirus-covid-19-spread-map/>. Accessed ■■■■.
4. Lee H, Park SJ, Lee GR, et al. The relationship between trends in COVID-19 prevalence and traffic levels in South Korea. *Int J Infect Dis*. 2020;96:399-407.
5. Centers for Disease Control and Prevention. National vital statistics reports. Deaths: leading causes for 2017. Available at: https://www.cdc.gov/nchs/data/nvsr/nvsr68/nvsr68_06-508.pdf. Accessed ■■■■.
6. Centers for Disease Control and Prevention. Traumatic brain injury in the United States, 2002-2006. Available at: https://www.cdc.gov/traumaticbraininjury/pdf/blue_book.pdf. Accessed ■■■■.
7. Jain NB, Ayers GD, Peterson EN, et al. Traumatic spinal cord injury in the United States, 1993-2012. *JAMA*. 2015;313:2236-2243.
8. Pichard R, Kopel L, Lejeune Q, Masmoudi R, Masmajea EH. Impact of the CoronaVirus Disease 2019 lockdown on hand and upper limb emergencies: experience of a referred university trauma hand centre in Paris, France [e-pub ahead of print]. *Int Orthop*. <https://doi.org/10.1007/s00264-020-04654-2>, accessed ■■■■.
9. Zagra L, Faraldi M, Pregliasco F, et al. Changes of clinical activities in an orthopaedic institute in North Italy during the spread of COVID-19 pandemic: a seven-week observational analysis [e-pub ahead of print]. *Int Orthop*. <https://doi.org/10.1007/s00264-020-04590-1>, accessed ■■■■.
10. Houshyar R, Tran-Harding K, Glavis-Bloom J, et al. Effect of shelter-in-place on emergency department radiology volumes during the COVID-19 pandemic [e-pub ahead of print]. *Emerg Radiol*. <https://doi.org/10.1007/s10140-020-01797-y>, accessed ■■■■.
11. Antony J, James WT, Neriamparambil AJ, Barot DD, Withers T. An Australian response to the COVID-19 pandemic and its implications on the practice of neurosurgery. *World Neurosurg*. 2020;20, 31107-31104.
12. Bram JT, Johnson MA, Magee LC, et al. Where have all the fractures gone? The epidemiology of pediatric fractures during the COVID-19 pandemic. *J Pediatr Orthop*. 2020;40:373-379.
13. von Dercks N, Körner C, Heyde CE, Theopold J. How badly is the coronavirus pandemic affecting orthopaedic and trauma surgery clinics? An analysis of the first 5 weeks. *Orthopade*. 2020;49:494-501.
14. Hassan K, Prescher H, Wang F, Chang DW, Reid RR. Evaluating the effects of COVID-19 on plastic surgery emergencies: protocols and analysis from a Level I trauma center. *Ann Plast Surg*. 2020;85(suppl 2):S161-S165.
15. Zhu W, Li X, Wu Y, et al. Community quarantine strategy against coronavirus disease 2019 in Anhui: an evaluation based on trauma center patients. *Int J Infect Dis*. 2020;96:417-421.
16. Nuñez JH, Sallent A, Lakhani K, et al. Impact of the COVID-19 pandemic on an emergency traumatology service: experience at a tertiary trauma centre in Spain. *Injury*. 2020;51:1414-1418.
17. Forrester JD, Liou R, Knowlton LM, Jou RM, Spain DA. Impact of shelter-in-place order for COVID-19 on trauma activations: Santa Clara County, California, March 2020. *Trauma Surg Acute Care Open*. 2020;5:e000505.
18. Comelli I, Scioscioli F, Cervellini G. Impact of the COVID-19 epidemic on census, organization and activity of a large urban emergency department. *Acta Biomed*. 2020;91:45-49.
19. de Girolamo G, Cerveri G, Clerici M, Monzani E, Spinogatti F, Starace F. Mental health in the coronavirus disease 2019 emergency—the Italian response [e-pub ahead of print]. *JAMA Psychiatry*. <https://doi.org/10.1001/jamapsychiatry.2020.1276>, accessed ■■■■.
20. Chevance A, Gourion D, Hoertel N, Llorca P, Thomas P, Bocher B. Ensuring mental health care during the SARS-CoV-2 epidemic in France: a narrative review. *Encephale*. 2020;46:193-201.
21. Ashby M. Initial evidence on the relationship between the coronavirus pandemic and crime in the United States. *Crime Sci*. 2020;9:6.
22. Gunnell D, Appleby L, Arensman E, Hawton K, John A, Kapur N. Suicide risk and prevention during the COVID-19 pandemic. *Lancet Psychiatry*. 2020;7:468-471.

Conflict of interest statement: The authors declare that the article content was composed in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received 18 June 2020; accepted 12 August 2020

Citation: World Neurosurg. (2020).

<https://doi.org/10.1016/j.wneu.2020.08.083>

Journal homepage: www.journals.elsevier.com/world-neurosurgery

Available online: www.sciencedirect.com

1878-8750/\$ - see front matter © 2020 Published by Elsevier Inc.