J. Dairy Sci. 106:5196–5196
https://doi.org/10.3168/jds.2023-106-7-5196
© 2023, The Authors. Published by Elsevier Inc. and Fass Inc. on behalf of the American Dairy Science Association®.
This is an open access article under the CC BY license (http://creativecommons.org/licenses/by/4.0/).

Corrigendum to "Effects of supplementing a *Saccharomyces cerevisiae* fermentation product during the periparturient period on performance of dairy cows fed fresh diets differing in starch content" (J. Dairy Sci. 102:3082–3096)

W. Shi,[1] C. E. Knoblock,[1] K. V. Murphy,[1] T. C. Bruinjé,[1] I. Yoon,[2] D. J. Ambrose,[3] and M. Oba[1]
[1]Department of Agricultural, Food and Nutritional Science, University of Alberta, Edmonton, AB, Canada T6G 2P5
[2]Diamond V, Cedar Rapids, IA 52404
[3]Livestock Research and Extension Branch, Alberta Agriculture and Forestry, Edmonton, AB, Canada T6H 5T6

In Table 9, MUN values were incorrect. The correct values are shown in bold below. The authors regret the error.

Table 9. Effects of supplementation of a Saccharomyces cerevisiae fermentation product (SCFP) and carryover effects of starch content of fresh diets (ST) on DMI, BW and BCS changes, milk yield and composition during the post-fresh period (d 24 ± 3 to 44 ± 3 after calving), and the interval from calving to first ovulation postpartum[1]

	LS		HS			*P*-value		
Variable	CON	SCFP	CON	SCFP	SE	ST	SCFP	ST × SCFP
DMI, kg/d	20.7	19.9	20.6	19.8	0.47	0.77	0.10	0.98
Initial BW,[2] kg	576	578	585	588	9.94	0.33	0.78	0.94
BW change, kg/d	0.48	0.61	0.32	0.32	0.28	0.42	0.82	0.82
Initial BCS[2]	2.79	2.83	2.88	2.83	0.06	0.39	0.95	0.44
BCS change, /21d	−0.18	−0.13	−0.15	−0.11	0.03	0.48	0.16	0.77
Yield, kg/d								
Milk	42.6	41.9	39.4	41.2	1.13	0.09	0.59	0.26
Fat	1.56	1.66	1.60	1.63	0.05	0.91	0.21	0.46
CP	1.21	1.24	1.18	1.24	0.03	0.64	0.23	0.64
Lactose	1.98	1.95	1.87	1.96	0.06	0.34	0.53	0.28
Total solids	5.16	5.27	5.05	5.24	0.13	0.63	0.26	0.78
3.5% FCM	43.8	45.2	43.4	44.6	1.17	0.69	0.25	0.94
Solids-corrected milk	39.6	40.9	39.3	40.5	1.04	0.77	0.22	0.96
3.5% FCM/DMI	2.12	2.29	2.14	2.26	0.05	0.91	<0.01	0.61
Milk composition								
Fat, %	3.74[a]	4.00[ab]	4.05[b]	3.95[ab]	0.10	0.19	0.41	0.08
CP, %	2.86	2.95	2.95	2.96	0.04	0.15	0.17	0.27
Lactose, %	4.65	4.63	4.64	4.65	0.02	0.92	0.86	0.63
Total solids, %	12.2[a]	12.6[b]	12.6[b]	12.5[b]	0.13	0.09	0.26	0.08
MUN, mg/dL	**13.7[ab]**	**13.9[ab]**	**14.3[a]**	**13.1[b]**	**0.41**	**0.76**	**0.23**	**0.09**
SCC, 10^3 cells/mL	37.4[a]	88.3[ab]	164[b]	55.5[ab]	42.8	0.28	0.50	0.07
Apparent dietary NE_L[3]								
Mcal/d	40.7	40.8	39.8	40.8	1.16	0.71	0.65	0.68
Mcal/kg of DMI	1.98	2.07	1.96	2.07	0.05	0.84	0.05	0.86
NE_L balance,[4] Mcal/d	2.35	1.36	1.60	1.55	0.74	0.75	0.56	0.59
Days to FOP[5]	23.8	24.6	28.1	22.7	2.46	0.63	0.36	0.21

[a,b]Means in the same row with different superscripts differ significantly ($P < 0.05$) when an interaction of main treatment effect was observed.
[1]LS = low-starch fresh diets; HS = high-starch fresh diets; CON = control diet, no SCFP supplementation; SCFP = SCFP-supplemented diet.
[2]Measured on d 23 and 24 ± 3 before morning feeding.
[3]Calculated according to Dann et al. (2000).
[4]Apparent dietary NEL (Mcal/d) minus NEL required for milk output and maintenance (Mcal/d). The NEL required for milk and maintenance were calculated according to Dann et al. (2000).
[5]Interval from calving to first ovulation postpartum (d).

REFERENCE

Shi, W., C. E. Knoblock, K. V. Murphy, T. C. Bruinjé, I. Yoon, D. J. Ambrose, and M. Oba. 2019. Effects of supplementing a *Saccharomyces cerevisiae* fermentation product during the periparturient period on performance of dairy cows fed fresh diets differing in starch content. J. Dairy Sci. 102(4):3082–3096. https://doi.org/10.3168/jds.2018-15307.