

Solving nonlinear equations in one variable (II)

Mei-Chen Yeh

Yesterday

- Two iterative approaches for solving $f(x) = 0$
 - Bisection
 - Fixed-point iteration

Bisection

- Given a function $f(\cdot)$
- Given a range $[a, b]$
- Repeat computing the middle point until convergence, e.g., $(b-a)/2 < \text{TOL}$
- 猜數字

Bisection: How accurate and how fast?

- The interval length after n bisection steps is: $\frac{b-a}{2^n}$

$$\text{Solution error} = |x_n - x| < \frac{b-a}{2^{n+1}}$$

x_n : the midpoint of the n -th interval

- If we want the error to satisfy $|x_n - x| \leq \varepsilon$, it suffices to have $(b-a)/2^n \leq \varepsilon$, so that

$$n > \log_2 \left(\frac{b-a}{\varepsilon} \right)$$

Bisection: Properties

- Simple 😊
- Safe, robust 😊
- Requires only that f be continuous 😊
- Slow 😞
- Hard to generalize to systems 😞

Fixed-point iteration

- $f(x) = 0 \rightarrow g(x) = x$
- Given $g(\cdot)$
- Given an initial guess x_0
- Repeat evaluating $g(\cdot)$ until convergence, e.g., $|x_{i+1} - x_i| < \text{TOL}$



Not all of the fixed point form may converge!

$$f(x) = x^3 + x - 1 = 0$$

$$g(x) = 1 - x^3$$

$$g(x) = \sqrt[3]{1 - x}$$

$$g(x) = \frac{1 + 2x^3}{1 + 3x^2}$$

0.5000

0.8750

0.3301

0.9640

0.1041

0.9989

0.0034

1.0000

0.0000

1.0000

0.0000

1.0000

0.0000

0.5000

0.7937

0.5909

0.7424

0.6363

0.7138

0.6590

0.6986

0.6704

0.6907

0.6763

0.6866

0.6792

0.6845

0.6807

0.6835

0.6815

0.6829

0.6819

0.6826

0.6821

0.6825

0.6822

0.6824

0.6823

0.6824

0.5000

0.7143

0.6832

0.6823

0.6823

0.6823

0.6823

0.6823

0.6823

0.6823

0.6823

0.6823


0.6823




$$g(x) = 1 - x^3$$

$$g(x) = \sqrt[3]{1-x}$$


$$g(x) = \frac{1+2x^3}{1+3x^2}$$



(a)



(b)



(c)


Figure 1.3 Geometric view of FPI. The fixed point is the intersection of $g(x)$ and the diagonal line. Three examples of $g(x)$ are shown together with the first few steps of FPI. (a) $g(x) = 1 - x^3$ (b) $g(x) = (1-x)^{1/3}$ (c) $g(x) = (1+2x^3)/(1+3x^2)$

Convergence

- $\mathcal{S} = |g'(r)| < 1$

Mean Value Theorem

- Let f be a continuous function on the interval $[a, b]$. Then there exists a number c between a and b such that $f'(c) = (f(b) - f(a)) / (b - a)$



Convergence

- Let x_i denote the iterate at step i . There exists a number c_i between x_i and r such that

$$\begin{aligned} g'(c) &= (g(x_i) - g(r)) / (x_i - r) \\ &= (x_{i+1} - r) / (x_i - r) \end{aligned}$$

$$(x_{i+1} - r) = g'(c)(x_i - r) \quad \text{Define } e_i = |x_i - r|$$

$$e_{i+1} = |g'(c)| e_i$$

Linear convergence

- Let e_i denote the error at step i of an iterative method.
If

$$\lim_{i \rightarrow \infty} \frac{e_{i+1}}{e_i} = S < 1,$$

the method is said to obey **linear convergence** with rate S .

- **Theorem:** Assume that g is continuously differentiable, $g(r) = r$, and $S = |g'(r)| < 1$. Fixed-point iteration converges linearly with rate S to the fixed point r for initial guesses sufficiently close to r .
-

$$x^3 + x - 1 = 0, r \approx 0.6823$$

$$g(x) = 1 - x^3$$

$$g'(x) = -3x^2$$

$$|g'(r)| = 1.3966$$

$$g(x) = \sqrt[3]{1-x}$$

$$g'(x) = \frac{1}{3}(1-x)^{-2/3}(-1)$$

$$|g'(r)| = 0.0716$$

$$g(x) = \frac{1+2x^3}{1+3x^2}$$

$$g'(x) = \frac{6x^2(1+3x^2) - (1+2x^3)6x}{(1+3x^2)^2}$$

$$|g'(r)| = -4.7495\text{e-}005$$

$$|g'(r)| = 1.3966$$

$$g(x) = 1 - x^3$$

$$|g'(r)| = 0.0716$$

$$g(x) = \sqrt[3]{1-x}$$

$$|g'(r)| = -4.7495e-005$$

$$g(x) = \frac{1+2x^3}{1+3x^2}$$

0.5000

0.8750

0.3301

0.9640

0.1041

0.9989

0.0034

1.0000

0.0000

1.0000

0.0000

1.0000

0.0000

0.5000

0.7937

0.5909

0.7424

0.6363

0.7138

0.6590

0.6986

0.6704

0.6907

0.6763

0.6866

0.6792

0.6845

0.6807

0.6835

0.6815

0.6829

0.6819

0.6826

0.6821

0.6825

0.6822

0.6824

0.6823

0.6824

0.5000

0.7143

0.6832

0.6823

0.6823

0.6823

0.6823

0.6823

0.6823

0.6823

0.6823

0.6823

0.6823



Practice

- Explain why the fixed-point iteration $g(x) = \cos(x)$ converges.

$$r \approx 0.74$$

$$g'(x) = -\sin x$$

$$g'(r) = -\sin 0.74 \approx -0.67$$

$$|g'(r)| < 1$$

Bisection vs. Fixed-point iteration

- Which one is faster?
- Depending on $S = |g'(r)|$ is smaller or larger than $1/2$.

Today

- Newton's algorithm
 - A refined version of FPI where S is designed to be zero
- Secant method

The problem revisited

- Want to find solutions of the scalar nonlinear equation $f(x) = 0$
- We denote a solution of the equation (called **root**, or **zero**) by x^* .
- We looked at **bisection** and general **fixed point iterations** to determine solutions x^* of $f(x) = 0$.

Algorithm: Newton's iteration

Given a scalar differentiable function $f(x)$,

1. Start from an initial guess x_0 .
2. For $i = 0, 1, 2, \dots$, compute


$$x_{i+1} = x_i - \frac{f(x_i)}{f'(x_i)}$$


until x_{i+1} satisfies some termination criterion

Newton's iteration

- $f'(x_0) = f(x_0) / (x_0 - x_1)$
- $x_1 = x_0 - f(x_0) / f'(x_0)$
- $x_{i+1} = x_i - f(x_i) / f'(x_i), i = 0, 1, 2, \dots$
- Newton's method is a fixed point iteration with iteration function

$$g(x) = x - f(x) / f'(x)$$





Example

- Find the Newton's Method formula for $x^3 + x - 1 = 0$.
- $f(x) = x^3 + x - 1 = 0$
- $f'(x) = 3x^2 + 1$
- The iteration formula is

$$x_{i+1} = x_i - \frac{x_i^3 + x_i - 1}{3x_i^2 + 1}$$

i	x_i	$e_i = x_i - r $	e_i/e_{i-1}^2
0	-0.70000000	1.38232780	
1	0.12712551	0.55520230	0.2906
2	0.95767812	0.27535032	0.8933
3	0.73482779	0.05249999	0.6924
4	0.68459177	0.00226397	0.8214
5	0.68233217	0.00000437	0.8527
6	0.68232780	0.00000000	0.8541
7	0.68232780	0.00000000	

Order of Convergence (e_i vs. e_{i+1})

The method is

- **linearly convergent** if

$$\lim_{i \rightarrow \infty} \frac{e_{i+1}}{e_i} = S < 1,$$

- **quadratically convergent** if

$$\lim_{i \rightarrow \infty} \frac{e_{i+1}}{e_i^2} = M \text{ (a constant).}$$

Convergence of Newton's method

- Newton's method will converge with an initial guess close to the root.
- Recall that the convergence rate in fixed-point iteration is $|g'(r)|$, will show $|g'(r)| < 1$.

$$g(x) = x - \frac{f(x)}{f'(x)}$$

$$g'(x) = 1 - \frac{f'(x)^2 - f(x)f''(x)}{f'(x)^2} = \frac{f(x)f''(x)}{f'(x)^2}$$

$$g'(r) = 0 \quad (\text{Why?})$$

Theorem: Let f be twice continuously differentiable and $f(r) = 0$. If $f'(r) \neq 0$, then Newton's method is **quadratically convergent** to r , starting with x_0 close to r .

Proof

$$f(r) = f(x_i) + (r - x_i)f'(x_i) + \frac{(r - x_i)^2}{2} f''(x_i)$$

$$0 = f(x_i) + (r - x_i)f'(x_i) + \frac{(r - x_i)^2}{2} f''(x_i)$$

$$-\frac{f(x_i)}{f'(x_i)} = r - x_i + \frac{(r - x_i)^2}{2} \frac{f''(x_i)}{f'(x_i)}$$

$$\left(x_i - \frac{f(x_i)}{f'(x_i)} \right) - r = \frac{(r - x_i)^2}{2} \frac{f''(x_i)}{f'(x_i)}$$

x_{i+1}

$$e_{i+1} = e_i^2 \frac{f''(x_i)}{2f'(x_i)}$$

$$\lim_{i \rightarrow \infty} \frac{e_{i+1}}{e_i^2} = \left| \frac{f''(r)}{2f'(r)} \right|$$

$$\lim_{i \rightarrow \infty} \frac{e_{i+1}}{e_i^2} = M \text{ (a constant).}$$

Secant method

- Replaces the tangent line (the function's derivative) with the secant line.

$$f'(x_i) \approx \frac{f(x_i) - f(x_{i-1})}{x_i - x_{i-1}}$$

$$x_{i+1} = x_i - \frac{f(x_i)(x_i - x_{i-1})}{f(x_i) - f(x_{i-1})}$$

- Two starting guesses are needed to begin the Secant method.




Figure 1.11 Two steps of the Secant Method. Illustration of Example 1.16. Starting with $x_0=0$ and $x_1=1$, the Secant Method iterates are plotted along with the secant lines.

Convergence of Secant method

- The secant method is **superlinearly convergent**, meaning that it lies between linearly and quadratically convergent methods.

Without proof.

$$x_{i+1} = x_i - \frac{f(x_i)(x_i - x_{i-1})}{f(x_i) - f(x_{i-1})}$$

Example

- Apply the Secant method with $x_0 = 0$, $x_1 = 1$ to find the root of $f(x) = x^3 + x - 1$.
- The iteration formula is

$$x_{i+1} = x_i - \frac{(x_i^3 + x_i - 1)(x_i - x_{i-1})}{x_i^3 + x_i - (x_{i-1}^3 + x_{i-1})}$$

i	x_i
0	0.0000000000000000
1	1.0000000000000000
2	0.5000000000000000
3	0.6363636363636364
4	0.69005235602094
5	0.68202041964819
6	0.68232578140989
7	0.68232780435903
8	0.68232780382802
9	0.68232780382802

程式練習

And, please upload your program on moodle.

- Each equation has one root. Please use Newton (or Secant) method to find the root of

$$x^5 + x = 3 \quad \text{①}$$

$$\ln x + 2x^2 = 3 \quad \text{②}$$

輸出至小數點以下四位