

Assignment 3

Aivar Abrashuly

June 13, 2019

Just a quick check, whether we can use the value $\frac{\nu}{D\gamma}$ as a parameter to describe the rotational velocity. From the figure below it seems that the value are not universal




Figure 0.1: Both band structures are for the same value of $\frac{\nu}{D\gamma}=0.2$. However, they are not same, because values $\frac{\nu}{D}$ and γ are differ from each other.

TASK GIVEN

1. Extend your unit cell by applying multiple of them from the previous assignment. Construct a bulk band structure $\omega(k_B)$

2. Given that you construct a scattering matrix from the previous problem, terminate it from the endpoints. It is called a Ribbon problem.

SOLUTION 1

The easiest way I can think for solving such problem is to apply a recursive functions. Initially, I'll make a scattering matrix for $(n - 1)$ unit cell structure and combine it with another unit cell. Finally, I should get a $n \times n$ matrix. See Fig.

$$\begin{pmatrix} b_1 \\ b_2 \\ \vdots \\ b_{2n} \end{pmatrix} = M^{n-1} \begin{pmatrix} a_1 \\ a_2 \\ \vdots \\ a_{2n} \end{pmatrix} \quad (0.1)$$

While for the another unit cell

$$\begin{pmatrix} B_1 \\ B_2 \\ B_3 \\ B_4 \end{pmatrix} = S \begin{pmatrix} A_1 \\ A_2 \\ A_3 \\ A_4 \end{pmatrix} \quad (0.2)$$

At the interface of these structures, we have $B_1 = a_{2n}$ and $A_1 = b_{2n}$. By combining equations (0.1) and (0.2) we can now get $(2n + 2) \times (2n + 1)$ matrix.

$$\begin{pmatrix} b_1 \\ b_2 \\ \vdots \\ b_{2N+2} \end{pmatrix} = M^N \begin{pmatrix} a_1 \\ a_2 \\ \vdots \\ a_{2N+2} \end{pmatrix} \quad (0.3)$$

For simplicity, let us use notation $M = M^n$

The Bloch matrix \tilde{B} can be found using relations

$$\begin{aligned} b_{2n+1} &= a_{2n} e^{i\mathbf{k}_b \cdot \mathbf{a}} \\ b_{2n} &= a_{2n+1} e^{-i\mathbf{k}_b \cdot \mathbf{a}} \end{aligned}$$

where n varies from 1 to N . At the lower and upper boundaries

$$\begin{aligned} b_{2N+2} &= a_1 e^{iN\mathbf{k}_b \cdot \mathbf{b}} \\ b_1 &= a_{2N+2} e^{-iN\mathbf{k}_b \cdot \mathbf{b}} \end{aligned}$$




Figure 0.2: Our recursive strategy.

The bulk band structure, namely $-\log|\det(M - \tilde{B})|$ represented in the figure below. It is important that we considered new unit cell which is $N = 10$ times of a previous unit cell. Also, the other parameters are $a = 0.6m$, $\gamma = 10Hz$, $\omega_R = 800Hz$ and speed of sound $c = 340m/s$.




Figure 0.3: Bulk band structure for no bias velocity.




Figure 0.4: Bulk band structure for $\frac{v}{D} a = 2 m/s$.




Figure 0.5: Bulk band structure for $\frac{v}{D} a = 5 m/s$.

SOLUTION 2

Here in Ribbon problem, let's say that we have a same unit cell and we just need to equate lower and upper bands. To be exact

$$\begin{pmatrix} b_{1'} \\ b_1 \\ \vdots \\ b_{2N} \\ b_{2'} \end{pmatrix} = M \begin{pmatrix} a_{1'} \\ a_1 \\ \vdots \\ a_{2N} \\ a_{2'} \end{pmatrix} \quad (0.4)$$

By using the relations $b_{1'} = a_{1'}$ and $b_{2'} = a_{2'}$, we should transfer it to $2N \times 2N$ matrix

$$\begin{pmatrix} b_1 \\ \vdots \\ b_{2N} \end{pmatrix} = T \begin{pmatrix} a_1 \\ \vdots \\ a_{2N} \end{pmatrix} \quad (0.5)$$

After using "simple" calculations, we can actually find the coefficients of T matrix

$$T_{ij} = M_{ij} + \frac{M_{i2'}[M_{2'j}(1 - M_{1'1'}) + M_{2'1'}M_{1'j}] + M_{i1'}[M_{1'j}(1 - M_{2'2'}) + M_{1'2'}M_{2'j}]}{(1 - M_{1'1'})(1 - M_{2'2'}) + M_{1'2'}M_{2'1'}} \quad (0.6)$$

Below, bulk band structure for Ribbon problem are represented. All the parameters are the same as previous part.




Figure 0.6: Bulk band structure for (Ribbon problem) for no bias velocity.




Figure 0.7: Bulk band structure for (Ribbon problem) for $\frac{v}{D}a = 2 \text{ m/s}$.




Figure 0.8: Bulk band structure for (Ribbon problem) for $\frac{v}{D}a = 5 \text{ m/s}$.

REFERENCES

- [1] Romain Fleury, MS. *Breaking Temporal Symmetries in Metamaterials and Metasurfaces*, August 2015.
- [2] Charles Kittel. *Introduction to Solid State Physics*, 1914.

- [3] Alexander B. Khanikaev, Romain Fleury, S. Hossein Mousavi and Andrea Alu. *Topologically robust sound propagation in an angular-momentum-biased graphene-like resonator lattice*, Nat. Comm. 2015