

The singleton action from the supermembrane¹

Gianguido Dall'Agata², Davide Fabbri², Christophe Fraser³,
Pietro Fré², Piet Termonia² and Mario Trigiante⁴

² Dipartimento di Fisica Teorica, Università di Torino, via P. Giuria 1, I-10125 Torino,
Istituto Nazionale di Fisica Nucleare (INFN) - Sezione di Torino, Italy

³ Wellcome Trust Centre for the Epidemiology of Infectious Diseases, University of Oxford,
South Parks Road, OXFORD OX1 3PS, UK.

⁴ Department of Physics, University of Wales Swansea, Singleton Park,
Swansea SA2 8PP, United Kingdom

Abstract

We derive the free $Osp(8|4)$ singleton action by sending the $M2$ brane to the Minkowski boundary of an $AdS_4 \times \mathcal{M}_7$ background. We do this by means of the solvable Lie algebra parametrization of the coset space. We also give some comments on singleton actions from membranes on $AdS_4 \times G/H$ backgrounds.

¹Talk given by P. T. at the TMR meeting of the project “*Quantum Aspects of Gauge Theories, Supersymmetry and Unification*” TMR contract ERBFMRX-CT96-0045, Corfu September 20–26, 1998. To appear in the proceedings.
This work is also supported in part by EEC under TMR contract ERBFMRX-CT96-0012, in which M. Trigiante is associated to Swansea University.