

expected to proceed as a complex evolution of expansion and cooling of the plasma, perhaps through a mixed parton-hadron phase, until a purely hadronic phase is reached [53]. Therefore, a fully dynamical and microscopic hadronization scheme as proposed in this paper is needed to trace the space-time-dependent cooling and expansion process from the parton to the hadron phase.

References

- [1] P. Hasenfratz and J. Kuti, Phys. Rep. **40**, 75 (1998); R. Goldflam and L. Wilets, Phys. Rev. **D25**, 1951 (1982); H. B. Nielsen and A. Patkos, Nucl. Phys. **B195**, 137 (1982).
- [2] See e.g.: A. H. Mueller (ed.), *Perturbative Quantum Chromodynamics*, Advanced Series on Directions in High Energy Physics, Vol. 5 (World Scientific, Singapore 1989).
- [3] J. Gasser and H. Leutwyler, Phys. Rep. **87**, 77 (1982); E. Witten, Nucl. Phys. **B223**, 422, 433 (1983); J. Gasser and H. Leutwyler, Nucl. Phys. **B250**, 465 (1985); U. Vogl and W. Weise, Prog. Part. Nucl. Phys. **27**, 195 (1991).
- [4] M. A. Shifman, A. I. Vainshtein, and V. A. Zakharov, Nucl. Phys. **B147**, 448 (1979).
- [5] B. Webber, Ann. Rev. Nucl. Part. Sci. **36**, 253 (1986); preprint Cavendish-HEP-943/17.
- [6] F. Karsch, Nucl. Phys. (Proc. Suppl.) **B9**, 357 (1989).
- [7] J. Ellis and K. Geiger, preprint CERN-TH. 95-34.
- [8] K. Geiger, Phys. Rev. **D51**, 2345 (1995).
- [9] R. Friedberg and T. D. Lee, Phys. Rev. **D15**, 1694 (1977); **D16**, 1096 (1977); T. D. Lee, *Particle Physics and Introduction to Field Theory* (Harwood Academic Press, New York, 1981).
- [10] G. Fai, J. Perry and L. Wilets, Phys. Lett. **B208**, 1 (1988).
- [11] G. Krein, P. Tang, L. Wilets and A.G. Williams, Nucl. Phys. **A523**, 548 (1991).
- [12] J. Schechter, Phys. Rev. **D21**, 3393 (1981).