

The expressions for t_i and g_{ij} (eq.(7) in §2) can be obtained from eq.(A4). For example, g_d is given by the coefficients of $|\Delta_d|^2$ in $\Delta F = \Delta F_1 + \Delta F_2$.

The meaning of the J_2 -terms is now obvious. Since we got these terms by picking up $|\Delta_p^R|^2$ in G_{11}^R and G_{22}^R (diagonal components of the Green's functions), only one particle is transferred from (L) to (R) (or (R) to (L)) in this process. This means that one of the electrons consisting of a Cooper pair tunnels to the other side, while the other one is reflected when the Cooper is scattered at the interface. Hence the process including t_{kp} (tunneling matrix element) can lead to the suppression (g_d and g_s) and the interference (g_{ds}) of the superconducting order parameters.

- 1) D.A. Wollman *et al.*, Phys. Rev. Lett. **71**, 2134 (1993) ; D. Brawner and H.R. Ott, Phys. Rev. B **50**, 6530 (1994); A. Mathai *et al.*, Phys. Rev. Lett. **74**, 4523 (1995) ; C.C. Tsuei *et al.*, Phys. Rev. Lett. **73**, 593 (1994) ; I. Iguchi and Z. Wen, Phys. Rev. B **49**, 12388 (1994).
- 2) M. Sigrist and T.M. Rice, J. Phys. Soc. Jpn. **61**, 4283 (1992).
- 3) M. Sigrist, D. Bailey and R. Laughlin, Phys. Rev. Lett. **74**, 3249 (1995).
- 4) K. Kuboki and M. Sigrist, J. Phys. Soc. Jpn., **65** 361 (1996) ; Czech. J. of Phys. **46**-Suppl., 1039-1040 (1996).
- 5) M. Sigrist, K. Kuboki, P.A. Lee, A.J. Millis and T.M. Rice, Phys. Rev. **B53**, 2835 (1996).
- 6) M. Sigrist, K. Kuboki, A.B. Kuklov, D.B. Bailey and R.B. Laughlin, Czech. J. of Phys. **46**-Suppl., 3159-3164 (1996).
- 7) In 2D Bose condensation in a usual sense would not occur. However, there exists a crossover temperature, T_{BE} , below which bosons are almost condensed to the lowest energy state. The true condensation occurs with a small three dimensionality which is always present in real systems.
- 8) In this state there is a (pseudo-) gap in the spin, but not in the charge excitations.
- 9) In this theory the Fermi liquid state is represented as an ordered state where holons are Bose condensed. In this case the system has only fermionic degrees of freedom (spinons), and thus represent a Fermi liquid state.
- 10) A similar result was obtained from quasiclassical theory in M. Matsumoto and H. Shiba, J. Phys. Soc. Jpn., **64**, 3384 (1995).
- 11) S. Yip, Phys. Rev. **B52**, 3087 (1995).
- 12) Y. Suzumura, Y. Hasegawa and H. Fukuyama, J. Phys. Soc. Jpn., **57**, 580 (1988).
- 13) G. Kotliar and J. Liu, Phys. Rev., **B38**, 5142 (1988).
- 14) T. Tanamoto, K. Kuboki and H. Fukuyama, J. Phys. Soc. Jpn., **60**, 3072 (1991).
- 15) H. Fukuyama, H. Kohno, B. Normand and T. Tanamoto, J. Low Temp. Phys., **99**, 429 (1995).
- 16) N. Nagaosa and P.A. Lee, Phys. Rev. Lett., **64** (1990) 2450; P.A. Lee and N. Nagaosa, Phys. Rev. **B46**, 5621 (1992).
- 17) For a review see M. Sigrist and K. Ueda, Rev. Mod. Phys., **63**, 239 (1991).
- 18) C. Bruder, A. van Otterlo and G.T. Zimanyi, Phys. Rev. **B51**, 12904 (1995).
- 19) Yu.S. Barash, A.V. Galaktionov and A.D. Zaikin, Phys. Rev. **B52**, 665 (1995).
- 20) At very low temperature, of the order of $10^{-3}J$, α_s can be negative. However, the GL theory could not be applied to such a low T .
- 21) Y. Tanuma, Y. Tanaka, M. Ogata and S. Kashiwaya, to appear in J. Phys. Soc. Jpn.: In this paper the surface state of the $t-J$ model (surface is faced to vacuum) is studied at $T=0$, using Bogoliubov - de Gennes method based on a Gutzwiller approximation. The \mathcal{T} -breaking state is found at $\varphi = \pi/4$, but not for $\varphi = 0$.
- 22) G.E. Volovik and L.P. Gorkov, Zh. EKsp. Teor. Fiz. **88**, 1412 (1985) [Sov. Phys. JETP **61**, 843 (1985)].

- 23) M. Sigrist, T.M. Rice and K. Ueda, Phys. Rev. Lett., **63**, 1727 (1989).
- 24) Y. Ohashi and T. Momoi, J. Phys. Soc. Jpn, **65**, 3254 (1996).
- 25) M. Sigrist and Y.B. Kim, J. Phys. Soc. Jpn. **63**, 4314 (1994).
- 26) J.R. Kirtley *et al.*, Phys. Rev. **B51**, 12057 (1995).
- 27) A.B. Kuklov and M. Sigrist, Int. J. Mod. Phys. **11**, 1113 (1997).
- 28) A. Amato, Rev. Mod. Phys. **69**, 1119 (1997).
- 29) M. Covington, M. Aprili, E. Paraoanu, L.H. Greene, F. Xu, J. Zhu and C.A. Mirkkin, Phys. Rev. Lett. **79**, 277 (1997).
- 30) M. Fogelström, D. Rainer and J.A. Sauls, Phys. Rev. Lett. **79**, 281 (1997).
- 31) A. Huck, A. van Otterlo and M. Sigrist, Phys. Rev. **B56**, 14163 (1997).

Fig. 1. Schematic phase diagram of the $t-J$ model within a slave-boson mean-field approximation. T_χ , T_{RVB} and T_{BE} are the transition temperature for the bond order, singlet RVB order and the Bose condensation, respectively.

Fig. 2. Interface between d -wave superconductors. The lines on both sides indicate the crystalline a -axis (the c -axis points out of the plane).

Fig. 3. Numerical results of the coefficients in F_I for D/D -junction: (a) t_d and (b) g_{ij} ($i, j = d, s$). Here $\delta = 0.15$, $T = 0.8T_c$, $\tilde{t}^2 = 8.0$ and $\tilde{r}^2 = 0.2$.

Fig. 4 Spatial variation of the order parameters and their relative phase. Here $\delta = 0.15$, $\tilde{t}^2 = 8.0$ and $\tilde{r}^2 = 0.2$ and $\varphi = \pi/4$.

Fig. 5. Phase diagram of surface states in the plane of T and φ . Here $\delta = 0.15$, $\tilde{t}^2 = 8.0$ and $\tilde{r}^2 = 0.2$. The line is for the D/D (S/D)-junction. (The \mathcal{T} -breaking region for D/D -junction is invisible in this scale.)

Fig. 6. Distributions of the surface current J_y (A/cm²) and the local magnetic field B (G). Here $\delta = 0.15$, $T = 0.8T_c$, $\tilde{t}^2 = 8.0$, $\tilde{r}^2 = 0.2$ and $\varphi = \pi/4$.

Fig.7 Numerical results of the coefficients in F_I for S/D -junction. Here $\delta = 0.15$, $T = 0.8T_c$, $\tilde{t}^2 = 8.0$ and $\tilde{r}^2 = 0.2$.

Fig.8 The relative phases between Δ_d and Δ_s (Δ_0), ϕ_{ds} (ϕ_0) as a function of φ .