

Theory of GPCR motion

Meeting I

Aiyan B.

November 20, 2025

Two components of non-interacting GPCR motion

- GPCRs exhibit two non-trivial motions in the membrane:
 - ① Time-averaged anomalous diffusion
 - ② Localization to particular regions; “hot-spots”

Two components of non-interacting GPCR motion

- GPCRs exhibit two non-trivial motions in the membrane:
 - ① Time-averaged anomalous diffusion
 - ② Localization to particular regions; “hot-spots”
- Four physical mechanisms to realize anomalous diffusion¹:
 - ① Fractional Brownian motion
 - ② Continuous time random walks
 - ③ Obstructed diffusion
 - ④ Heterogeneous diffusion

Two components of non-interacting GPCR motion

- GPCRs exhibit two non-trivial motions in the membrane:
 - ① Time-averaged anomalous diffusion
 - ② Localization to particular regions; “hot-spots”
- Four physical mechanisms to realize anomalous diffusion¹:
 - ① Fractional Brownian motion
 - ② Continuous time random walks
 - ③ Obstructed diffusion
 - ④ Heterogeneous diffusion
- Note: Markovian state-switching does NOT allow for anomalous diffusion or localization over long timescales
- Observe heterogeneous behaviours in GPCRs – switching between immobile/confined/free – so the picture that seems the most tempting is 4

Heterogeneous diffusivity

- Instead of keeping the diffusivity κ fixed in space, we promote it to a scalar field $0 < \lambda < \kappa(x) < \Lambda < \infty, \forall x \in \mathcal{M} \subset \mathbb{R}^3$

Heterogeneous diffusivity

- Instead of keeping the diffusivity κ fixed in space, we promote it to a scalar field $0 < \lambda < \kappa(x) < \Lambda < \infty, \forall x \in \mathcal{M} \subset \mathbb{R}^3$
- The corresponding Fokker-Planck equation reads²

$$\frac{\partial p(x, t)}{\partial t} = \nabla \cdot [\kappa(x) \nabla p(x, t)] \quad (1)$$

Heterogeneous diffusivity

- Instead of keeping the diffusivity κ fixed in space, we promote it to a scalar field $0 < \lambda < \kappa(x) < \Lambda < \infty, \forall x \in \mathcal{M} \subset \mathbb{R}^3$
- The corresponding Fokker-Planck equation reads²

$$\frac{\partial p(x, t)}{\partial t} = \nabla \cdot [\kappa(x) \nabla p(x, t)] \quad (1)$$

- With appropriate choice of κ , can lead to anomalous diffusion³

Heterogeneous diffusivity

- Instead of keeping the diffusivity κ fixed in space, we promote it to a scalar field $0 < \lambda < \kappa(x) < \Lambda < \infty, \forall x \in \mathcal{M} \subset \mathbb{R}^3$
- The corresponding Fokker-Planck equation reads²

$$\frac{\partial p(x, t)}{\partial t} = \nabla \cdot [\kappa(x) \nabla p(x, t)] \quad (1)$$

- With appropriate choice of κ , can lead to anomalous diffusion³
- Issues:
 - ① The steady state for Neumann or periodic BC is uniquely the Lebesgue measure \implies cannot observe localization effects!

Heterogeneous diffusivity

- Instead of keeping the diffusivity κ fixed in space, we promote it to a scalar field $0 < \lambda < \kappa(x) < \Lambda < \infty, \forall x \in \mathcal{M} \subset \mathbb{R}^3$
- The corresponding Fokker-Planck equation reads²

$$\frac{\partial p(x, t)}{\partial t} = \nabla \cdot [\kappa(x) \nabla p(x, t)] \quad (1)$$

- With appropriate choice of κ , can lead to anomalous diffusion³
- Issues:
 - 1 The steady state for Neumann or periodic BC is uniquely the Lebesgue measure \implies cannot observe localization effects!
 - 2 Our numerical simulations have shown that anomalous diffusion seems to be uniquely due to κ being bounded

Heterogeneous diffusivity

- Instead of keeping the diffusivity κ fixed in space, we promote it to a scalar field $0 < \lambda < \kappa(x) < \Lambda < \infty, \forall x \in \mathcal{M} \subset \mathbb{R}^3$
- The corresponding Fokker-Planck equation reads²

$$\frac{\partial p(x, t)}{\partial t} = \nabla \cdot [\kappa(x) \nabla p(x, t)] \quad (1)$$

- With appropriate choice of κ , can lead to anomalous diffusion³
- Issues:
 - ① The steady state for Neumann or periodic BC is uniquely the Lebesgue measure \implies cannot observe localization effects!
 - ② Our numerical simulations have shown that anomalous diffusion seems to be uniquely due to κ being bounded
- Tried to rectify both of these by introducing a drift such that $p_\infty(x) \propto e^{-\beta\kappa(x)}$, however this still did not give rise to anomalous diffusive behaviour

So how exactly do we get anomalous diffusion?

- Need one of spatial/temporal correlations to be power-law

So how exactly do we get anomalous diffusion?

- Need one of spatial/temporal correlations to be power-law
- Hot-spots do not get large enough to see anomalous diffusion on the observed timescales \implies heterogeneous diffusion likely not the principal contributor to anomalism

So how exactly do we get anomalous diffusion?


- Need one of spatial/temporal correlations to be power-law
- Hot-spots do not get large enough to see anomalous diffusion on the observed timescales \implies heterogeneous diffusion likely not the principal contributor to anomalism
- Instead, we propose that there are two states – mobile M and immobile S – whose rates depend upon their location

So how exactly do we get anomalous diffusion?

- Need one of spatial/temporal correlations to be power-law
- Hot-spots do not get large enough to see anomalous diffusion on the observed timescales \implies heterogeneous diffusion likely not the principal contributor to anomalism
- Instead, we propose that there are two states – mobile M and immobile S – whose rates depend upon their location
 - Space is partitioned into two domains, high \mathcal{H} and low \mathcal{L} activity, i.e. $\mathcal{M} = \mathcal{H} \cup \mathcal{L}$:

So how exactly do we get anomalous diffusion?

- Need one of spatial/temporal correlations to be power-law
- Hot-spots do not get large enough to see anomalous diffusion on the observed timescales \implies heterogeneous diffusion likely not the principal contributor to anomalism
- Instead, we propose that there are two states – mobile M and immobile S – whose rates depend upon their location
 - Space is partitioned into two domains, high \mathcal{H} and low \mathcal{L} activity, i.e. $\mathcal{M} = \mathcal{H} \cup \mathcal{L}$ ⁴:




So how exactly do we get anomalous diffusion?

- The rate of converting to mobile state from immobile is power-law distributed, i.e.
 $\rho_{M \rightarrow S}(\tau) \sim \tau^{-\alpha-1}$

So how exactly do we get anomalous diffusion?

- The rate of converting to mobile state from immobile is power-law distributed, i.e.
 $\rho_{M \rightarrow S}(\tau) \sim \tau^{-\alpha-1}$
- Rate of conversion to the immobile phase is increased in \mathcal{H} domains, leading to an decrease in α in these regions



References

- [1] D. Krapf, in *Current Topics in Membranes*, Vol. 75 (Elsevier, 2015) pp. 167–207.
- [2] A. W. C. Lau and T. C. Lubensky, *Physical Review E* **76**, 011123 (2007).
- [3] A. G. Cherstvy and R. Metzler, *Physical Review E* **90**, 012134 (2014).
- [4] P. Massignan, C. Manzo, J. A. Torreno-Pina, M. F. García-Parajo, M. Lewenstein, and G. J. Lapeyre, *Physical Review Letters* **112**, 150603 (2014), arXiv:1401.6110 [cond-mat].

The unique steady state for heterogeneous diffusion

- As in the presentation, consider the steady-state equation of the heterogeneous diffusion equation

$$\nabla \cdot [\kappa(x) \nabla p_\infty(x)] = 0 \quad (2)$$

- Let $\tilde{p}_\infty(x) = C$. Clearly $\nabla \tilde{p}_\infty(x) = 0$ so $\tilde{p}_\infty(x)$ is a steady-state solution. Normalization enforces that $C = 1/|\mathcal{M}|$, which specifies \tilde{p}_∞ as the Lebesgue measure on \mathcal{M}
- Let ρ_∞ be any sufficiently smooth solution of Eq. (2) with periodic BC (as in the case of a closed manifold) that is not the Lebesgue measure
- Define the fluctuation field about the mean $\psi(x) := p_\infty(x) - \bar{p}_\infty$. ψ also satisfies Eq. (2) with the same BC

The unique steady state for heterogeneous diffusion

- Multiplying by Eq. (2) by ψ and using integration by parts,

$$\int_{\mathcal{M}} \psi(x) \nabla \cdot [\kappa(x) \nabla \psi(x)] dx = - \int_{\mathcal{M}} \kappa(x) |\nabla \psi(x)|^2 dx = 0, \quad (3)$$

since ψ is periodic and hence vanishes on the boundary.

- But the second representation of the integral is strictly positive since κ is bounded. Therefore the only steady state is \tilde{p}_{∞}