

Mammalia Raccoon Proximity Network Analysis

<https://github.com/aiyshwary/Raccoon-Network-Analysis>

Python

NetworkX

Jupyter

Graph Analysis

Epidemiology

OVERVIEW

A graph-based epidemiological study modelling rabies transmission risk in urban raccoon populations through proximity network analysis and centrality measures, supporting public health and wildlife management decisions.

IMPACT

Constructed and analyzed a raccoon proximity network using real-world contact data, applying centrality measures to identify high-risk individuals and quantify potential rabies spread pathways within urban populations.

TECHNICAL WALKTHROUGH

Mammalia Raccoon Proximity Network Analysis – Project

Explanation

Project Overview

This project analyzes social interaction patterns among urban raccoons using network analysis to understand how rabies and other pathogens can spread within raccoon populations. Since raccoons are a major vector of rabies, studying how often and how closely they interact helps estimate disease transmission risk and supports better public health and wildlife management strategies.

Objective

Model raccoon interactions as a graph/network

Identify high-risk raccoons that can spread pathogens quickly Use centrality measures to understand social structure and disease flow

Dataset & Network Construction

24 raccoons → represented as nodes

~2000 interactions → represented as weighted edges

Data collected using proximity-detecting collars

Edge weight = time spent in close proximity Monthly interaction data used to build the network

If two raccoons are connected, it means they had close physical

interaction, which is critical for rabies transmission.

Why Network Analysis?

Disease spread is not random.

It depends on

Who interacts with whom

How frequently

Who acts as a bridge between groups

Network analysis helps identify super-spreaders and critical connectors.

Centrality Measures Used & Interpretation

Degree Centrality

Measures number of direct connections

High degree = more interactions

Inference

- i) Raccoon #4 has the highest degree

Acts as a high-contact individual

High risk of initiating widespread infection

Closeness Centrality

Measures how close a node is to all others

Indicates how fast something can spread

Inference

- i) Raccoon #2 has highest closeness centrality

- ii) Can spread pathogens rapidly across the entire network

Betweenness Centrality

Measures how often a node lies on shortest paths

Identifies bridge nodes

Inference

- i) Raccoon #16 connects different subgroups

Acts as a key transmission bridge

Removing or vaccinating this raccoon can significantly slow disease spread

Clustering Coefficient

Measures how tightly a raccoon is embedded in a local group

Indicates family or group behavior

Inference

- i) Raccoon #2 has highest clustering

- ii) Infection here may cause localized outbreaks within families

Eigenvector Centrality

Measures influence based on connections to influential nodes

Inference

i) Raccoon #4 again ranks highest

Connected to other highly interactive raccoons

Considered a super-spreader candidate

PageRank Algorithm

Evaluates overall importance considering direction and weight of interactions

Helps rank raccoons by global influence

Key Findings

Raccoon populations are highly interconnected

Disease can spread much faster than expected

A few raccoons play disproportionately large roles in pathogen transmission

Targeted intervention (vaccination/tracking of key raccoons) is more effective than random control

Technologies & Concepts Used

Graph Theory

NetworkX (Python)

Weighted graphs

Centrality algorithms

Epidemiological modeling concepts

KEY DETAILS

1. Built a weighted proximity network from the mammalia-raccoon-proximity dataset using Python and NetworkX.

2. Computed degree, betweenness, closeness, and eigenvector centrality to rank individuals by transmission risk.

3. Identified superspreader nodes whose removal would most disrupt disease propagation chains.

4. Visualized network topology and centrality distributions to communicate findings clearly.

5. Results provide evidence-based recommendations for targeted wildlife intervention strategies.