

Audiometric hearing loss

Across research sites, care homes and participant's homes

Characteristic	Overall N = 423 ¹	Has Hearing Loss N = 196 ¹	Normal Hearing N = 227 ¹	p- value ²	q- value ³
Age	67 (60, 76)	75 (69, 82)	62 (56, 67)	<0.001	<0.001
Unknown	2	2	0		
Gender				>0.9	>0.9
Declined to respond	2 (0.5%)	1 (0.5%)	1 (0.4%)		
Female	229 (54%)	106 (54%)	123 (54%)		
Male	192 (45%)	89 (45%)	103 (45%)		
Occupation				0.011	0.018
Higher managerial, administrative and professional occupations	155 (50%)	72 (45%)	83 (55%)		
Intermediate occupations	68 (22%)	32 (20%)	36 (24%)		
Lower supervisory and technical occupations	25 (8.0%)	15 (9.4%)	10 (6.6%)		
Never worked and long-term unemployed	1 (0.3%)	1 (0.6%)	0 (0%)		
Semi-routine and routine occupations	39 (13%)	29 (18%)	10 (6.6%)		
Semi-routine and routine occupations, Intermediate occupations	2 (0.6%)	2 (1.3%)	0 (0%)		
Small employers and own account workers	21 (6.8%)	8 (5.0%)	13 (8.6%)		
Unknown	112	37	75		

¹ Median (Q1, Q3); n (%)

² Wilcoxon rank sum test; Fisher's exact test; Pearson's Chi-squared test

³ False discovery rate correction for multiple testing

Audiometric hearing loss

Across research sites, care homes and participant's homes

Characteristic	Overall N = 423 ¹	Has Hearing Loss N = 196 ¹	Normal Hearing N = 227 ¹	p- value ²	q- value ³
Were participants born in the UK?	294 (92%)	151 (93%)	143 (92%)	0.7	0.9
Unknown	104	33	71		
Language	308 (97%)	158 (96%)	150 (97%)	0.8	>0.9
Unknown	104	32	72		
Study Location				<0.001	<0.001
Research Site	319 (75%)	152 (78%)	167 (74%)		
Participant's home	84 (20%)	24 (12%)	60 (26%)		
Care Home	20 (4.7%)	20 (10%)	0 (0%)		
Self Reported Hearing Loss				<0.001	<0.001
Yes	138 (36%)	97 (57%)	41 (19%)		
No	237 (61%)	68 (40%)	169 (78%)		
Unknown/unsure	12 (3.1%)	4 (2.4%)	8 (3.7%)		
Unknown	36	27	9		
Difficulty hearing the television				<0.001	<0.001
Yes	113 (33%)	87 (60%)	26 (13%)		

¹ Median (Q1, Q3); n (%)

² Wilcoxon rank sum test; Fisher's exact test; Pearson's Chi-squared test

³ False discovery rate correction for multiple testing

Audiometric hearing loss

Across research sites, care homes and participant's homes

Characteristic	Overall N = 423 ¹	Has Hearing Loss N = 196 ¹	Normal Hearing N = 227 ¹	p- value ²	q- value ³
No	227 (66%)	58 (40%)	169 (86%)		
Declined question	2 (0.6%)	1 (0.7%)	1 (0.5%)		
Unknown	81	50	31		
Need to speak loudly				0.033	0.049
Yes	236 (69%)	109 (74%)	127 (66%)		
No	95 (28%)	32 (22%)	63 (33%)		
Don't Know	8 (2.3%)	6 (4.1%)	2 (1.0%)		
Refused to answer	2 (0.6%)	1 (0.7%)	1 (0.5%)		
Unknown	82	48	34		
Difficulty hearing in group conversation	140 (41%)	93 (63%)	47 (24%)	<0.001	<0.001
Unknown	81	49	32		
Do you wear a hearing aid?	74 (19%)	68 (40%)	6 (2.8%)	<0.001	<0.001
Unknown	41	28	13		
Do they report tinnitus	110 (34%)	61 (38%)	49 (31%)	0.2	0.2
Unknown	101	34	67		

¹ Median (Q1, Q3); n (%)

² Wilcoxon rank sum test; Fisher's exact test; Pearson's Chi-squared test

³ False discovery rate correction for multiple testing