



Review article

Management of HER2 alterations in non-small cell lung cancer – The past, present, and future

Jorn Nützinger^{a,1}, Jii Bum Lee^{b,1}, Jia Li Low^a, Puey Ling Chia^c, Silvana Talisa Wijaya^a,
Byoung Chul Cho^b, Sun Min Lim^b, Ross A Soo^{a,*}

^a Department of Haematology-Oncology, National University Cancer Institute, Singapore

^b Division of Medical Oncology, Department of Internal Medicine, Yonsei University College of Medicine, Seoul, South Korea

^c Department of Medical Oncology, Tan Tock Seng Hospital, Singapore



ARTICLE INFO

Keywords:

HER2

Non-small cell lung cancer

Exon 20 mutation

Targeted therapy

ABSTRACT

HER2 mutations, which account for 2–4% of non-small cell lung cancer (NSCLC), are distinct molecular alterations identified via next generation sequencing (NGS). Previously, treatment outcomes in *HER2*-mutant metastatic NSCLC were dismal, showing limited clinical benefit with platinum-based chemotherapy with or without immunotherapy. In contrast to *HER2*-altered breast and gastric cancer, *HER2*-mutant NSCLC does not benefit from *HER2* targeting agents such as trastuzumab or TDM1. *HER2* mutations are also inherently different from *HER2* overexpression and amplification. Currently, trastuzumab deruxtecan, a *HER2* targeting antibody drug conjugate (ADC) is the first and only approved treatment option for patients with *HER2*-mutant metastatic NSCLC after failure with standard treatment. In this review, we summarized the biology of *HER2* and detection of *HER2* overexpression, amplification and mutations, as well as general landscape of landmark and ongoing clinical trials encompassing from chemotherapy to targeted agents, including tyrosine kinase inhibitors (TKIs), ADCs and investigational agents.

1. Introduction

Non-small cell lung cancer (NSCLC) is a heterogeneous disease composed of distinct driver oncogenes such as mutations of the epidermal growth factor receptor gene (*EGFR*), Kirsten rat sarcoma virus gene (*KRAS*), and gene rearrangement of anaplastic lymphoma kinase (*ALK*) and *ROS1*. Molecular targeted therapies targeting such molecular alterations have led to major changes in the treatment landscape for NSCLC.

The Human epidermal growth factor receptor 2, also known as *HER2* or *erbB-2/neu*, a well-established oncogene in breast cancer, has recently been identified as a therapeutic target in NSCLC. Several *HER2* activating mechanisms have been described including gene mutation,

amplification, and overexpression [1].

HER2 mutations in NSCLC were first discovered in 2004 [2], however until recently, developing effective anti-*HER2* therapies has been a challenge (Fig. 1) [3]. In August 2022, based on the data from the DESTINY-Lung02, the U.S Food and Drug Administration (FDA) granted accelerated approval to the antibody drug conjugate trastuzumab deruxtecan, making it the first targeted agent approved for *HER2*-mutant NSCLC.

In this review, we provide an overview of the biology of *HER2* including various alterations of *HER2*, their detection methods, and the treatment of *HER2*-mutant NSCLC (Fig. 2).

Abbreviations: ADC, antibody drug conjugate; ADCC, antibody-dependent cellular cytotoxicity; BID, twice daily; CNS, central nervous system; DAR, drug-antibody ratio; DoR, duration of response; ECD, extracellular domain; EGFR, epidermal growth factor receptor; FDA, Food and Drug Administration; FISH, fluorescence in situ hybridization; ICI, immune checkpoint inhibitor; IHC, immunohistochemistry; ILD, interstitial lung disease; mAbs, monoclonal antibodies; NGS, next generation sequencing; NSCLC, non-small cell lung cancer; ORR, overall response rate; OS, overall survival; PFS, progression-free survival; QD, once daily; qPCR, quantitative polymerase chain reaction; qRT-PCR, real-time quantitative reverse transcriptase polymerase chain reaction; T-DXd, trastuzumab deruxtecan; TEAE, Treatment-emergent adverse event; TKD, tyrosine kinase domain; TKIs, tyrosine kinase inhibitors; TMB, tumor mutational burden; TMD, transmembrane domain.

* Corresponding author.

E-mail address: ross_soo@nuhs.edu.sg (R.A. Soo).

¹ These authors contributed equally as first authors.

<https://doi.org/10.1016/j.lungcan.2023.107385>

Received 6 August 2023; Received in revised form 21 September 2023; Accepted 22 September 2023

Available online 28 September 2023

0169-5002/© 2023 Elsevier B.V. All rights reserved.

2. Biology and detection of *HER2*

2.1. Biology

Discovered in the mid 1980's (Fig. 1), the *HER2* proto-oncogene is located on the long arm of human chromosome 17 (17q11.2-q12) [4]. The *HER2* transcript encodes a transmembrane glycoprotein of 185 kDa (kD) belonging to the ErbB-family of type 1 transmembrane growth factor receptors which consists of HER1 (ErbB1, EGFR), HER2 (ErbB2, HER2/neu), HER3 (ErbB3) and HER4 (ErbB4). Each receptor is composed of three main components: an extracellular ligand binding domain, an α -helical transmembrane segment and an intracellular tyrosine kinase domain. Under physiologic conditions, upon ligand binding to their extracellular domains, HER proteins undergo hetero- or homodimerization between the receptors hereby adhering to a distinct hierarchical order that favors heterodimers over homodimers. Subsequent intracellular transphosphorylation of tyrosine residues initiates a variety of signaling pathways including MAPK, PI3K/AKT, PKC and STAT, leading to cellular proliferation, differentiation, and migration. *HER2* exhibits significant structural homology with all other members of its family, but, unlike other members, no direct ligand has been identified for *HER2* [5]. Instead, *HER2* remains in its active conformation, enabling continuous availability for dimerization, therefore rendering it the favored partner for heterodimerization. *HER2* also displays reduced internalization and degradation, leading to prolonged activation on the cell membrane [6]. Combined with *HER3*, which is devoid of intrinsic tyrosine kinase activity and therefore restricted to heterodimerizing with other HER-receptors, the *HER2/HER3* heterodimer is the most potent in terms of strength of their interaction, ligand-induced tyrosine phosphorylation and downstream signaling [7].

Its superior capability for heterodimerization, combined with a distinctive basal tyrosine kinase activity, grants *HER2* a crucial function in signal transduction and consequently plays a significant role in carcinogenesis when its function is dysregulated (Fig. 2).

2.2. *HER2* alterations

In contrast to breast cancer which is triggered by *HER2* amplification, NSCLC can be activated by gene mutation, amplification, and overexpression of the *HER2* leading to downstream activation of the MAPK, PI3K/Akt, phospholipase C, protein kinase C, and the Janus kinase (Jak-STAT) signaling pathways [8,9]. In the following sections, we

will discuss the different alterations and their detection methods.

2.3. *HER2* overexpression

Prevalence ranges widely from 2.4 to 38 % and is more common in well-differentiated adenocarcinomas [10–12]. The identification and distinction remain debatable due to difference in applied methods, patient groups studied and the different definitions of *HER2* positivity [13–15].

The underlying molecular pathogenesis is unclear, but increased *HER2* presence on the cell surface increases the formation of *HER2*-containing heterodimers, consequently altering cell polarity and adhesion and activating several above-mentioned oncogenic signaling pathways [16]. The mechanism of overexpression is attributed to increased copy number from chromosome duplication and polysomy (*HER2* gene copy number higher than 5 or 6, but *HER2/CEP17* < 2) [13,17]. Bunnet et al. discovered a strong correlation between *HER2* protein expression, assessed by IHC, and *HER2* gene copy number, evaluated by fluorescence in situ hybridization (FISH). Of the cell lines tested, 32 % were *HER2* IHC positive. Interestingly, polysomy, rather than genuine amplification, was often detected by FISH. Therefore, in case of moderate or strong *HER2* overexpression an additional FISH analysis is advised to distinguish between these possibilities [6,18]. *HER2* overexpression is associated with poorer survival, which may be due to chemo-radiotherapy resistance, but results are conflicting [19–21].

There is no consensus on scoring and categorizing *HER2* protein overexpression in lung cancer and a universal definition and test are urgently needed. In the absence of a specific testing method, two immunohistochemistry (IHC) scoring systems are used, including scoring using 0–3+ and the H-score [22]. Another method for quantifying *HER2* overexpression is the use of quantitative polymerase chain reaction (qPCR) or real-time quantitative reverse transcriptase polymerase chain reaction (qRT-PCR).

2.4. *HER2* amplification

HER2 amplification in NSCLC occurs less frequently compared to breast cancer [13], but the reported frequency varies due to a lack of a clear definition and the existence of two distinct mechanisms: amplification as a primary driver in patients who did not receive targeted therapy (3 %) and amplification from acquired EGFR-TKI resistance (13 %) [23].




Fig. 1. Timeline of development of *HER2* targeted therapies in *HER2*-mutant NSCLC.

HER2 amplification increases NSCLC cell invasiveness in vitro and constitutively activates both *HER2* and *EGFR* independent of ligands [24]. It is strongly associated with pleural invasion [1], never smoking and female sex [25,26] but its prognostic significance is unclear [42].

The definition of *HER2* amplification is derived from breast cancer diagnostics and is typically defined as the average ratio of the *HER2* gene copy number to centromeres (*HER2*/CEP ratio ≥ 2) and/or gene copy number > 6 . It must be noted that other cut off values have also been used, resulting in variations among different studies [18]. Furthermore, it is posed to modify the current used breast cancer classification as it may not be adequate for NSCLC [27]. As stated earlier, polysomy can be found and while it is believed not to drive oncogenesis, its predictive or prognostic value is unclear [14].

FISH is used as the gold standard for detection [28,29] and evaluation of *HER2* amplification is advised in clinical studies or in case of *EGFR* TKI resistance [30]. Another method to assess *HER2* amplification in clinical practice is NGS, which also lacks universal standard for defining amplification across NGS platforms but may offer advantages because it can simultaneously test alterations in several hundred genes [31].

2.5. *HER2* mutations

HER2 mutations comprise heterogeneous alterations in the extracellular (ECD), transmembrane (TMD), and tyrosine kinase domains (TKD). They are found in 2–4 % of NSCLC [1,5,32], have low genetic diversity and are associated with female gender, never smokers, adenocarcinoma histology and brain metastases [33]. *HER2*-mutant NSCLC patients have a worse prognosis than patients with *EGFR* mutations or *ALK* rearrangements due to a lack of effective selective targeted agents.

HER2 mutations occur mostly in the intracellular tyrosine kinase domain (TKD) and induce conformational changes in the ATP-binding pocket, which increases kinase activity and downstream induction of PI3K/AKT and MEK oncogenic pathways [34]. The most frequent mutations occur as in-frame insertions within exon 20. The 12 base-pair duplicated insertion at codon 775 - A775_G776insYVMA, which affects the α C-B4 loop of the kinase domain, is the most common followed by G776delinsVC and G778_P780insGSP [5,35,36]. The exon 20 YVMA

mutation, in particular, is associated with a high incidence of brain metastasis and inferior outcome of chemotherapy [37]. *HER2* exon 20 insertions share structural homology with *EGFR* exon 20 insertions and affect the same structural regions: the α C-helix, at residues 770–774, and the loop region comprising residues 775–783. Similar to *EGFR*, *HER2* activity is regulated by the α C-helix, which switches between an inactive, outward conformation and an active, inward conformation. *HER2* exon 20 insertions exhibit greater activation of signal transduction, *EGFR* phosphorylation, cell proliferation, and invasiveness compared to wild-type *HER2* signaling. Different insertions have unique behavior and drug resistance profiles depend on their conformational configuration (Figs. 3 and 4) [36].

More TKD-affecting point mutations exist with lower prevalence and more recent a few less common mutations affecting the transmembrane and the juxtamembrane domains (G660D, R678Q, E693K and Q709L) have been described and can be targeted [38,39].

Although earlier studies have proposed *HER2* mutations to be mutually exclusive to other oncogenic drivers, the widespread use of NGS has revealed accompanying mutations in *EGFR* mutations, *ALK* translocations and *ROS* translocations [40]. While most studies do not detail the use of previous targeted treatments, a small percentage of *HER2* mutations have been described as a possible mechanism of primary or acquired resistance during *EGFR* TKI treatment [18,41,42].

As stated earlier, *HER2* overexpression detected by FISH or IHC staining is not a reliable substitute marker for *HER2* mutation [14] and although targeted assays like RT-PCR are highly specific, they are limited in their ability to detect only known and well characterized gene alterations, potentially missing less common alterations. Therefore, sequencing methods remain the gold standard when testing for *HER2* mutations [30].

2.6. Genomic profile and tumor microenvironment of *HER2*-mutant NSCLC

Whole exome sequencing of NSCLC harbouring *HER2* exon 20 insertion have revealed low tumor mutational burden (TMB) and few co-driver mutations, findings similar to *EGFR*-mutated NSCLC [43]. Other non-exon 20 insertion *HER2* mutations show features similar to *EGFR*-




Fig. 2. Mechanism of action for *HER2* targeted therapies in *HER2*-mutant NSCLC.

HER2 variant	Afatinib [35]	Dacomitinib [35]	Neratinib [35]	Pozotinib [35]	Pyrotinib [35]	Osimertinib [35]	Tucatinib [84]	Furmonertinib [72]	Mobocertinib [114]	BAY2927088 [114]	TAS0728 [82]	JB1-08-176-01 [82]	JIN-A04 [83]	ELVN-002 [84]
WT EGFR	20.2	19.26	38.75	6.49	639.95	142.27	>10,000	162.6	15.9	221	399.15	368.14	NA	>10,000
HER2 S310F	NA	NA	NA	NA	NA	NA	11	NA	0.8	2.8	238.2	41.49	NA	3
HER2 S310Y	NA	NA	NA	NA	NA	NA	12	NA	NA	NA	NA	NA	NA	3
HER2 S335C	NA	NA	NA	NA	NA	NA	NA	NA	2.5	5.9	NA	NA	NA	NA
HER2 V659E	NA	NA	NA	NA	NA	NA	NA	NA	10.1	13.4	NA	NA	NA	NA
HER2 R678Q	NA	NA	NA	NA	NA	NA	29	NA	NA	NA	NA	NA	NA	5
HER2 L755P	20.54	25.16	19.71	3.65	55.18	69.58	1284	NA	5.7	11.6	NA	NA	NA	21
HER2 L755S	4.37	12.49	3.28	0.53	41.6	12.41	418	NA	5.9	1.9	3757	139.92	NA	8
HER2 D769N	0.81	2.65	0.51	0.12	4.53	8.1	7	NA	NA	NA	NA	NA	NA	2
HER2 D769Y	1.48	3.67	1.02	0.36	0.78	7.38	NA	NA	NA	NA	NA	NA	NA	NA
HER2 D769H	2.71	7.04	2.5	0.35	5.36	13.8	NA	NA	NA	NA	NA	NA	NA	NA
HER2 Y772dupYVMA	42.05	143.43	31.08	1.87	118.8	577.77	NA	NA	NA	NA	NA	NA	NA	NA
HER2 V773M	3.27	15.19	2.99	1.07	0.57	29.44	64	NA	NA	NA	NA	NA	NA	4
HER2 M774delinsWLV	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	894.16	21.03	NA	NA
HER2 L775A	NA	NA	NA	NA	NA	NA	NA	NA	4.7	7	NA	NA	NA	NA
HER2 V777L	1.17	3.42	1.62	0.13	0.74	6.96	11	NA	NA	NA	19.24	2.71	NA	3
HER2 A775_G776insYVMA	NA	NA	NA	NA	NA	NA	225	118	NA	NA	256.45	13.58	11.1	11
HER2 775_757LREdelinsRP	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	7.24	NA	NA
HER2 A775-G776-ins-C	NA	NA	NA	NA	NA	NA	24	NA	NA	NA	NA	NA	NA	2
HER2 A775-G776-ins-YVMS	NA	NA	NA	NA	NA	NA	510	NA	NA	NA	NA	NA	NA	15
HER2 A775-G776-ins-SVMA	NA	NA	NA	NA	NA	NA	157	NA	NA	NA	NA	NA	NA	6
HER2 A775-G776-ins-VVMA	NA	NA	NA	NA	NA	NA	294	NA	NA	NA	NA	NA	NA	12
HER2 A775-G776-ins-MIMAY	NA	NA	NA	NA	NA	NA	287	NA	NA	NA	NA	NA	NA	7
HER2 A775-G776-ins-YVMA-R678Q	NA	NA	NA	NA	NA	NA	642	NA	NA	NA	NA	NA	NA	14
HER2 G776delinsVC	29.34	84.87	33.59	4.85	89.82	412.6	NA	NA	NA	NA	625.55	6.01	0.3	NA
HER2 G776delinsLC	7.01	25.11	6.55	1.13	11.42	46.96	88	NA	NA	NA	NA	NA	NA	13
HER2 G776delinsVV	14.9	45.37	20.85	2.99	1663.65	188.06	1239	NA	NA	NA	NA	NA	NA	34
HER2 A776VC	NA	NA	NA	NA	NA	NA	499	NA	NA	NA	NA	NA	NA	17
HER2 P776delinsLC	NA	NA	NA	NA	NA	NA	1104	NA	NA	NA	NA	NA	NA	41
HER2 P776-V777delinsCVC	NA	NA	NA	NA	NA	NA	209	NA	NA	NA	NA	NA	NA	13
HER2 G776-delinsAVGC	NA	NA	NA	NA	NA	NA	438	NA	NA	NA	NA	NA	NA	14
HER2 L777L	NA	NA	NA	NA	NA	NA	NA	NA	1	1.1	NA	NA	NA	NA
HER2 V777_G778insGC	NA	NA	NA	NA	NA	NA	20	25	NA	NA	NA	NA	NA	5
HER2 G778insLPS	1.67	3.39	1.99	0.18	0.96	16.9	NA	NA	NA	NA	NA	NA	NA	NA
HER2 P780insGSP	27.59	95.57	32.27	4.12	23.1	440.31	NA	NA	NA	NA	80.08	7.38	1.4	NA
HER2 P780-P781insGSP	NA	NA	NA	NA	NA	NA	29	NA	NA	NA	NA	NA	NA	3
HER2 L786V	5.02	9.78	1.41	1.22	2.26	8.3	NA	NA	NA	NA	NA	NA	NA	NA
HER2 T789M	NA	NA	NA	NA	NA	NA	3412	NA	NA	NA	NA	NA	NA	194
HER2 V842I	4.05	6.48	1.51	0.45	0.65	24.31	21	NA	NA	NA	NA	NA	NA	4
HER2 V862I	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	1315.4	284.03	NA	NA
HER2 L869R	1.44	3.07	0.96	0.81	4.56	27.61	148	NA	NA	NA	NA	NA	NA	2
HER2 L869R/T7987I	NA	NA	NA	NA	NA	NA	2524	NA	NA	NA	NA	NA	NA	43

Fig. 3. Heat map representation of IC₅₀ for proliferation (nmol/L) of indicated TKIs for Ba/F3 cells harboring each HER2 mutation subtypes, including non-selective pan-HER TKIs (afatinib [35], dacomitinib [35], neratinib [35], poziotinib [35], pyrotinib [35], osimertinib [35], tucatinib [84], furmonertinib [72], mobocertinib [114], BAY2927088 [114]) and selective HER2 TKIs (TAS0728 [82], JBI-08-176-01 [82], JIN-A04 [83], ELVN-002 [84]). For NVL-330 and ENT-H1 there was no detailed IC₅₀ data available at the time of publication. The colors mark the sensitivity to different agents: sensitive (green, <10), intermediate (yellow, 10–99) and resistant (red, ≥100). (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

wild type NSCLC with relatively higher TMB, co-altering mutations, and smoking mutational signature. Concurrent mutations such as *TP53* and *PIK3CA* alterations are identified with *HER2* mutations [44].

HER2-altered NSCLC has an uninfamed TME, is associated with lower rates of high PD-L1 expression and a lower median tumor mutational burden (TMB) [74,76,90]. *HER2* exon 20 mutated NSCLC express low levels of mature NK cells, and upregulate *TNFSF4* (*OX-40*) and *CXCR4*. Preclinical studies show that HER2-altered NSCLC inhibit cAS-STING pathway regulating innate immune mechanism, to prevent TBK1-STING signaling [45]. Taken together, these characteristics result in immune-cold environment in HER2-altered NSCLC.

3. HER2 targeted therapy

3.1. Chemotherapy

In patients with *HER2*-mutant NSCLC treated with first line pemetrexed based chemotherapy, the overall response rate (ORR) was 36 % and the PFS of 5.1 months, similar to the *KRAS*-mutant and *EGFR*-mutant group with a PFS of 5.0 and 6.5 months, respectively but was

significantly shorter than the *ALK/ROS1*-rearranged group with a PFS of 9.2 months ($p = 0.004$). Among the *HER2* mutation variants, there was also a trend towards inferior PFS in A775_G776insYVMA group compared with other variants (4.2 vs 7.2 months, $p = 0.085$) [46].

Several phase II studies have reported the efficacy of chemotherapy in combination with HER2 targeting agents (Table 1). Treatment was well tolerated and results were consistent with those from chemotherapy alone [47]. Similarly, a randomized phase 2 study assessed the addition of trastuzumab to cisplatin and gemcitabine in HER2 positive NSCLC with no clear clinical benefit [48]. In the subset of NSCLC with HER2 overexpression of + 3, the ORR was 25 % [49] but these results should be viewed with caution given the small sample size.

3.2. Immunotherapy

While ICI therapy has become a key component of the treatment for non-oncogene driver NSCLC, its role in oncogene-driven NSCLC such as those with HER2 alterations is less certain. Responses to ICI as a monotherapy appear to be modest at best, with ORR ranging from 0 to 27.3 % and PFS ranging from 1.88 to 2.5 months [44,50–53]. The




Fig. 4. ORR of HER2 agents versus HER2 mutation subtypes. Trastuzumab deruxtecan [99,101], trastuzumab emtansine [92,94–96], trastuzumab [88], trastuzumab and pertuzumab [90], afatinib [58,115], dacomitinib [61], poziotinib for treatment naïve* [35,116–118], poziotinib for treatment refractory** [66], pyrotinib [68], and pyrotinib and apatinib [69].

IMMUNOTARGET registry reported an ORR of 7.4 % and PFS of 2.5 months [52], whereas The French Lung Cancer Group reported an ORR of 27.3 % and a median PFS of 2.2 months [53].

A retrospective study of first-line ICI plus chemotherapy reported an ORR of 52 % and a median PFS of 6 months [54], results comparable to that of an unselected NSCLC cohort in KEYNOTE-189 [55]. Another retrospective study reported an ORR of 28.9 % and median PFS of 5.2 months. While the ORR was higher, there was no statistically significant improvement in median PFS compared to chemotherapy alone (5.2 vs 4.03 months, $p = 0.20$) [56].

Overall, the current evidence does not support the use of ICIs as a single agent in the treatment of HER2-altered NSCLC. Immunotherapy combinations, albeit with limited supporting evidence, remain a viable first-line treatment option.

3.3. HER2 targeted tyrosine kinase inhibitors (TKIs)

3.3.1. Non-selective HER2 inhibitors

Non-selective HER2 inhibitors include the pan-HER TKIs: afatinib, dacomitinib, neratinib, poziotinib, pyrotinib, tucatinib, furmonertinib, mobocertinib, BAY2927088, tarloxotinib and tuxobertinib (Figs. 2, 3).

Afatinib

Afatinib, an irreversible pan-HER TKI, has modest clinical activity with a partial response (PR) rate of 13–19 % [57]. In a single arm phase 2 study of afatinib in pre-treated patients with HER2 exon 20 mutations (ETOP NICHE), the ORR was 0 %, and the study was halted as it did not meet prespecified efficacy threshold [58].

Responses to afatinib may vary according to the HER2 mutation variant (Fig. 4). The most frequent variant, p.A775_G776insYVMA, found in up to 42 % of HER2 mutations has been reported to be resistant to afatinib [59]. Afatinib may be active in non-YVMA variants such as p.

G776delinsVC, p.Y772_A775dup and p.G778_P780dup with a higher ORR (50 % vs 0 %, $p = 0.077$) and longer PFS (9.53 vs 1.8 months, $p = 0.057$) than other HER2 mutation subtypes [60]. TP53 co-mutations and PI3K/AKT/mTOR pathway activations may also confer additional resistance to afatinib [59].

Dacomitinib

In a phase 2 trial of dacomitinib, an irreversible pan-HER TKI, the PR rate was 12 % in patients harbouring HER2 mutations and 0 % with HER2 amplification. No responses were seen in the common YVMA subtype [61]. The IC₅₀ of dacomitinib was higher in Ba/F3 cells harbouring YVMA insertion whereas the IC₅₀ was lower with HER2 variants with GSP insertion, WLW insertion, and CPG insertion (Fig. 3).

Neratinib

Neratinib, an irreversible pan-HER TKI, was evaluated in the SUMMIT phase II basket trial across multiple cancer types with HER2 or HER3 mutations. Of the 26 patients with NSCLC, only one patient with a kinase domain missense mutation (L755S) had a PR [3].

A randomized phase II study of neratinib with or without temsirinolimus in patients with advanced HER2-mutant NSCLC reported an ORR of 19 % with combination therapy and no responses were seen in patient who received neratinib alone [62].

The activity of non-selective HER2-TKIs has yielded disappointing results although some responses have been reported in certain HER2 mutant variants which may be explained by differences in drug binding affinity. For instance, 3D modelling of the YVMA insertion demonstrated the presence of two bulky side chains leading to steric hindrance, thus affecting afatinib binding to the binding pocket [59]. Dacomitinib sensitive HER2 mutation variants contained a glycine in a position two residue beyond the end of the C helix instead of Asp770 found in wild-

Table 1
Pivotal trials of HER2 targeted therapies in *HER2*-mutant NSCLC.

Study	Patient population	N	Agents	ORR (%)	PFS	G3 + TRAE	Dose reduction, %	Dose discontinuation, %	Reference
Chemotherapy									
ECOG 2598	Untreated <i>HER2</i> 1–3+	56	Carboplatin, paclitaxel, trastuzumab	24.50 %	3.25 months	35.8 % G4	NR	NR	[49]
MSKCC	untreated <i>HER2</i> 0–3+	64	Trastuzumab with docetaxel or paclitaxel	28 %	2.4 months for Docetaxel + T, 3.9 months for Paclitaxel + T	G3 diarrhea (10 %, docetaxel)	NR	NR	[47]
U Gatzemeier et al	Untreated <i>HER2</i> 1–3+	101	Cisplatin, gemcitabine +/- trastuzumab	36 % (vs 41 % in control)	6.3 months (vs 7.2 in control)	NR	NR	NR	[48]
Immunotherapy									
Negrao et al, MDACC, Retrospective	<i>HER2</i> mutation (codons 775 and 770–785)	15	ICI monotherapy (pembrolizumab, atezolizumab, nivolumab, or durvalumab)	8 %	1.88 months	NR	NR	NR	[44]
Chen et al, Retrospective	<i>HER2</i> mutation (Exon20 insertion)	6	ICI monotherapy (drug not specified)	0 %	NA	NR	NR	NR	[51]
Lai et al, Retrospective	<i>HER2</i> mutation	26	ICI monotherapy (drug not specified)	12 %	1.9 months	NR	NR	NR	[52]
Mazieres et al, Retrospective	<i>HER2</i> mutation (Exon20 insertion)	29	ICI monotherapy (pembrolizumab, atezolizumab, nivolumab, or durvalumab)	7.4 %	2.5 months	NR	NA	NA	[52]
Guisier et al, Retrospective	<i>HER2</i> mutation (Exon20 insertion)	23	ICI monotherapy (nivolumab, pembrolizumab, or other)	27.3 %	2.2 months	NR	NA	NA	[53]
Saalfeld et al, Retrospective	<i>HER2</i> mutation (Exons 8, 19, 20)	22	Immuno-chemotherapy combination	52 %	6 months	NR	NA	NA	[54]
Chu et al, Retrospective	<i>HER2</i> mutation	16	Immuno-chemotherapy combination	37.5 %	8.4 months	11.5 %	NR	NR	[119]
Yang 2022, Retrospective	<i>HER2</i> -altered (mutant or amplification) advanced NSCLC	46	Immuno-chemotherapy combination	28.9 %	5.2 months	NR	NR	NR	[56]
Non-selective <i>HER2</i> TKIs									
Niche et al	<i>HER2</i> mutation (Exon20 insertion)	13	Afatinib	53.80 %	15.9 weeks	<10 %	NR	NR	[58]
Victoria Lai et al	<i>HER2</i> mutation (Exon20 insertion)	27	Afatinib	13 % (3/23 evaluable patients)	NR	NR	NR	NR	[57]
MG Kris et al	<i>HER</i> mutations/ amplification	30	Dacomitinib	12 % (3/26 with <i>HER2</i> mutations), 0 % (<i>HER2</i> amplification)	3 months	NR	17 %	13 %	[61]
B. Besse et al	<i>HER</i> mutations	27	Neratinib +/- temsirolimus	0 % (neratinib alone), 21 % (3/14 in combination group)	2.9 months in neratinib alone, 4 months in combination	G3 diarrhea (8 % in neratinib alone, 14 % in combination)	NR	NR	[120]
PUMA-NER-4201	<i>HER</i> mutations	62	Neratinib +/- temsirolimus	0 % (neratinib alone), 19 % (8/43 in	3 months in neratinib alone, 4.1	G3 diarrhea (12 % in neratinib alone, 14 % in combination)	NR	NR	[67]

(continued on next page)

Table 1 (continued)

Study	Patient population	N	Agents	ORR (%)	PFS	G3 + TRAE	Dose reduction, %	Dose discontinuation, %	Reference
Summit	HER2/3 mutations	26 (lung cohort)	Neratinib	combination group) 4 % (1/26)	months in combination 5.5 months	G3 diarrhea (22 %)	NR	2.8 % DD due to diarrhea	[50]
ZENITH-20 (cohort 2)	Previously treated HER2 mutation (Exon20 insertion)	90	Pozotinib	27.8 %	5.5 months	G3 rash (48.9 %), G3 diarrhea (25.6 %), G3 stomatitis (24.4 %)	76.7 %	13.3 %	[116]
ZENITH-20 (cohort 4)	Treatment naïve HER2 mutation (Exon20 insertion)	80	Pozotinib	39 %	5.6 months	G3 rash (QD, 45 %; BID, 39 %), stomatitis (QD, 21 %; BID, 15 %), diarrhea (QD, 15 %; BID, 21 %)	75 % (QD, 79 % BID, 70 %)	88 % (QD, 89 %; BID, 85 %)	[66]
Wang et al	HER2 mutation	15	Pyrotinib	53.3 % (8/15)	6.4 months	NR	NR	NR	[121]
Zhou et al	HER2 mutation	60	Pyrotinib	30 %	6.9 months	G3 or G4 (28.3 %), G3 diarrhea (20 %)	5 % due to diarrhea	1.7 %	[67]
Song et al	HER2 mutation	78	Pyrotinib	19.2 % (exon 20, 17.7 %; non-exon 20 25 %)	5.6 months	G3 (20.5 %), G3 diarrhea (16.7 %)	2.5 % (2/78)	5.1 % (4/78)	[68]
PATHER2	HER mutations/amplification	33	Pyrotinib with apatinib	51.5 %	6.9 months	G3 diarrhea (3 %), G3 hypertension (9.1 %)	30.3 % (pyrotinib)	3 % (both), 15 % (apatinib)	[69]
Zhao et al	HER2 mutation	48	Pyrotinib with inetetamab	0 % (pyrotinib 240 mg), 36.6 % (pyrotinib 320 mg)	5.5 months	14.5 % (7/48; 1 in pyrotinib 240 mg; 6 in pyrotinib 320 mg)	NR	NR	[70]
Selective HER2 TKIs									
Beamion Lung 1	HER mutations/amplification		BI 1,810,631	46 %	NR	G3 (9.3 %), alanine aminotransferase increased (7 %)	4.7 %	4.7 %	[81]
Monoclonal antibody									
HOT1303-B Trial	HER2 IHC 3 + or IHC 2+/- DISH +, and/or presence of mutation	10	Trastuzumab	0 %	5.2 months	10 % (n = 1, pneumonitis)	NR	NR	[88]
California Cancer Consortium Screening and Phase II Trial	HER2 IHC 3 + or IHC 2+	13	Trastuzumab, Docetaxel	31 % (single agent followed by combination therapy)	4.3 months	NR	NR	NR	[87]
Zinner et al	HER2 IHC 1, 2, 3 + or > 15 ng/mL of serum shed antigen ELISA	21	Trastuzumab, Gemcitabine, Cisplatin	38 %	8.5 months	NR	NR	NR	[89]
ECOG Study 2598	HER2 IHC ≥ 1+	56	Trastuzumab, Paclitaxel, Carboplatin	24.5 %	3.3 months	35.8 %*G4	NR	NR	[49]
Gatzemeier et al	HER2 IHC ≥ 2+, HER2/CEP17 ratio ≥ 2, or serum HER2 ECD 15 ng/mL by ELISA	103	Trastuzumab, Gemcitabine, Cisplatin	36 %	6.1 months (8.5 months in IHC 3 +)	G3/G4 neutropenia 57 %, thrombocytopenia 35 %, leukopenia 33 %	NR	NR	[48]
Herbst et al	NSCLC with previous treatment with chemotherapy	43	Pertuzumab	0 %	6.1 weeks	9.3 %	NR	NR	[122]
IFCT 1703-R2D2	HER2 alteration (exon 20 mutation or insertion)	47	Pertuzumab, Trastuzumab, Docetaxel	29 %	6.8 months	64 %	47 % (Docetaxel)	None	[123]

(continued on next page)

Table 1 (continued)

Study	Patient population	N	Agents	ORR (%)	PFS	G3 + TRAE	Dose reduction, %	Dose discontinuation, %	Reference
Targeted Agent and Profiling Utilization Registry Study	HER mutations/ amplification	28	Pertuzumab, Trastuzumab	11 %	16 weeks	18 %	NR	NR	[90]
Antibody drug conjugate									
Phase II basket trial (MSKCC)	HER2 mutation	18	Ado Trastuzumab- Emtansine	44 %	5 months	6 %	None	None	[92]
Iwama et al	HER2 mutation (Exon20 insertion)	22	Ado Trastuzumab- Emtansine	38.1 %	2.8 months	22.7 %	None	14 %	[94]
Hotta et al	HER2 IHC ≥ 2+, HER2/CEP17 ratio > 2, or HER2 exon 20 mutation	15	Ado Trastuzumab- Emtansine	6.7 %	2 months	G3/G4 thrombocytopenia 40 %	NR	NR	[95]
Peters et al	HER2 IHC ≥ 2+	49	Ado Trastuzumab- Emtansine	0 % (IHC 2 +), 20 % (IHC 3 +)	2.6 months (IHC 2 +), 2.7 months (IHC 3 +)	22 %	NR	4 %	[96]
Tsurutani et al, Phase I	HER2 IHC ≥ 1 + or HER2 mutation	18	Trastuzumab deruxtecan	55.6 %	11.3 months	11.1 %*serious TRAE	23.7 % (all tumors)	8.5 % (all tumors)	[99]
DESTINY-Lung01	HER2 IHC 2, 3+	49	Trastuzumab deruxtecan	24.5 %	5.4 months	73.5 %	34.7 %	22.4 %	[100]
DESTINY-Lung01	HER2 mutation	91	Trastuzumab deruxtecan	55 %	8.2 months	46 %	34 %	25 %	[101]
DESTINY-Lung02	HER2 mutation	80	Trastuzumab deruxtecan at 5.4 mg/kg and 6.4 mg/kg	53.8 % (5.4 mg/kg), 42.9 % (6.4 mg/kg)	NR	31.7 % (5.4 mg/kg),			
58 % (6.4 mg/kg)	9.9 % (5.4 mg/kg), 26 % (6.4 mg/kg)	7.9 % (5.4 mg/kg), 16 % (6.4 mg/kg)	[104]						

Abbreviations: BID, twice daily; DD, dose delay; DISH, dual color in situ hybridization; ELISA, enzyme-linked immunosorbent assay; G3 + TRAE, >3 grade 3, treatment-related adverse events; HER2 ECD, HER2 extracellular domain; ICI, immune checkpoint inhibitor; IHC, immunohistochemistry; N, number; NR, not reported; NSCLC, non-small cell lung cancer; ORR, overall response rate; PFS, progression-free survival; QD, once daily; T, Trastuzumab.

type EGFR. The presence of glycine reduces steric hindrance which would have occurred in patients with wild-type Asp 770 and acquired *HER2* exon 20 insertions, facilitates C-helix conformational changes, allowing binding of drug [63]. In view of these challenges, other novel pan-*HER2* TKIs have been developed.

Pozitotinib

Pozitotinib is an irreversible pan-*HER* TKI. The phase II ZENITH-20 study reported an ORR of 27.8 % and median PFS of 5.5 months in patients with previously treated NSCLC with *HER2* exon 20 insertion mutations [64]. Grade 3 or higher treatment-related adverse events (TRAEs) included rash (48.9 %), diarrhea (25.6 %), and stomatitis (24.4 %) and dose reduction in 77 % of patients. FDA approval was not granted based as the benefits of pozitotinib did not outweigh its toxicities and the development of pozitotinib was subsequently de-prioritized [65].

The activity of pozitotinib 16 mg QD and 8 mg BID in treatment naïve *HER2*-mutant patients have been reported recently [66]. A BID dosing was studied as it could lower C_{max} and maintain C_{trough} and overall drug exposure, potentially reducing toxicity. The ORR was 39 % (QD 45 % vs BID 30 %), DCR was 73 % (QD 75 % vs BID 70 %), and the median PFS was 5.6 months (QD 5.6 vs BID 5.6). The G778_P780dupGSP

subtype was particularly sensitive to pozitotinib with an ORR of 71 %. The most frequent grade 3 TRAE was rash (43 %) with 45 % in the QD group and 39 % in the BID group. Other grade 3 TRAEs included stomatitis (overall 19 %; QD: 21 %; BID: 15 %), diarrhea (overall 18 %; QD: 15 %; BID: 21 %), and grade 5 pneumonitis (one patient in QD group), leading to treatment discontinuation in 5 % (4/80) of the treated patients.

Pyrotinib

Pyrotinib is an irreversible pan *HER* TKI. In a phase II study of pyrotinib in patients with previously treated *HER2*-mutant NSCLC, the ORR was 30 % (95 % CI, 18.8–43.2 %) [67]. The ORR for A775_G776insYVMA and M774_A775insAYVM combined was 27 % (Fig. 4). The mPFS was 6.9 months (95 % CI, 5.5–8.3), and mOS was 14.4 months (95 % CI, 12.3–21.3) in the overall population. The most common TRAE was grade 3diarrhea (20 %). In treatment naïve patients, the ORR and PFS was 19.2 % and 5.6 months, respectively [68]. The ORR was 50 % and 100 % for A755_G776insYVMA and G776_V777.VCV variants, respectively. A phase III trial, Pyramid-1 evaluating pyrotinib vs docetaxel in patients after failure with platinum based chemotherapy in NSCLC harboring *HER2* exon 20 mutation is underway

(NCT04447118).

Pyrotinib has been combined with apatinib and inेतetamab [69,70]. PATHER2, a phase 2 study of pyrotinib in combination with apatinib, a TKI vascular endothelial growth factor receptor-2 (VEGFR2) TKI reported an ORR of 51.5 % (95 % CI, 33.5–69.2 %), and mPFS (6.9 months) and mOS (14.8 months) [69]. Inेतetamab, a recombinant humanized anti-HER2 mAb, was evaluated in *HER2*-mutant NSCLC in a phase IB study [70]. There were no DLT and adverse events were mainly grade 1–2 diarrhea, rash, and vomiting. Preliminary efficacy reported an ORR and median PFS of 5.5 months.

Tucatinib

Tucatinib is a reversible pan-HER TKI that was recently granted FDA approval in combination with trastuzumab for the treatment of *HER2* positive metastatic colorectal cancer [71]. In NSCLC, the combination's efficacy is currently being evaluated in a phase II basket trial (NCT04579380).

Furmonertinib

Furmonertinib (AST5902) is an irreversible, highly brain-penetrant pan-EGFR inhibitor with demonstrated activity against *HER2* exon 20 insertion mutations [72]. The efficacy and safety of furmonertinib is currently being evaluated in a phase I clinical trial (NCT05364073).

Mobocertinib

Mobocertinib (TAK-788), an irreversible TKI targeting *EGFR* and *HER2* exon 20 insertion mutations, was granted FDA approval in 2021 for the second line treatment of *EGFR* exon 20 insertion mutant NSCLC [73]. Preclinical data in *HER2*-mutant NSCLC showed promising results in vitro and in mouse models, with varying activity against specific variants [74]. Currently, there are no published clinical studies investigating the effects of mobocertinib on NSCLC harboring *HER2* exon 20 insertion [75].

BAY2927088

BAY2927088 is a reversible, highly potent TKI targeting *EGFR* and *HER2* driver mutations [76]. Preclinically, it showed significant dose-dependent inhibition of in vivo tumor growth in NSCLC *EGFR* ex20ins and C797S mutations. Phase 1 testing, which includes *HER2*-mutant lung cancer patients, is implemented (NCT05099172).

Tarloxotinib

Tarloxotinib is a hypoxia-activated prodrug of a pan-HER kinase inhibitor. First analysis of the phase II RAIN-701 study showed an ORR of 22 % in the *HER2* activating mutation group [77]. However, its development was terminated mid-2022 and it is no longer under clinical investigation.

Tuxobertinib (BDTX-189)

Tuxobertinib (BDTX-189) is an irreversible selective inhibitor of allosteric *EGFR* and *HER2* oncogenic mutations, including *EGFR/HER2* exon 20 insertion mutation [78]. The MasterKey-01 study showed promising anti-cancer activity in a heavily pre-treated patient population, including patients who had received prior *EGFR/HER2* TKI's with the added benefit of novel new generation TKIs being designed to inhibit *HER* mutations while sparing *HER* wildtypes with aims to reduce off target toxicities [79].

3.3.2. Selective *HER2* inhibitors

Novel *HER2* TKIs that are highly *HER2* selective with *EGFR* sparing

activity currently being developed include TAS0728, BI 1810631, JBJ-08-176-01, JIN-A04, ELVN-002, NVL-330 and ENT-H1 (Fig. 2,3).

TAS0728

TAS0728 is a *HER2*-selective inhibitor that specifically targets *HER2*. It has demonstrated potent inhibitory activity against overexpressed/amplified *HER2* and mutated *HER2* in cancer cells [80]. Assessment of TAS0728 in ongoing clinical trials is awaited (NCT03410927).

BI 1810361

BI 1,810,361 (zongertinib) is an orally active, potent, covalent, and highly selective *HER2* inhibitor [111,112]. Encouraging anti-tumor activity was observed in pantumor harboring *HER2* aberrations in the ongoing Phase I Beamion Lung 1 (NCT04886804) [81]. In *HER2*-mutant NSCLC refractory to platinum-based chemotherapy, the ORR was 46 %. Dose taken into dose optimization for BI 1810631 are 240 mg and 120 mg. Diarrhea was the most common adverse event (32.6 %) that were grade 1 or 2.

BJJ-08-176-01

BJJ-08-176-01 is a *HER2* selective, covalent TKI, that targets multiple *HER2* activating mutations including exon 20 insertion as well as amplification. JBJ-08-178-01 displayed strong selectivity toward *HER2* mutations while sparing wild type *EGFR*, compared to other *EGFR/HER2* TKIs [82]. JBJ-08-178-01 covalently binds to the *HER2* kinase domain, suggesting that the selectivity for *HER2* stems from a hydrogen bond between the triazolopyridine ring of JBJ-08-178-01 and Ser783. The efficacy of JBJ-08-178-01 was seen in various *HER2* mutant cell lines including patient-derived cells, and in vivo.

JIN-A04

JIN-A04 is a TKI selectively targeting *HER2* exon 20 insertion mutations, including *HER2* VVMA and *HER2* P780_780_Y781insGSP mutations [83]. JIN-A04 significantly inhibited Ba/F3 cell lines harboring these mutations with an IC₅₀ of 11.1 and 1.4 nM for *HER2* YVMA and GSP, respectively.

ELVN-002

ELVN-002 is a CNS-penetrant, irreversible potent *HER2* inhibitor against key *HER2* mutations, including exon 20 insertion mutations in NSCLC whilst sparing *EGFR* [84]. A phase I study evaluating ELVN-002 in patients with solid cancers harboring an abnormal *HER2* gene is ongoing (NCT05650879).

NVL-330

NVL-330 is a highly selective, CNS-penetrant, covalent *HER2* inhibitor with promising in vivo activity targeting crucial *HER2* mutations, particularly exon 20 insertions in NSCLC. Evaluation in clinical setting is awaited [85].

ENT-H1

ENT-H1 is an irreversible *HER2* inhibitor that targets both wild-type and mutated *HER2*, including exon 20 insertion mutation. In vitro analysis show potency in various cell lines, including A775G776insYVMA and G776VC [86].

3.4. *HER2*-targeted monoclonal antibodies

Early attempts to target NSCLC with *HER2* amplification have been

unsuccessful with modest responses (Table 1). Previously, HER2 overexpression (IHC 2+ or 3+) identified in ~30% of cases of NSCLC was associated with worse prognosis [87]. Initially, trastuzumab, a humanized monoclonal antibody (mAb) directly targeting HER2, was administered as monotherapy in the H01303-B Trial [88]. The ORR was 0% and median PFS was 5.2 months. Subsequently, various phase II studies included HER2-overexpressed NSCLC, and chemotherapy combination such as gemcitabine and cisplatin [48,89], paclitaxel and carboplatin [49], or docetaxel [87] to overcome limitations to trastuzumab monotherapy. Although a more favorable ORR (24.5–38%) was noted with the addition of chemotherapy, the PFS benefit was modest with a median of 3.3–8.5 months.

Pertuzumab, a HER2 humanized mAb, combined with trastuzumab in HER2 mutation and amplification resulted in a modest ORR of 11% in heavily pretreated patients [90]. Responses were reported in HER2 exon 20 mutation variants Y772_A775dup, A775_G776insYVMA and in HER2 amplification.

In a phase II study (IFCT-1703) of patients with HER2-mutated, advanced NSCLC after ≥ 1 platinum-based treatment treated with pertuzumab, trastuzumab and docetaxel, the ORR and PFS was 29% and 6.8 months, respectively [21].

3.5. HER2-targeted antibody-drug conjugates (ADCs)

An exciting emerging therapeutic class of agents are ADCs which is composed of a monoclonal antibody, payload and chemical linker designed to selectively target cancer cells [91]. By increasing efficacy via cleaving, internalizing, and releasing payload to HER2-positive cancer cells to induce cancer cell death, ADCs are able to maximize clinical efficacy and minimize unwanted toxicities, thereby overcoming the disadvantages of conventional chemotherapeutic agents.

3.5.1. T-DM1

T-DM1 (ado-trastuzumab emtastine), is a HER2-targeted ADCs linking trastuzumab with emtastine, an antimicrotubule agent with a drug-to-antibody ratio (DAR) of 3 to 4 [92,93]. In a phase II basket trial of T-DM1 in heavily pre-treated patients with HER2-mutant cancers, the ORR and PFS in patients with NSCLC was 44% and 5 months, respectively. TRAEs were mainly grade 1 or grade 2 infusion reactions, thrombocytopenia and elevated AST and ALT. Biomarker analysis showed that responders were those with HER2 mutations with differences in ORR according to mutation subtypes: 55% in HER2 exon-20 insertion mutation, 29% in other subtypes, and no responses in exon-19 mutation. Although concurrent HER2 amplification was seen in 11% of HER2-mutant NSCLC, no association was seen between HER2 IHC and clinical response. This landmark study highlighted the importance of molecular selection by HER2 mutation and the clinical utility of NGS.

A subsequent phase II trial of T-DM1 included patients with NSCLC harbouring HER2 exon20 insertion mutations with A775_G776insYVMA the most frequently identified mutation (86.4%) [94]. The ORR and PFS was 38.1%, and 2.8 months, respectively. In contrast to the efficacy observed in HER2-overexpressed breast or gastric cancer, studies of T-DM1 in metastatic NSCLC with HER2 overexpression have showed modest efficacy with an ORR of 7–20% (Table 1) [95,96].

3.5.2. Trastuzumab deruxtecan

Trastuzumab deruxtecan (T-DXd; DS-8201a) is an ADC consisting of humanized anti-HER2 antibody, peptide-based cleavable linker and topoisomerase I inhibitor payload with a DAR of approximately 8 [97]. The membrane-permeable cytotoxic payload enables efficient delivery of payload to HER2-expressing tumor cells as well as cytotoxic bystander effect on nearby tumor cells [98].

In a dose-expansion, phase 1 study of trastuzumab deruxtecan in advanced HER2-altered NSCLC, the ORR and median PFS was 55.8% and 11.3 months, respectively [99]. The ORR in the HER2-mutant subset was 72.7% and the median PFS was 11.3 months. Four patients

experienced interstitial lung disease (ILD) and pneumonitis, and 3 cases were adjudicated as related to trastuzumab deruxtecan.

In the phase II study of trastuzumab deruxtecan (DESTINY-Lung01) in patients with metastatic NSCLC with HER2 overexpression or HER2-activating mutation, the ORR was 24.5% and 20.0% for IHC 2+ and 3+ cohorts, respectively [100]. In patients with HER2-mutant NSCLC, the ORR was 55%. With HER2 mutation variants A776insYVMA and Tyr772_Ala775dup/p.Ala775_Gl associated with an ORR 61% (Fig. 4). The median PFS and median OS were 8.2 months and 17.8 months, respectively [101]. Preclinical studies have shown activating HER2 mutations enhance receptor internalization and intracellular uptake of trastuzumab deruxtecan [102], and the DESTINY-Lung01 study affirms this finding with trastuzumab deruxtecan more active in HER2-mutant NSCLC than HER2-overexpressing NSCLC. TEAE of grade 3 or more related to trastuzumab deruxtecan were reported in 55.1% of patients, including neutropenia (20.4%) and adjudicated ILD (6.6%). Dose interruption, reduction and treatment discontinuation were reported in 53.1%, 34.7%, and 22.4% of patients, respectively. Increasing awareness on trastuzumab deruxtecan related ILD/pneumonitis is crucial for better patient outcomes. [103].

Subsequently, a non-comparative, randomized phase II trial (DESTINY-Lung02) of trastuzumab deruxtecan 5.4 or 6.4 mg/kg every 3 weeks was conducted to assess the benefit-risk profile for HER2-mutant NSCLC [104]. Interim analysis reported an ORR of 53.8% and 42.9% for 5.4 mg and 6.4 mg/kg, respectively. However, safety profile was more tolerable in the 5.4 mg/kg cohort with adjudicated ILD rates of 5.9%, versus 14% with the 6.4 mg/kg. Trastuzumab deruxtecan is the first FDA approved HER2 ADC for treatment refractory HER2-mutant NSCLC [105]. Companion diagnostic test include Guardant360 CDx (Guardant Health, Redwood, CA) and Oncomine Dx Target Test (Life Technologies Corporation, MD) for plasma-based and tissue-based assay, respectively.

4. Mechanisms of resistance

Similar to other molecular targeted therapies, acquired resistance to HER2-targeted therapy is inevitable. While there are resistance mechanisms to HER2-targeted therapy in gastroesophageal adenocarcinoma [106] and breast cancer [107], there is currently insufficient data to describe the landscape of acquired resistance to HER2-targeted therapy in lung cancer. Theoretically, acquired resistance may evolve both HER2-dependent and HER2-independent pathways [108]. More specifically, HER2 secondary mutations may occur, or bypass activation such as RAS/MAPK signaling pathway or PI3K/AKT pathways may evolve. Evidence on the acquired resistance to ADCs in HER2-mutant NSCLC patients is scarce and further investigations are needed.

5. Future perspectives and conclusion

The optimal sequencing strategy and exploration of the best combination treatment still awaits to be clarified for HER2-altered NSCLC patients. Currently, only NSCLC patients with HER2 mutations are indicated for the use of HER2 directed targets as overexpression and amplifications of HER2 do not have the same benefits [30].

Identifying HER2 mutations is pivotal for NSCLC patients to benefit from these targeted therapies. Platinum based chemotherapy currently remains the preferred first line treatment for patients with HER2-mutant lung cancer. Trastuzumab deruxtecan remains the next line treatment option for patients with previously treated, advanced NSCLC with HER2 mutation on the basis of DESTINY-Lung01 and -Lung02 studies [101,104]. Upfront use of trastuzumab deruxtecan will be evaluated in the currently ongoing DESTINY-Lung 04 study in the first-line treatment of advanced NSCLC with HER2 Exon 19 or 20 mutations (NCT05048797) [30]. The intracranial efficacy for the HER2 ADCs is also not fully established. In DESTINY-Lung04, CNS PFS is being explored as one of the secondary outcome measures [30].

Combinations of HER2 directed antibody-drug conjugates and ICIs

are a promising approach, and a few phase I trials are currently underway to investigate their safety and efficacy in previously treated *HER2*-mutant advanced NSCLC patients. DESTINY-Lung03 is an ongoing phase Ib study evaluating the combination of trastuzumab deruxtecan and durvalumab with chemotherapy as a 1st line treatment in NSCLC patients with *HER2* overexpression (NCT04686305) [109]. Trastuzumab deruxtecan plus durvalumab are given as a second-line treatment (HUDSON study phase II basket trial), and trastuzumab deruxtecan in combination with pembrolizumab for *HER2*-expressing and *HER2*-mutant lung cancer (NCT04042701) are awaiting results [110].

The addition of immunotherapy to *HER2* TKIs is also being explored with *HER2*-mutated NSCLC after failure of 1st line chemotherapy in a phase II trial using pyrotinib and PD-1 inhibitors (NCT04144569). The main concern for the combination strategies would be potential overlapping toxicities such as ILD.

Other novel targets such as bi-specific antibodies have now been explored to provide opportunities to enable dual targeting of different epitopes or antigens. Zenocutuzumab (MCLA-128) is a humanized IgG1bAb that targets the extracellular domains of *HER2* and *HER3* [111]. It also has enhanced antibody-dependent cellular cytotoxicity (ADCC) activity.

Newer concepts in development include dual-payload ADCs which utilize two separate payloads, with different mechanisms of action with aims of synergistic payloads delivery for a more potent cytotoxic response [112]. Several other *HER2* ADCs such as A166, ARX788, SHR-A1811, and MRG002 are being evaluated in NSCLC with *HER2* alteration in Phase I/II clinical trials (NCT03602079, NCT03255070, NCT04818333, NCT05141786) [113].

Many exciting opportunities have emerged in the arena of *HER2*-mutant NSCLC since the initial FDA approval of the first *HER2* ADC. Ongoing clinical trials may provide further streamlining of the sequencing, viable combinations and understanding of resistance mechanisms of these novel agents.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

MID (Medical Illustration & Design), a part of the Medical Research Support Services of Yonsei University College of Medicine, for providing excellent support with medical illustrations, and Sujin Choi, who supported us with figure illustration.

References

- [1] E.K. Kim, K.A. Kim, C.Y. Lee, H.S. Shim, The frequency and clinical impact of *HER2* alterations in lung adenocarcinoma, *PLoS One* 12 (2) (2017) e0171280.
- [2] P. Stephens, C. Hunter, G. Bignell, S. Edkins, H. Davies, J. Teague, C. Stevens, S. O'Meara, R. Smith, A. Parker, A. Barthorpe, M. Blow, L. Brackenbury, A. Butler, O. Clarke, J. Cole, E. Dicks, A. Dike, A. Drozd, K. Edwards, S. Forbes, R. Foster, K. Gray, C. Greenman, K. Halliday, K. Hills, V. Kosmidou, R. Lugg, A. Menzies, J. Perry, R. Petty, K. Raine, L. Ratford, R. Shepherd, A. Small, Y. Stephens, C. Tofts, J. Varian, S. West, S. Widaa, A. Yates, F. Brasseur, C. Cooper, A.M. Flanagan, M. Knowles, S.Y. Leung, D.N. Louis, L.H. Looijenga, B. Malkowicz, M.A. Pierotti, B. Teh, G. Chenevix-Trench, B.L. Weber, S.T. Yuen, G. Harris, P. Goldstraw, A.G. Nicholson, P.A. Futreal, R. Wooster, M.R. Stratton, Lung cancer: intragenic *ERBB2* kinase mutations in tumours, *Nature* 431 (7008) (2004) 525–526, <https://doi.org/10.1038/431525b>.
- [3] D.M. Hyman, S.A. Piha-Paul, H. Won, J. Rodon, C. Saura, G.I. Shapiro, D. Juric, D.I. Quinn, V. Moreno, B. Doger, I.A. Mayer, V. Boni, E. Calvo, S. Loi, A. C. Lockhart, J.P. Erinjeri, M. Scaltriti, G.A. Ulaner, J. Patel, J. Tang, H. Beer, S. D. Selcuklu, A.J. Hanrahan, N. Bouvier, M. Melcer, R. Murali, A.M. Schram, L. M. Smyth, K. Jhaveri, B.T. Li, A. Drilon, J.J. Harding, G. Iyer, B.S. Taylor, M. F. Berger, R.E. Cutler Jr., F. Xu, A. Butturini, L.D. Eli, G. Mann, C. Farrell, A. S. Lalani, R.P. Bryce, C.L. Arteaga, F. Meric-Bernstam, J. Baselga, D.B. Solit, *HER* kinase inhibition in patients with *HER2*- and *HER3*-mutant cancers, *Nature* 554 (7691) (2018) 189–194, <https://doi.org/10.1038/nature25475>.
- [4] C.I. Bargmann, M.-C. Hung, R.A. Weinberg, The neu oncogene encodes an epidermal growth factor receptor-related protein, *Nature* 319 (6050) (1986) 226–230, <https://doi.org/10.1038/319226a0>.
- [5] M.E. Arcila, J.E. Chaff, K. Nafa, S. Roy-Chowdhuri, C. Lau, M. Zaidinski, P. K. Paik, M.F. Zakowski, M.G. Kris, M. Ladanyi, Prevalence, clinicopathologic associations, and molecular spectrum of *ERBB2* (*HER2*) tyrosine kinase mutations in lung adenocarcinomas, *Clin Cancer Res* 18 (18) (2012) 4910–4918, <https://doi.org/10.1158/1078-0432.Ccr-12-0912>.
- [6] M. Riudavets, I. Sullivan, P. Abdayem, D. Planchard, Targeting *HER2* in non-small-cell lung cancer (NSCLC): a glimpse of hope? An updated review on therapeutic strategies in NSCLC harbouring *HER2* alterations, *ESMO Open* 6 (5) (2021), 100260, <https://doi.org/10.1016/j.esmoop.2021.100260>.
- [7] I.A. Vathiotis, A. Charpidou, N. Gavrielatou, K.N. Syrigos, *HER2* Aberrations in Non-Small Cell Lung Cancer: From Pathophysiology to Targeted Therapy, *Pharmaceuticals* (Basel) 14 (12) (2021), <https://doi.org/10.3390/ph14121300>.
- [8] K.M. Ferguson, Structure-based view of epidermal growth factor receptor regulation, *Annu Rev Biophys* 37 (2008) 353–373, <https://doi.org/10.1146/annurev.biophys.37.032807.125829>.
- [9] Y. Yarden, M.X. Sliwkowski, Untangling the ErbB signalling network, *Nat Rev Mol Cell Biol* 2 (2) (2001) 127–137, <https://doi.org/10.1038/35052073>.
- [10] C. Pellegrini, M. Falleni, A. Marchetti, B. Cassani, M. Miozzo, F. Buttitta, M. Roncalli, G. Coggi, S. Bosari, *HER-2/Neu* alterations in non-small cell lung cancer: a comprehensive evaluation by real time reverse transcription-PCR, fluorescence in situ hybridization, and immunohistochemistry, *Clin Cancer Res* 9 (10 Pt 1) (2003) 3645–3652.
- [11] A. Yoshizawa, S. Sumiyoshi, M. Sonobe, M. Kobayashi, T. Uehara, M. Fujimoto, T. Tsuruyama, H. Date, H. Haga, *HER2* status in lung adenocarcinoma: a comparison of immunohistochemistry, fluorescence in situ hybridization (FISH), dual-ISH, and gene mutations, *Lung Cancer* 85 (3) (2014) 373–378, <https://doi.org/10.1016/j.lungcan.2014.06.007>.
- [12] H. Nakamura, N. Kawasaki, M. Taguchi, K. Kabasawa, Association of *HER-2* overexpression with prognosis in non-small cell lung carcinoma: a metaanalysis, *Cancer* 103 (9) (2005) 1865–1873, <https://doi.org/10.1002/cncr.20957>.
- [13] F.R. Hirsch, M. Varella-Garcia, W.A. Franklin, R. Veve, L. Chen, B. Helfrich, C. Zeng, A. Baron, P.A. Bunn Jr., Evaluation of *HER-2/neu* gene amplification and protein expression in non-small cell lung carcinomas, *Br J Cancer* 86 (9) (2002) 1449–1456, <https://doi.org/10.1038/sj.bjc.6600286>.
- [14] B.T. Li, D.S. Ross, D.L. Aisner, J.E. Chaff, M. Hsu, S.L. Kako, M.G. Kris, M. Varella-Garcia, M.E. Arcila, *HER2* Amplification and *HER2* Mutation Are Distinct Molecular Targets in Lung Cancers, *Journal of thoracic oncology : official publication of the International Association for the Study of Lung Cancer* 11 (3) (2016) 414–419, <https://doi.org/10.1016/j.jtho.2015.10.025>.
- [15] G.S. Herter-Sprie, H. Greulich, K.K. Wong, Activating Mutations in *ERBB2* and Their Impact on Diagnostics and Treatment, *Front Oncol* 3 (2013) 86, <https://doi.org/10.3389/fonc.2013.00086>.
- [16] M.A. Olaiyoye, Intracellular signaling pathways of *ErbB2/HER-2* and family members, *Breast Cancer Res* 3 (6) (2001) 385, <https://doi.org/10.1186/bcr327>.
- [17] P.A. Bunn Jr., B. Helfrich, A.F. Soriano, W.A. Franklin, M. Varella-Garcia, F. R. Hirsch, A. Baron, C. Zeng, D.C. Chan, Expression of *Her-2/neu* in human lung cancer cell lines by immunohistochemistry and fluorescence in situ hybridization and its relationship to in vitro cytotoxicity by trastuzumab and chemotherapeutic agents, *Clin Cancer Res* 7 (10) (2001) 3239–3250.
- [18] M. Jebbink, A.J. de Langen, M.C. Boelens, K. Monkhorst, E.F. Smit, The force of *HER2* - A druggable target in NSCLC? *Cancer Treat Rev* 86 (2020), 101996, <https://doi.org/10.1016/j.ctrv.2020.101996>.
- [19] Z. Calikusu, Y. Yildirim, Z. Akcali, H. Sakalli, N. Bal, I. Unal, O. Ozyilkcan, The effect of *HER2* expression on cisplatin-based chemotherapy in advanced non-small cell lung cancer patients, *J Exp Clin Cancer Res* 28 (1) (2009) 97, <https://doi.org/10.1186/1756-9966-28-97>.
- [20] F. Cappuzzo, C. Ligorio, L. Toschi, E. Rossi, R. Trisolini, D. Paioli, E. Magrini, G. Finocchiaro, S. Bartolini, A. Cancellieri, F.R. Hirsch, L. Crino, M. Varella-Garcia, *EGFR* and *HER2* gene copy number and response to first-line chemotherapy in patients with advanced non-small cell lung cancer (NSCLC), *J Thorac Oncol* 2 (5) (2007) 423–429, <https://doi.org/10.1097/JTO.0000268676.79872.9b>.
- [21] S. Kuyama, K. Hotta, M. Tabata, Y. Segawa, Y. Fujiwara, N. Takigawa, K. Kiura, H. Ueoka, K. Eguchi, M. Tanimoto, Impact of *HER2* gene and protein status on the treatment outcome of cisplatin-based chemoradiotherapy for locally advanced non-small cell lung cancer, *J Thorac Oncol* 3 (5) (2008) 477–482, <https://doi.org/10.1097/JTO.0b013e31816e2ea3>.
- [22] F.R. Hirsch, M. Varella-Garcia, P.A. Bunn Jr., M.V. Di Maria, R. Veve, R. M. Bremmes, A.E. Barón, C. Zeng, W.A. Franklin, Epidermal growth factor receptor in non-small-cell lung carcinomas: correlation between gene copy number and protein expression and impact on prognosis, *J Clin Oncol* 21 (20) (2003) 3798–3807, <https://doi.org/10.1200/jco.2003.11.069>.
- [23] J. Zhao, Y. Xia, Targeting *HER2* Alterations in Non-Small-Cell Lung Cancer: A Comprehensive Review, *JCO Precis Oncol* 4 (2020) 411–425, <https://doi.org/10.1200/po.19.00333>.
- [24] J. Zeng, W. Ma, R.B. Young, T. Li, Targeting *HER2* genomic alterations in non-small cell lung cancer, *J. Natl. Cancer Center* 1 (2) (2021) 58–73, <https://doi.org/10.1016/j.jncc.2021.04.001>.
- [25] L. Liu, X. Shao, W. Gao, J. Bai, R. Wang, P. Huang, Y. Yin, P. Liu, Y. Shu, The role of human epidermal growth factor receptor 2 as a prognostic factor in lung cancer: a meta-analysis of published data, *J Thorac Oncol* 5 (12) (2010) 1922–1932, <https://doi.org/10.1097/jto.0b013e3181f26266>.

- [26] F. Cappuzzo, M. Varella-Garcia, H. Shigematsu, I. Domenichini, S. Bartolini, G. L. Ceresoli, E. Rossi, V. Ludovini, V. Gregorc, L. Toschi, W.A. Franklin, L. Crino, A. F. Gazdar, P.A. Bunn Jr., F.R. Hirsch, Increased HER2 gene copy number is associated with response to gefitinib therapy in epidermal growth factor receptor-positive non-small-cell lung cancer patients, *J Clin Oncol* 23 (22) (2005) 5007–5018, <https://doi.org/10.1200/jco.2005.09.111>.
- [27] K. Hotta, H. Yanai, K. Ohashi, K. Ninomiya, H. Nakashima, H. Kayatani, M. Takata, K. Kiura, Pilot evaluation of a HER2 testing in non-small-cell lung cancer, *J Clin Pathol* 73 (6) (2020) 353–357, <https://doi.org/10.1136/jclinpath-2019-206204>.
- [28] M.F. Press, D.J. Slamon, K.J. Flom, J. Park, J.Y. Zhou, L. Bernstein, Evaluation of HER-2/neu gene amplification and overexpression: comparison of frequently used assay methods in a molecularly characterized cohort of breast cancer specimens, *J Clin Oncol* 20 (14) (2002) 3095–3105, <https://doi.org/10.1200/jco.2002.09.094>.
- [29] M.F. Press, G. Sauter, L. Bernstein, I.E. Villalobos, M. Mirlacher, J.Y. Zhou, R. Wardeh, Y.T. Li, R. Guzman, Y. Ma, J. Sullivan-Halley, A. Santiago, J.M. Park, A. Riva, D.J. Slamon, Diagnostic evaluation of HER-2 as a molecular target: an assessment of accuracy and reproducibility of laboratory testing in large, prospective, randomized clinical trials, *Clin Cancer Res* 11 (18) (2005) 6598–6607, <https://doi.org/10.1158/1078-0432.Ccr-05-0636>.
- [30] S. Ren, J. Wang, J. Ying, T. Mitsudomi, D.H. Lee, Z. Wang, Q. Chu, P.C. Mack, Y. Cheng, J. Duan, Y. Fan, B. Han, Z. Hui, A. Liu, J. Liu, Y. Lu, Z. Ma, M. Shi, Y. Shu, Q. Song, X. Song, Y. Song, C. Wang, X. Wang, Z. Wang, Y. Xu, Y. Yao, L. Zhang, M. Zhao, B. Zhu, J. Zhang, C. Zhou, F.R. Hirsch, Consensus for HER2 alterations testing in non-small-cell lung cancer, *ESMO Open* 7 (1) (2022), 100395, <https://doi.org/10.1016/j.esmoop.2022.100395>.
- [31] S. Zhang, W. Wang, C. Xu, Y. Zhang, X. Cai, Q. Wang, Z. Song, Z. Li, J. Yu, W. Zhong, Z. Wang, J. Liu, A. Liu, W. Li, P. Zhan, H. Liu, T. Lv, L. Miao, L. Min, G. Lin, L. Huang, J. Yuan, Z. Jiang, X. Pu, C. Rao, D. Lv, Z. Yu, X. Li, C. Tang, C. Zhou, J. Zhang, H. Guo, Q. Chu, R. Meng, X. Liu, J. Wu, J. Zhou, Z. Zhu, W. Pan, X. Dong, F. Pang, K. Wang, C. Yao, G. Lin, S. Li, Z. Yang, J. Luo, H. Jia, X. Nie, L. Wang, Y. Zhu, X. Hu, Y. Xie, X. Lin, J. Cai, Y. Xia, H. Feng, L. Wang, Y. Du, W. Yao, X. Shi, X. Niu, D. Yuan, Y. Yao, J. Kang, J. Zhang, C. Zhang, W. Gao, J. Huang, Y. Zhang, P. Sun, H. Wang, M. Ye, D. Wang, Z. Wang, B. Wan, D. Lv, G. Yu, L. Shi, Y. Xia, F. Gao, X. Zhang, T. Xu, W. Zhou, H. Wang, Z. Liu, N. Yang, L. Wu, Q. Wang, G. Wang, Z. Hong, J. Wang, M. Fang, Y. Fang, Y. Zhang, Y. Song, S. Ma, W. Fang, Y. Lu, Chinese expert consensus on the diagnosis and treatment of HER2-altered non-small cell lung cancer, *Thorac Cancer* 14 (1) (2023) 91–104, <https://doi.org/10.1111/1759-7714.14743>.
- [32] L.M. Sholl, D.L. Aisner, M. Varella-Garcia, L.D. Berry, D. Dias-Santagata, H. Wistuba II, J. Chen, K. Fujimoto, W.A. Kugler, A.J. Franklin, M. Iafra, M. G. Ladanyi, B.E. Kris, P.A. Johnson, J.D. Bunn, D.J.K. Minna, Multi-institutional Oncogenic Driver Mutation Analysis in Lung Adenocarcinoma: The Lung Cancer Mutation Consortium Experience, *J Thorac Oncol* 10 (5) (2015) 768–777, <https://doi.org/10.1097/jto.0000000000000516>.
- [33] R.N. Pillai, M. Behera, L.D. Berry, M.R. Rossi, M.G. Kris, B.E. Johnson, P.A. Bunn, S.S. Ramalingam, F.R. Khuri, HER2 mutations in lung adenocarcinomas: A report from the Lung Cancer Mutation Consortium, *Cancer* 123 (21) (2017) 4099–4105, <https://doi.org/10.1002/cncr.30869>.
- [34] S. Peters, S. Zimmermann, Targeted therapy in NSCLC driven by HER2 insertions, *Transl Lung Cancer Res* 3 (2) (2014) 84–88, <https://doi.org/10.3978/j.issn.2218-6751.2014.02.06>.
- [35] J.P. Robichaux, Y.Y. Elamin, R.S.K. Vijayan, M.B. Nilsson, L. Hu, J. He, F. Zhang, M. Pisegna, A. Poteete, H. Sun, S. Li, T. Chen, H. Han, M.V. Negrao, J.R. Ahnert, L. Diao, J. Wang, X. Le, F. Meric-Bernstam, M. Routbort, B. Roeck, Z. Yang, V. M. Raymond, R.B. Lanman, G.M. Frampton, V.A. Miller, A.B. Schrock, L. A. Albacker, K.K. Wong, J.B. Cross, J.V. Heymach, Pan-Cancer Landscape and Analysis of ERBB2 Mutations Identifies Pzoitinib as a Clinically Active Inhibitor and Enhancer of T-DM1 Activity, *Cancer Cell* 36 (4) (2019) 444–457.e447, <https://doi.org/10.1016/j.ccell.2019.09.001>.
- [36] A. Friedlaender, V. Subbiah, A. Russo, G.L. Banna, U. Malapelle, C. Rolfo, A. Addeo, EGFR and HER2 exon 20 insertions in solid tumours: from biology to treatment, *Nat Rev Clin Oncol* 19 (1) (2022) 51–69, <https://doi.org/10.1038/s41571-021-00558-1>.
- [37] S. Yang, Y. Wang, C. Zhao, X. Li, Q. Liu, S. Mao, Y. Liu, X. Yu, W. Wang, Q. Tian, Y. Pan, A. Xiong, B. Chen, G. Gao, W. Li, Y. He, F. Wu, X. Chen, C. Su, S. Ren, C. Zhou, Exon 20 YVMA insertion is associated with high incidence of brain metastasis and inferior outcome of chemotherapy in advanced non-small cell lung cancer patients with HER2 kinase domain mutations, *Transl Lung Cancer Res* 10 (2) (2021) 753–765, <https://doi.org/10.21037/tlcr-20-559>.
- [38] S.I. Ou, A.B. Schrock, E.V. Bocharov, S.J. Klempner, C.K. Haddad, G. Steinecker, M. Johnson, B.J. Gitlitz, J. Chung, P.V. Campregher, J.S. Ross, P.J. Stephens, V. A. Miller, J.H. Suh, S.M. Ali, V. Velcheti, HER2 Transmembrane Domain (TMD) Mutations (V659/G660) That Stabilize Homo- and Heterodimerization Are Rare Oncogenic Drivers in Lung Adenocarcinoma That Respond to Afatinib, *J Thorac Oncol* 12 (3) (2017) 446–457, <https://doi.org/10.1016/j.jtho.2016.11.2224>.
- [39] K.B. Pahuja, T.T. Nguyen, B.S. Jaiswal, K. Prabhaskar, T.M. Thaker, K. Senger, S. Chaudhuri, N.M. Kljavin, A. Antony, S. Phalke, P. Kumar, M. Mravic, E. W. Stawski, D. Vargas, S. Durinck, R. Gupta, A. Khanna-Gupta, S.E. Trabucco, E. S. Sokol, R.J. Hartmaier, A. Singh, A. Chougule, V. Trivedi, A. Dutt, V. Patil, A. Joshi, V. Noronha, J. Ziai, S.D. Banavali, V. Ramprasad, W.F. DeGrado, R. Bueno, N. Jura, S. Seshagiri, Actionable Activating Oncogenic ERBB2/HER2 Transmembrane and Juxtamembrane Domain Mutations, *Cancer Cell* 34 (5) (2018) 792–806.e795, <https://doi.org/10.1016/j.ccell.2018.09.010>.
- [40] J. Mazières, F. Barlesi, T. Filleron, B. Besse, I. Monnet, M. Beau-Faller, S. Peters, E. Dansin, M. Früh, M. Pless, R. Rosell, M. Wislez, P. Fournel, V. Westeel, F. Cappuzzo, A. Cortot, D. Moro-Sibilot, J. Milia, O. Gautschi, Lung cancer patients with HER2 mutations treated with chemotherapy and HER2-targeted drugs: results from the European EUHER2 cohort, *Ann Oncol* 27 (2) (2016) 281–286, <https://doi.org/10.1093/annonc/mdv573>.
- [41] R. Chen, J. Zhao, G. Lin, L. Liu, L. Chen, X. Hu, X. Ai, Z. Fan, C. Xu, W. Wang, W. Zhuang, M. Fang, Y. Zhu, G. Chen, Y. Guan, L. Yang, X. Xia, X. Yi, Ma, 07.13 NGS Sequencing Based Liquid / Tissue Biopsy Identified Coexistence of HER2 Amplification and Mutation in Advanced NSCLC Patients, *Journal of Thoracic Oncology*, S1830 12 (2017), <https://doi.org/10.1016/j.jtho.2017.09.513>.
- [42] S.S. Ramalingam, J.C. Yang, C.K. Lee, T. Kurata, D.W. Kim, T. John, N. Nogami, Y. Ohe, H. Mann, Y. Rukaznikov, S. Ghiorghiu, D. Stetson, A. Markovets, J. C. Barrett, K.S. Thress, P.A. Jänne, Osimertinib As First-Line Treatment of EGFR Mutation-Positive Advanced Non-Small-Cell Lung Cancer, *J Clin Oncol* 36 (9) (2018) 841–849, <https://doi.org/10.1200/jco.2017.74.7576>.
- [43] A.C. Tan, S.P.L. Saw, J. Chen, G.G.Y. Lai, H.N. Oo, A. Takano, D.P.X. Lau, J.P. S. Yeong, G.S. Tan, K.H. Lim, A.J. Skanderup, J.W.K. Chan, Y.L. Teh, T. Rajasekaran, A. Jain, W.L. Tan, Q.S. Ng, R. Kanesvaran, W.T. Lim, M.K. Ang, D. S.W. Tan, Clinical and Genomic Features of HER2 Exon 20 Insertion Mutations and Characterization of HER2 Expression by Immunohistochemistry in East Asian Non-Small-Cell Lung Cancer, *JCO Precis Oncol* 6 (2022) e2200278.
- [44] K. Chen, G. Pan, G. Cheng, F. Zhang, Y. Xu, Z. Huang, Y. Fan, Immune microenvironment features and efficacy of PD-1/PD-L1 blockade in non-small cell lung cancer patients with EGFR or HER2 exon 20 insertions, *Thorac Cancer* 12 (2) (2021) 218–226, <https://doi.org/10.1111/1759-7714.13748>.
- [45] S. Wu, Q. Zhang, F. Zhang, F. Meng, S. Liu, R. Zhou, Q. Wu, X. Li, L. Shen, J. Huang, J. Qin, S. Ouyang, Z. Xia, H. Song, X.-H. Feng, J. Zou, P. Xu, HER2 recruits AKT1 to disrupt STING signalling and suppress antiviral defence and antitumour immunity, *Nat Cell Biol* 21 (8) (2019) 1027–1040, <https://doi.org/10.1038/s41556-019-0352-z>.
- [46] Y. Wang, S. Zhang, F. Wu, J. Zhao, X. Li, C. Zhao, S. Ren, C. Zhou, Outcomes of Pemertrex-based chemotherapies in HER2-mutant lung cancers, *BMC Cancer* 18 (1) (2018) 326, <https://doi.org/10.1186/s12885-018-4277-x>.
- [47] L.M. Krug, V.A. Miller, J. Patel, J. Crapanzano, C.G. Azzoli, J. Gomez, M.G. Kris, R.T. Heelan, B. Pizzo, L. Tyson, C. Sheehan, J.S. Ross, E. Venkatraman, Randomized phase II study of weekly docetaxel plus trastuzumab versus weekly paclitaxel plus trastuzumab in patients with previously untreated advanced nonsmall cell lung carcinoma, *Cancer* 104 (10) (2005) 2149–2155, <https://doi.org/10.1002/cncr.21428>.
- [48] U. Gatzemeier, G. Groth, C. Butts, N. Van Zandwijk, F. Shepherd, A. Ardizzone, C. Barton, P. Ghahramani, V. Hirsh, Randomized phase II trial of gemcitabine-cisplatin with or without trastuzumab in HER2-positive non-small-cell lung cancer, *Ann Oncol* 15 (1) (2004) 19–27, <https://doi.org/10.1093/annonc/mdh031>.
- [49] C.J. Langer, P. Stephenson, A. Thor, M. Vangel, D.H. Johnson, Trastuzumab in the treatment of advanced non-small-cell lung cancer: is there a role? Focus on Eastern Cooperative Oncology Group study 2598, *J Clin Oncol* 22 (7) (2004) 1180–1187, <https://doi.org/10.1200/jco.2004.04.105>.
- [50] M.V. Negrao, F. Skoulidis, M. Montesin, K. Schulze, I. Bara, V. Shen, H. Xu, S. Hu, D. Sui, Y.Y. Elamin, X. Le, M.E. Goldberg, K. Murugesan, C.J. Wu, J. Zhang, D.S. Barreto, J.P. Robichaux, A. Reuben, T. Cascone, C.M. Gay, K.G. Mitchell, L. Hong, W. Rinsurongkawong, J.A. Roth, S.G. Swisher, J. Lee, A. Tsao, V. Papadimitrakopoulou, D.L. Gibbons, B.S. Glisson, G. Singal, V.A. Miller, B. Alexander, G. Frampton, L.A. Albacker, D. Shames, J. (2021), <https://doi.org/10.1136/jitc-2021-002891>. Zhang, J.V. Heymach, Oncogene-specific differences in tumor mutational burden, PD-L1 expression, and outcomes from immunotherapy in non-small cell lung cancer, *J Immunother Cancer* 9 (8).
- [51] W.-C.-V. Lai, D.L. Feldman, D.J. Buonocore, E.B. Brzostowski, H. Rizvi, A. J. Plodkowski, A. Ni, J.K. Sabari, M.D. Offin, M.G. Kris, A.E. Drilon, M.E. Arcila, M.D. Hellmann, B.T. Li, PD-L1 expression, tumor mutation burden and response to immune checkpoint blockade in patients with HER2-mutant lung cancers, *J Clin Oncol* 36 (15 suppl) (2018) 9060, https://doi.org/10.1200/JCO.2018.36.15_suppl.9060.
- [52] J. Mazieres, A. Drilon, A. Lusque, L. Mhanna, A.B. Cortot, L. Mezquita, A.A. Thai, C. Masciaux, S. Couraud, R. Veillon, M. Van den Heuvel, J. Neal, N. Peled, M. Früh, T.L. Ng, V. Gounant, S. Popat, J. Diebold, J. Sabari, V.W. Zhu, S. I. Rothschild, P. Bironzo, A. Martinez-Marti, A. Curioni-Fontecedro, R. Rosell, M. Lattuca-Truc, M. Wiesweg, B. Besse, B. Solomon, F. Barlesi, R.D. Schouten, H. Wakelee, D.R. Camidge, G. Zalcman, S. Novello, S.I. Ou, J. Milia, O. Gautschi, Immune checkpoint inhibitors for patients with advanced lung cancer and oncogenic driver alterations: results from the IMMUNOTARGET registry, *Ann Oncol* 30 (8) (2019) 1321–1328, <https://doi.org/10.1093/annonc/mdz167>.
- [53] F. Guisier, C. Dubos-Arvis, F. Vinas, H. Dobre, C. Ricordel, S. Robert, H. Janicot, M. Bernardi, P. Fournel, R. Lamy, M. Pérol, J. Dauba, G. Gonzales, L. Falchero, C. Decroisette, P. Assouline, C. Chouaid, O. Bylicki, Efficacy and Safety of Anti-PD-1 Immunotherapy in Patients With Advanced NSCLC With BRAF, HER2, or MET Mutations or RET Translocation: GFPC 01–2018, *J Thorac Oncol* 15 (4) (2020) 628–636, <https://doi.org/10.1016/j.jtho.2019.12.129>.
- [54] F.C. Saalfeld, C. Wenzel, P. Christopoulos, S. Merkelbach-Bruse, T.M. Reissig, S. Laßmann, S. Thiel, J.A. Stratmann, R. Marienfeld, J. Berger, A. Desuki, J. L. Velthaus, D. Kauffmann-Guerrero, A. Stenzinger, S. Michels, T. Herold, M. Kramer, S. Herold, A. Tufman, S. Loges, J. Alt, M. Joosten, G. Schmidtke-Schrezenmeier, M. Sebastian, S. Stephan-Falkenau, C.F. Waller, M. Wiesweg, J. Wolf, M. Thomas, D.E. Aust, M. Wermke, Efficacy of Immune Checkpoint Inhibitors Alone or in Combination With Chemotherapy in NSCLC Harboring

- ERBB2 Mutations, *J Thorac Oncol* 16 (11) (2021) 1952–1958, <https://doi.org/10.1016/j.jtho.2021.06.025>.
- [55] L. Gandhi, D. Rodríguez-Abreu, S. Gadgeel, E. Esteban, E. Felip, F. De Angelis, M. Domine, P. Clingan, M.J. Hochmair, S.F. Powell, S.Y. Cheng, H.G. Bischoff, N. Peled, F. Grossi, R.R. Jennens, M. Reck, R. Hui, E.B. Garon, M. Boyer, B. Rubio-Viqueira, S. Novello, T. Kurata, J.E. Gray, J. Vida, Z. Wei, J. Yang, H. Raftopoulos, M.C. Pietanza, M.C. Garassino, Pembrolizumab plus Chemotherapy in Metastatic Non-Small-Cell Lung Cancer, *N Engl J Med* 378 (22) (2018) 2078–2092, <https://doi.org/10.1056/NEJMoa1801005>.
 - [56] G. Yang, Y. Yang, R. Liu, W. Li, H. Xu, X. Hao, J. Li, P. Xing, S. Zhang, X. Ai, F. Xu, Y. Wang, First-line immunotherapy or angiogenesis inhibitor plus chemotherapy for HER2-altered NSCLC: a retrospective real-world POLISH study, *Ther Adv Med Oncol* 14 (2022), <https://doi.org/10.1177/17588359221082339>, 17588359221082339.
 - [57] W.V. Lai, L. Lebas, T.A. Barnes, J. Milia, A. Ni, O. Gautschi, S. Peters, R. Ferrara, A.J. Plodkowski, J. Kavanagh, J.K. Sabari, S.J. Clarke, N. Pavlakakis, A. Drilon, C. M. Rudin, M.E. Arcila, N.B. Leigh, F.A. Shepherd, M.G. Kris, J. Mazieres, B.T. Li, Afatinib in patients with metastatic or recurrent HER2-mutant lung cancers: a retrospective international multicentre study, *Eur J Cancer* 109 (2019) 28–35, <https://doi.org/10.1016/j.ejca.2018.11.030>.
 - [58] R. Dziadziuszko, E.F. Smit, U. Dafni, J. Wolf, B. Wasag, W. Biernat, S.P. Finn, R. Kamlmer, Z. Tsourti, M. Rabaglio, B. Ruepp, H. Roschitzki-Voser, R.A. Stahel, E. Felip, S. Peters, Afatinib in NSCLC With HER2 Mutations: Results of the Prospective, Open-Label Phase II NICHE Trial of European Thoracic Oncology Platform (ETOP), *J Thorac Oncol* 14 (6) (2019) 1086–1094, <https://doi.org/10.1016/j.jtho.2019.02.017>.
 - [59] W. Fang, S. Zhao, Y. Liang, Y. Yang, L. Zhang, X. Dong, L. Zhang, Y. Tang, S. Wang, Y. Yang, X. Ma, M. Wang, W. Wang, S. Zhao, K. Wang, S. Gao, L. Zhang, Mutation Variants and Co-Mutations as Genomic Modifiers of Response to Afatinib in HER2-Mutant Lung Adenocarcinoma, *Oncologist* 25 (3) (2020) e545–e554, <https://doi.org/10.1634/theoncologist.2019-0547>.
 - [60] S. Zhao, W. Fang, Y. Yang, L. Zhang, P1-01-110 Three Specific HER2 Mutations Predict Favorable Outcomes in Advanced Lung Cancer Patients Treated with Afatinib, *J Thorac Oncol* 13 (10) (2018) S506–S507, <https://doi.org/10.1016/j.jtho.2018.08.667>.
 - [61] M.G. Kris, D.R. Camidge, G. Giaccone, T. Hida, B.T. Li, J. O'Connell, I. Taylor, H. Zhang, M.E. Arcila, Z. Goldberg, P.A. Jänne, Targeting HER2 aberrations as actionable drivers in lung cancers: phase II trial of the pan-HER tyrosine kinase inhibitor dacomitinib in patients with HER2-mutant or amplified tumors, *Ann Oncol* 26 (7) (2015) 1421–1427, <https://doi.org/10.1093/annonc/mdv186>.
 - [62] L. Gandhi, B. Besse, J. Mazieres, S. Waqar, A. Cortot, F. Barlesi, E. Quoix, G. Otterson, D. Ettinger, L. Horn, D. Moro-Sibilot, M. Socinski, K. Gold, J. Gray, A. Oton, R.S. Heist, D. Costa, L. McCulloch, J. Bechuk, R. Bryce, M. Kris, MA04.02 Neratinib ± Temozolimumab in HER2-Mutant Lung Cancers: An International, Randomized Phase II Study, *J Thorac Oncol* 12 (1) (2017) S358–S359, <https://doi.org/10.1016/j.jtho.2016.11.398>.
 - [63] T. Kosaka, J. Tanizaki, R.M. Paranal, H. Endoh, C. Lydon, M. Capelletti, C. E. Repellin, J. Choi, A. Ogino, A. Calles, D. Ercan, A.J. Redig, M. Bahcall, G. R. Oxnard, M.J. Eck, P.A. Jänne, Response Heterogeneity of EGFR and HER2 Exon 20 Insertions to Covalent EGFR and HER2 Inhibitors, *Cancer Res* 77 (10) (2017) 2712–2721, <https://doi.org/10.1158/0008-5472.Can-16-3404>.
 - [64] X. Le, R. Cornelissen, M. Garassino, J.M. Clarke, N. Tchekmedyan, J. W. Goldman, S.Y. Leu, G. Bhat, F. Lebel, J.V. Heymach, M.A. Socinski, Poziotinib in Non-Small-Cell Lung Cancer Harboring HER2 Exon 20 Insertion Mutations After Prior Therapies: ZENITH20-2 Trial, *J Clin Oncol* 40 (7) (2022) 710–718, <https://doi.org/10.1200/jco.21.01323>.
 - [65] N. Tucker, FDA Denies Approval of Poziotinib for Advanced/Metastatic NSCLC With HER2 Exon 20 Insertions. Targeted Oncology.
 - [66] R. Cornelissen, A. Prelaj, S. Sun, C. Baik, M. Wollner, E.B. Haura, H. Mamdani, J. W. Riess, F. Cappuzzo, M.C. Garassino, J.V. Heymach, M.A. Socinski, S.Y. Leu, G. Bhat, F. Lebel, X. Le, Poziotinib in Treatment-Naïve NSCLC Harboring HER2 Exon 20 Mutations: ZENITH20-4, A Multicenter, Multicohort, Open-Label, Phase 2 Trial (Cohort 4), *J Thorac Oncol* (2023), <https://doi.org/10.1016/j.jtho.2023.03.016>.
 - [67] C. Zhou, X. Li, Q. Wang, G. Gao, Y. Zhang, J. Chen, Y. Shu, Y. Hu, Y. Fan, J. Fang, G. Chen, J. Zhao, J. He, F. Wu, J. Zou, X. Zhu, X. Lin, Pyrotinib in HER2-Mutant Advanced Lung Adenocarcinoma After Platinum-Based Chemotherapy: A Multicenter, Open-Label, Single-Arm, Phase II Study, *J Clin Oncol* 38 (24) (2020) 2753–2761, <https://doi.org/10.1200/jco.20.00297>.
 - [68] Z. Song, Y. Li, S. Chen, S. Ying, S. Xu, J. Huang, D. Wu, D. Lv, T. Bei, S. Liu, X. Huang, C. Xie, X. Wu, J. Fu, F. Hua, W. Wang, C. Xu, C. Gao, S. Cai, S. Lu, Y. Zhang, Efficacy and safety of pyrotinib in advanced lung adenocarcinoma with HER2 mutations: a multicenter, single-arm, phase II trial, *BMC Med* 20 (1) (2022) 42, <https://doi.org/10.1186/s12916-022-02245-z>.
 - [69] G. Yang, H. Xu, Y. Yang, S. Zhang, F. Xu, X. Hao, J. Li, P. Xing, X. Hu, Y. Liu, L. Wang, L. Lin, Z. Wang, J. Duan, J. Wang, Y. Wang, Pyrotinib combined with apatinib for targeting metastatic non-small cell lung cancer with HER2 alterations: a prospective, open-label, single-arm phase 2 study (PATHER2), *BMC Med* 20 (1) (2022) 277, <https://doi.org/10.1186/s12916-022-02470-6>.
 - [70] Y. Zhao, Y. Huang, Y. Zhang, S. Hong, Y. Huang, H. Zhao, Y. Yang, S. Zhao, G. Chen, H. Zhou, Y. Ma, N. Zhou, L. Zhang, Safety and efficacy of inetinamab in combination with pyrotinib in HER2 mutant patients with non-small cell lung cancer (NSCLC): An open-label, phase Ib trial, *J Clin Oncol* 41 (16_suppl) (2023) 9105, https://doi.org/10.1200/JCO.2023.41.16_suppl.9105.
 - [71] S.J. Casak, M.N. Horiba, M. Yuan, J. Cheng, S.J. Lemery, Y.L. Shen, W. Fu, J. N. Moore, Y. Li, Y. Bi, D. Auth, N. Fesenko, P.G. Kluetz, R. Pazdur, L.A. Fashoyin-Aje, FDA Approval Summary: Tucatinib with trastuzumab for advanced unresectable or metastatic, chemotherapy refractory, HER2 positive RAS wild type colorectal cancer, *Clin Cancer Res* (2023), <https://doi.org/10.1158/1078-0432.Ccr-23-1041>.
 - [72] L. Musib, M. Kowanetz, Q. Li, H. Luo, J. Hu, S. Lutzker, PP01.11 Furmonertinib is an Oral, Irreversible, Highly Brain-Penetrant Pan-EGFR Inhibitor with Activity Against Classical and Atypical EGFR Mutations, *J Thorac Oncol* 18 (3, Supplement) (2023) e14–e15, <https://doi.org/10.1016/j.jtho.2022.09.037>.
 - [73] C. Zhou, S.S. Ramalingam, T.M. Kim, S.W. Kim, J.C. Yang, G.J. Riely, T. Mekhail, D. Nguyen, M.R. Garcia Campelo, E. Felip, S. Vincent, S. Jin, C. Griffin, V. Bunn, J. Lin, H.M. Lin, M. Mehta, P.A. Jänne, Treatment Outcomes and Safety of Mobocertinib in Platinum-Pretreated Patients With EGFR Exon 20 Insertion-Positive Metastatic Non-Small Cell Lung Cancer: A Phase 1/2 Open-label Nonrandomized Clinical Trial, *JAMA Oncol* 7 (12) (2021) e214761.
 - [74] H. Han, S. Li, T. Chen, M. Fitzgerald, S. Liu, C. Peng, K.H. Tang, S. Cao, J. Chouitar, J. Wu, D. Peng, J. Deng, Z. Gao, T.E. Baker, F. Li, H. Zhang, Y. Pan, H. Ding, H. Hu, V. Pyon, C. Thakurdin, E. Papadopoulos, S. Tang, F. Gonzalez, H. Chen, V.M. Rivera, R. Brake, S. Vincent, K.K. Wong, Targeting HER2 Exon 20 Insertion-Mutant Lung Adenocarcinoma with a Novel Tyrosine Kinase Inhibitor Mobocertinib, *Cancer Res* 81 (20) (2021) 5311–5324, <https://doi.org/10.1158/0008-5472.Can-21-1526>.
 - [75] D. Sentana-Lledo, E. Academia, H. Viray, D. Rangachari, S.S. Kobayashi, P. A. VanderLaan, D.B. Costa, EGFR exon 20 insertion mutations and ERBB2 mutations in lung cancer: a narrative review on approved targeted therapies from oral kinase inhibitors to antibody-drug conjugates, *Transl Lung Cancer Res* 12 (7) (2023) 1590–1610, <https://doi.org/10.21037/tlcr-23-98>.
 - [76] F. Siegel, G. Karsli-Uzunbas, K. Kotynkova, Q. McVeigh, S. Siegel, D. Korr, V. Schulze, M. Berger, G. Beckmann, A. Cherniack, M. Meyerson, H. Greulich, Abstract 4035: Preclinical activity of BAY 2927088 in HER2 mutant non-small cell lung cancer, *Cancer Res* 83 (7_Supplement) (2023) 4035, <https://doi.org/10.1158/1538-7445.Am2023-4035>.
 - [77] S.V. Liu, L.C. Villaruz, V.H.F. Lee, V.W. Zhu, C.S. Baik, A. Sacher, C.E. McCoach, D. Nguyen, J.Y.C. Li, J.M. Pacheco, C. Kim, T.F. Burns, E.L. Schenk, N. Leigh, L. Tozzi, D.R. Camidge, LBA61 First analysis of RAIN-701: Study of tarloxotinib in patients with non-small cell lung cancer (NSCLC) EGFR Exon 20 insertion, HER2-activating mutations & other solid tumours with NRG1/ERBB gene fusions, *Ann Oncol* 31 (2020) S1189, <https://doi.org/10.1016/j.annonc.2020.08.2294>.
 - [78] A.M. Schram, J.R. Ahnert, M.R. Patel, S. Jauhari, J.C. Sachdev, V.W. Zhu, P. LoRusso, D. Nguyen, X. Le, M. O'Connor, N. Waters, C. Cook, K. Witt, R. W. Humphrey, P.A. Janne, E.P. Hamilton, Safety and preliminary efficacy from the phase 1 portion of MasterKey-01: A First-in-human dose-escalation study to determine the recommended phase 2 dose (RP2D), pharmacokinetics (PK) and preliminary antitumor activity of BDTX-189, an inhibitor of allosteric ErbB mutations, in patients (pts) with advanced solid malignancies, *J Clin Oncol* 39 (15_suppl) (2021) 3086, https://doi.org/10.1200/JCO.2021.39.15_suppl.3086.
 - [79] E.P. Hamilton, M.R. Patel, J. Rodon, D.S. Hong, A.M. Schram, P.A. Janne, P. LoRusso, J.C. Sachdev, S.H. Ou, E.A. Buck, M. O'Connor, N. Waters, K. Witt, C. Cook, Masterkey-01: Phase I/II, open-label multicenter study to assess safety, tolerability, pharmacokinetics, and antitumor activity of BDTX-189, an inhibitor of allosteric ErbB mutations, in patients with advanced solid malignancies, *J Clin Oncol* 38 (15_suppl) (2020), https://doi.org/10.1200/JCO.2020.38.15_suppl.TPS3665.
 - [80] H. Irie, K. Ito, Y. Fujioka, K. Oguchi, A. Fujioka, A. Hashimoto, H. Ohsawa, K. Tanaka, K. Funabashi, H. Arai, Y. Kawai, T. Shimamura, R. Wadhwani, S. Ohkubo, K. Matsuo, TAS0728, A Covalent-binding, HER2-selective Kinase Inhibitor Shows Potent Antitumor Activity in Preclinical Models, *Mol Cancer Ther* 18 (4) (2019) 733–742, <https://doi.org/10.1158/1535-7163.Mct-18-1085>.
 - [81] F. Opdam, M.A. Barve, H.-Y. Tu, Y.-L. Wu, D. Berz, N. Gibson, B. Sadrolhefazi, J. Serra, K. Yoh, N. Yamamoto, Phase I Beamion Lung 1 trial of BI 1810631, a HER2 tyrosine kinase inhibitor (TKI), as monotherapy in patients (pts) with advanced/metastatic solid tumors with HER2 aberrations: Updated data, *J Clin Oncol* 41 (16_suppl) (2023) 8545, https://doi.org/10.1200/JCO.2023.41.16_suppl.8545.
 - [82] J. Son, J. Jang, T.S. Beyett, Y. Eum, H.M. Haikala, A. Verano, M. Lin, J. M. Hatcher, N.P. Kwiatkowski, P. Eser, M.J. Poitras, S. Wang, M. Xu, P. C. Gokhale, M.D. Cameron, M.J. Eck, N.S. Gray, P.A. Jänne, A Novel HER2-Selective Kinase Inhibitor Is Effective in HER2 Mutant and Amplified Non-Small Cell Lung Cancer, *Cancer Res* 82 (8) (2022) 1633–1645, <https://doi.org/10.1158/0008-5472.Can-21-2693>.
 - [83] M.R. Yu, M.R. Yun, J.B. Lee, J.Y. Lee, S.W. Aum, S.J. Choi, J.Y. Park, S.Y. Oh, E. J. Lee, K.B. Duggirala, K. Lee, M.H. Hong, S.M. Lim, A. Jo, E. Seah, C. Kim, B. C. Cho, Abstract 4029: JIN-A04, highly effective tyrosine kinase inhibitor targeting HER2 exon 20 insertion mutations in NSCLC, *Cancer Res* 83 (7_Supplement) (2023) 4029, <https://doi.org/10.1158/1538-7445.Am2023-4029>.
 - [84] M. Aujay, A.J. Broad, S.D. Gross, L. Ren, J.P. Lyssikatos, S. Kintz, Q. Wang, H. Collins, Abstract 4019: Preclinical activity of ELVN-002: A potent, selective, and irreversible HER2 and pan-HER2 mutant small molecule inhibitor for the treatment of HER2 driven malignancies, *Cancer Res* 83 (7_Supplement) (2023) 4019, <https://doi.org/10.1158/1538-7445.Am2023-4019>.
 - [85] K.L. Andrews, Y. Sun, B. Gerard, A. Tangperachaiikul, S. Mente, R.A. Kemper, C. M. Martin, N.E. Kohl, J.C. Horan, H.E. Pelish, NVL-330 is a selective, brain-penetrant inhibitor of oncogenic HER2 exon 20 insertion mutations in preclinical models, *Eur J Cancer* 174 (2022) S76–S77, [https://doi.org/10.1016/S0959-8049\(22\)01003-6](https://doi.org/10.1016/S0959-8049(22)01003-6).
 - [86] C. Zhao, L. Kulyk, I. Botrous, K. Chen, C. Zhao, A. Banisadr, M.L. Anderson, F. Manby, T. Miller, C. Zhang, L. Gomez, Z. Huang, Abstract 4034: A potent and

- highly selective irreversible HER2 inhibitor for treating HER2-driven cancers, *Cancer Res* 83 (7, Supplement) (2023) 4034, <https://doi.org/10.1158/1538-7445.Am2023-4034>.
- [87] P.N. Lara Jr., L. Laptalo, J. Longmate, D.H. Lau, R. Gandour-Edwards, P. H. Gumerlock, J.H. Doroshow, D.R. Gandara, Trastuzumab plus docetaxel in HER2/neu-positive non-small-cell lung cancer: a California Cancer Consortium screening and phase II trial, *Clin Lung Cancer* 5 (4) (2004) 231–236, <https://doi.org/10.3816/clc.2004.n.004>.
- [88] I. Kinoshita, T. Goda, K. Watanabe, M. Maemondo, S. Oizumi, T. Amano, Y. Hatanaka, Y. Matsuno, H. Nishihara, H. Asahina, T. Harada, K. Goto, H. Isobe, M. Nishimura, H. Dosaka-Akita, A phase II study of trastuzumab monotherapy in pretreated patients with non-small cell lung cancers (NSCLCs) harboring HER2 alterations: H0T1303-B trial, *Annals of Oncology* 29 (2018) viii540, <https://doi.org/10.1093/annonc/mdy292.112>.
- [89] R.G. Zinner, B.S. Glisson, F.V. Fossella, K.M. Pisters, M.S. Kies, P.M. Lee, E. Massarelli, B. Sabloff, H.A. Fritsche Jr., J.Y. Ro, N.G. Ordóñez, H.T. Tran, Y. Yang, T.L. Smith, R.D. Mass, R.S. Herbst, Trastuzumab in combination with cisplatin and gemcitabine in patients with Her2-overexpressing, untreated, advanced non-small cell lung cancer: report of a phase II trial and findings regarding optimal identification of patients with Her2-overexpressing disease, *Lung Cancer* 44 (1) (2004) 99–110, <https://doi.org/10.1016/j.lungcan.2003.09.026>.
- [90] A.K. Ganti, M. Rothe, P.K. Mangat, E. Garrett-Mayer, E.G. Dib, H.L. Duvivier, E. R. Ahn, D. Behl, H. Borghaei, A.S. Balmanoukian, A. Gaba, R. Li, K. Osei-Boateng, R. Thota, G.N. Grantham, A. Gregory, S. Halabi, R.L. Schilsky, Pertuzumab Plus Trastuzumab in Patients With Lung Cancer With ERBB2 Mutation or Amplification: Results From the Targeted Agent and Profiling Utilization Registry Study, *JCO Precis Oncol* 7 (2023) e2300041.
- [91] C.H. Chau, P.S. Steeg, W.D. Figg, Antibody-drug conjugates for cancer, *Lancet* 394 (10200) (2019) 793–804, [https://doi.org/10.1016/S0140-6736\(19\)31774-X](https://doi.org/10.1016/S0140-6736(19)31774-X).
- [92] B.T. Li, R. Shen, D. Buonocore, Z.T. Olah, A. Ni, M.S. Ginsberg, G.A. Ulaner, M. Offin, D. Feldman, T. Hembrough, F. Cecchi, S. Schwartz, N. Pavlakis, S. Clarke, H.H. Won, E.B. Brzostowski, G.J. Riely, D.B. Solit, D.M. Hyman, A. Drilon, C.M. Rudin, M.F. Berger, J. Baselga, M. Scaltriti, M.E. Arcila, M.G. Kris, Ado-Trastuzumab Emtansine for Patients With HER2-Mutant Lung Cancers: Results From a Phase II Basket Trial, *J Clin Oncol* 36 (24) (2018) 2532–2537, <https://doi.org/10.1200/jco.2018.77.9777>.
- [93] G.D. Lewis Phillips, G. Li, D.L. Dugger, L.M. Crocker, K.L. Parsons, E. Mai, W. A. Blättler, J.M. Lambert, R.V. Chari, R.J. Lutz, W.L. Wong, F.S. Jacobson, H. Koepfen, R.H. Schwall, S.R. Kenkare-Mitra, S.D. Spencer, M.X. Sliwkowski, Targeting HER2-positive breast cancer with trastuzumab-DM1, an antibody-cytotoxic drug conjugate, *Cancer Res* 68 (22) (2008) 9280–9290, <https://doi.org/10.1158/0008-5472.Can-08-1776>.
- [94] E. Iwama, Y. Zenke, S. Sugawara, H. Daga, M. Morise, N. Yanagitani, T. Sakamoto, H. Murakami, J. Kishimoto, S. Matsumoto, Y. Nakanishi, K. Goto, I. Okamoto, Trastuzumab emtansine for patients with non-small cell lung cancer positive for human epidermal growth factor receptor 2 exon-20 insertion mutations, *Eur J Cancer* 162 (2022) 99–106, <https://doi.org/10.1016/j.ejca.2021.11.021>.
- [95] K. Hotta, K. Aoe, T. Kozuki, K. Ohashi, K. Ninomiya, E. Ichihara, T. Kubo, T. Ninomiya, K. Chikamori, D. Harada, N. Nogami, T. Hirata, S. Hinotsu, S. Toyooka, K. Kiura, A Phase II Study of Trastuzumab Emtansine in HER2-Positive Non-Small Cell Lung Cancer, *J Thorac Oncol* 13 (2) (2018) 273–279, <https://doi.org/10.1016/j.jtho.2017.10.032>.
- [96] S. Peters, R. Stahel, L. Bubendorf, P. Bonomi, A. Villegas, D.M. Kowalski, C. S. Baik, D. Isla, J.C. Carpeno, P. Garrido, A. Rittmeyer, M. Tiseo, C. Meyenberger, S. de Haas, L.H. Lam, M.W. Lu, T.E. Stinchcombe, Trastuzumab Emtansine (T-DM1) in Patients with Previously Treated HER2-Overexpressing Metastatic Non-Small Cell Lung Cancer: Efficacy, Safety, and Biomarkers, *Clin Cancer Res* 25 (1) (2019) 64–72, <https://doi.org/10.1158/1078-0432.Ccr-18-1590>.
- [97] T. Nakada, T. Masuda, H. Naito, M. Yoshida, S. Ashida, K. Morita, H. Miyazaki, Y. Kasuya, Y. Ogita, J. Yamaguchi, Y. Abe, T. Honda, Novel antibody drug conjugates containing exatecan derivative-based cytotoxic payloads, *Bioorg Med Chem Lett* 26 (6) (2016) 1542–1545, <https://doi.org/10.1016/j.bmcl.2016.02.020>.
- [98] Y. Ogita, K. Hagihara, M. Oitate, H. Naito, T. Agatsuma, Bystander killing effect of DS-8201a, a novel anti-human epidermal growth factor receptor 2 antibody-drug conjugate, in tumors with human epidermal growth factor receptor 2 heterogeneity, *Cancer Sci* 107 (7) (2016) 1039–1046, <https://doi.org/10.1111/cas.12966>.
- [99] J. Tsurutani, H. Iwata, I. Krop, P.A. Jänne, T. Doi, S. Takahashi, H. Park, C. Redfern, K. Tamura, T.M. Wise-Draper, K. Saito, M. Sugihara, J. Singh, T. Jikoh, G. Gallant, B.T. Li, Targeting HER2 with Trastuzumab Deruxtecane: A Dose-Expansion, Phase I Study in Multiple Advanced Solid Tumors, *Cancer Discov* 10 (5) (2020) 688–701, <https://doi.org/10.1158/2159-8290.Cd-19-1014>.
- [100] K. Nakagawa, M. Nagasaka, E. Felip, J. Pacheco, C. Baik, Y. Goto, A. Salto, B. Li, H. Udagawa, S. Gadgil, H. Murakami, D. Planchard, L. Bazhenova, L. Paz-Ares, M. Perol, J. Mazieres, F. Barlesi, K. Saxena, R. Shiga, S. Acharyya, Y. Cheng, J. Shahidi, P. Jänne, E. Smit, OA04.05 Trastuzumab Deruxtecane in HER2-Overexpressing Metastatic Non-Small Cell Lung Cancer: Interim Results of DESTINY-Lung01, *J Thorac Oncol* 16 (3) (2021) S109–S110, <https://doi.org/10.1016/j.jtho.2021.01.285>.
- [101] B.T. Li, E.F. Smit, Y. Goto, K. Nakagawa, H. Udagawa, J. Mazieres, M. Nagasaka, L. Bazhenova, A.N. Salto, E. Felip, J.M. Pacheco, M. Pérol, L. Paz-Ares, K. Saxena, R. Shiga, Y. Cheng, S. Acharyya, P. Vitazka, J. Shahidi, D. Planchard, P.A. Jänne, Trastuzumab Deruxtecane in HER2-Mutant Non-Small-Cell Lung Cancer, *N Engl J Med* 386 (3) (2021) 241–251, <https://doi.org/10.1056/NEJMoa2112431>.
- [102] B.T. Li, F. Micheline, S. Misale, E. Cocco, L. Baldino, Y. Cai, S. Shifman, H.Y. Tu, M.L. Myers, C. Xu, M. Mattar, I. Khodos, M. Little, B. Qeriqi, G. Weitsman, C. J. Wilhem, A.S. Lalani, I. Dila, R.A. Freedman, N.U. Lin, D.B. Solit, M.F. Berger, P.R. Barber, T. Ng, M. Offin, J.M. Isbell, D.R. Jones, H.A. Yu, S. Thyparambil, W. L. Liao, A. Bhalkikar, F. Cecchi, D.M. Hyman, J.S. Lewis, D.J. Buonocore, A.L. Ho, V. Makker, J.S. Reis-Filho, P. Razavi, M.E. Arcila, M.G. Kris, J.T. Poirier, R. Shen, J. Tsurutani, G.A. Ulaner, E. de Stanchina, N. Rosen, C.M. Rudin, M. Scaltriti, HER2-Mediated Internalization of Cytotoxic Agents in ERBB2 Amplified or Mutant Lung Cancers, *Cancer Discov* 10 (5) (2020) 674–687, <https://doi.org/10.1158/2159-8290.Cd-20-0215>.
- [103] S.M. Swain, M. Nishino, L.H. Lancaster, B.T. Li, A.G. Nicholson, B.J. Bartholmai, J. Naidoo, E. Schumacher-Wulf, K. Shitara, J. Tsurutani, P. Conte, T. Kato, F. Andre, C.A. Powell, Multidisciplinary clinical guidance on trastuzumab deruxtecane (T-DXd)-related interstitial lung disease/pneumonitis-Focus on proactive monitoring, diagnosis, and management, *Cancer Treat Rev* 106 (2022), 102378, <https://doi.org/10.1016/j.ctrv.2022.102378>.
- [104] K. Goto, K. Sang-We, T. Kubo, Y. Goto, M.J. Ahn, D. Planchard, D.W. Kim, J.C. H. Yang, T.Y. Yang, K.M.C. Pereira, K. Saxena, R. Shiga, Y. Cheng, M. Aregay, P. A. Jänne, LBA55 Trastuzumab deruxtecane (T-DXd) in patients (Pts) with HER2-mutant metastatic non-small cell lung cancer (NSCLC): Interim results from the phase 2 DESTINY-Lung02 trial, *Ann Oncol* 33 (2022) S1422, <https://doi.org/10.1016/j.annonc.2022.08.057>.
- [105] Discovery 12 (10) (2022) 2224, <https://doi.org/10.1158/2159-8290.Cd-nb2022-0053>.
- [106] D. Blangé, C.I. Strees, S. Derks, M.F. Bijlsma, H.W.M. van Laarhoven, Resistance mechanisms to HER2-targeted therapy in gastroesophageal adenocarcinoma: A systematic review, *Cancer Treat Rev* 108 (2022), 102418, <https://doi.org/10.1016/j.ctrv.2022.102418>.
- [107] M. Gámez-Chiachio, D. Sarrió, G. Moreno-Bueno, Novel Therapies and Strategies to Overcome Resistance to Anti-HER2-Targeted Drugs, *Cancers (basel)* 14 (18) (2022), <https://doi.org/10.3390/cancers14184543>.
- [108] D.S. Neel, T.G. Bivona, Resistance is futile: overcoming resistance to targeted therapies in lung adenocarcinoma, *npj Precis Oncol* 1 (2017), <https://doi.org/10.1038/s41698-017-0007-0>.
- [109] D. Planchard, J.C.H. Yang, J.R. Brahmer, A. Ragone, J. Chen, F. Liu, M. Saggese, 185TIP A phase Ib dose-escalation study evaluating trastuzumab deruxtecane (T-DXd) and durvalumab in combination with chemotherapy as first-line treatment in patients with advanced or metastatic nonsquamous non-small cell lung cancer (NSCLC) and HER2 overexpression (DESTINY-Lung03), *J Thorac Oncol* 16 (4) (2021) S798, [https://doi.org/10.1016/S1556-0864\(21\)02027-X](https://doi.org/10.1016/S1556-0864(21)02027-X).
- [110] B. Besse, M. Awad, P. Forde, M. Thomas, K. Park, G. Goss, N. Rizvi, F. Huemer, M. Hochmair, J. Bannoun, J. Cosaert, Z. Szucs, P. Mortimer, R. Hobson, K. Sachsenmeier, E. Dean, H. Ambrose, C. Hayward, M. Dressman, S. Barry, J. Heymach, OA07.08 HUDSON: An Open-Label, Multi-Drug, Biomarker-Directed, Phase II Platform Study in Patients with NSCLC, who Progressed on Anti-PD(L)1 Therapy, *J Thorac Oncol* 16 (3) (2021) S118–S119, <https://doi.org/10.1016/j.jtho.2021.01.299>.
- [111] C.A.W. Geuijen, C. De Nardis, D. Maussang, E. Rovers, T. Gallenle, L.J. A. Hendriks, T. Visser, R. Nijhuis, T. Logtenberg, J. de Kruif, P. Gros, M. Throsby, Unbiased Combinatorial Screening Identifies a Bispecific IgG1 that Potently Inhibits HER3 Signaling via HER2-Guided Ligand Blockade, *Cancer Cell* 33 (5) (2018) 922–936, <https://doi.org/10.1016/j.ccell.2018.04.003>.
- [112] F. Tang, Y. Yang, Y. Tang, S. Tang, L. Yang, B. Sun, B. Jiang, J. Dong, H. Liu, M. Huang, M.-Y. Geng, W. Huang, One-pot N-glycosylation remodeling of IgG with non-natural sialylglycopeptides enables glycosite-specific and dual-payload antibody–drug conjugates, *Org Biomol Chem* 14 (40) (2016) 9501–9518, <https://doi.org/10.1039/C6OB01751G>.
- [113] Y. Yu, Y. Yang, H. Li, Y. Fan, Targeting HER2 alterations in non-small cell lung cancer: Therapeutic breakthrough and challenges, *Cancer Treat Rev* 114 (2023), 102520, <https://doi.org/10.1016/j.ctrv.2023.102520>.
- [114] D. Tan, K. Goto, H.H. Loong, T.-Y. Yang, S. Gadgil, E. Massarelli, G. Daniele, X. Le, Y. Shinno, H. Murakami, T. Sakamoto, J.C. Ruffinelli, S. Lu, Y. Zhang, T. M. Kim, B.J. Brennan, C. Chen, M. Joosten, Z. Wang, S.-F. Pu, W. Bao, M. Kormacker, R. Ferraldeschi, P. Grassi, B.C. Goh, Abstract CT126: An open-label, first-in-human study of BAY2927088 in patients with advanced non-small cell lung cancer (NSCLC) harboring an EGFR and/or HER2 mutation, *Cancer Res* 83 (8, Supplement) (2023), <https://doi.org/10.1158/1538-7445.Am2023-ct126>.
- [115] Y. Fan, J. Chen, C. Zhou, H. Wang, Y. Shu, J. Zhang, H. Hua, D.C. Huang, C. Zhou, Afatinib in patients with advanced non-small cell lung cancer harboring HER2 mutations, previously treated with chemotherapy: A phase II trial, *Lung Cancer* 147 (2020) 209–213, <https://doi.org/10.1016/j.lungcan.2020.07.017>.
- [116] B.C. Cho, J. Lin, D.R. Camidge, V. Velcheti, B. Solomon, S. Lu, K.H. Lee, S.W. Kim, S. Kao, R. Dziadziuszko, M. Beg, M. Nagasaka, E. Felip, B. Besse, C. Springfield, S. Popat, J. Wolf, D. Trone, S. Stopatschinskaja, A. Drilon, Pivotal topline data from the phase 1/2 TRIDENT-1 trial of repotrectinib in patients with ROS1+ advanced non-small cell lung cancer (NSCLC), *Eur J Cancer* 174 (2022) S1–S2, [https://doi.org/10.1016/S0959-8049\(22\)00812-7](https://doi.org/10.1016/S0959-8049(22)00812-7).
- [117] A. Prelaj, A. Bottiglieri, C. Proto, G. Lo Russo, D. Signorelli, R. Ferrara, G. Galli, A. De Toma, G. Viscardi, M. Brambilla, R. Lobefaro, F. Nichetti, S. Manglaviti, M. Occhipinti, A. Labianca, M. Gannellini, R. Gallucci, N. Zilembo, G.F. Greco, V. Torri, F. de Braud, M.C. Garassino, Pozitotinib for EGFR and HER2 exon 20 insertion mutation in advanced NSCLC: Results from the expanded access

- program, *Eur J Cancer* 149 (2021) 235–248. <https://doi.org/10.1016/j.ejca.2021.02.038>.
- [118] Y.Y. Elamin, J.P. Robichaux, B.W. Carter, M. Altan, D.L. Gibbons, F.V. Fossella, V. K. Lam, A.B. Patel, M.V. Negrao, X. Le, F.E. Mott, J. Zhang, L. Feng, G. Blumenschein Jr., A.S. Tsao, J.V. Heymach, *Pozotinib for Patients With HER2 Exon 20 Mutant Non-Small-Cell Lung Cancer: Results From a Phase II Trial*, *J Clin Oncol* 40 (7) (2022) 702–709, <https://doi.org/10.1200/jco.21.01113>.
- [119] X. Chu, H. Qiang, M. Xie, X. Li, J. Zhao, Y. Wu, J. Zhou, J. Ye, C. Zhao, C. Han, T. Chu, C. Su, *Treatment efficacy of HER2-mutant lung adenocarcinoma by immune checkpoint inhibitors: a multicenter retrospective study*, *Cancer Immunol Immunother* 71 (7) (2022) 1625–1631, <https://doi.org/10.1007/s00262-021-03100-5>.
- [120] B. Besse, J.C. Soria, B. Yao, M. Kris, B. Chao, A. Cortot, J. Mazieres, M.A. Socinski, L. Horn, S. Waqar, F. Barlesi, J.E. Gray, D. Moro-Sibilot, A. Oton, E. Quoix, A. Lalani, L. McCulloch, R. Bryce, L. Gandhi, *Neratinib (N) with or Without Temsirolimus (Tem) in Patients (Pts) with Non-Small Cell Lung Cancer (NscLc) Carrying Her2 Somatic Mutations: an International Randomized Phase II Study*, *Ann Oncol* 25 (2014), v1, <https://doi.org/10.1093/annonc/mdu438.47>.
- [121] Y. Wang, T. Jiang, Z. Qin, J. Jiang, Q. Wang, S. Yang, C. Rivard, G. Gao, T.L. Ng, M.M. Tu, H. Yu, H. Ji, C. Zhou, S. Ren, J. Zhang, P. Bunn, R.C. Doebele, D. R. Camidge, F.R. Hirsch, *HER2 exon 20 insertions in non-small-cell lung cancer are sensitive to the irreversible pan-HER receptor tyrosine kinase inhibitor pyrotinib*, *Ann Oncol* 30 (3) (2019) 447–455, <https://doi.org/10.1093/annonc/mdy542>.
- [122] R.S. Herbst, A.M. Davies, R.B. Natale, T.P. Dang, J.H. Schiller, L.L. Garland, V. A. Miller, D. Mendelson, A.D. Van den Abbeele, Y. Melenevsky, D.J. de Vries, D. A. Eberhard, B. Lyons, S.G. Lutzker, B.E. Johnson, *Efficacy and Safety of Single-Agent Pertuzumab, a Human Epidermal Receptor Dimerization Inhibitor, Patients with Non-Small Cell Lung Cancer*, *Clinical Cancer Research* 13 (20) (2007) 6175–6181, <https://doi.org/10.1158/1078-0432.Ccr-07-0460>.
- [123] J. Mazieres, C. Lafitte, C. Ricordel, L. Greillier, E. Negre, G. Zalcman, C. Domblides, J. Madelaine, J. Bennouna, C. Mascaux, D. Moro-Sibilot, F. Pinquie, A.B. Cortot, J. Otto, J. Cadranel, A. Langlais, F. Morin, V. Westeel, B. Besse, *Combination of Trastuzumab, Pertuzumab, and Docetaxel in Patients With Advanced Non-Small-Cell Lung Cancer Harboring HER2 Mutations: Results From the IFCT-1703 R2D2 Trial*, *J Clin Oncol* 40 (7) (2022) 719–728, <https://doi.org/10.1200/jco.21.01455>.