


# Secant Method

# Secant Method

- Secant method is an open method. That is, no more interval under consideration needs to bracket the root, though it still requires two initial guesses. Let us denote these initial guesses by  $x_{-1}$  and  $x_0$ . The formula for generating the sequence of approximations is

$$x_{i+1} = \frac{x_{i-1}f(x_i) - x_i f(x_{i-1})}{f(x_i) - f(x_{i-1})}$$



# Secant Method

- Firstly, initial choices of  $x_{-1}$  and  $x_0$  need not be bracketing the root. So, that is, root need not lie within the endpoints  $x_{-1}$  and  $x_0$ . That is  $f(x_{-1})$  and  $f(x_0)$  can be of same sign. No more, one needs to ensure that  $f(x_{-1}) \cdot f(x_0) < 0$ .
- Though, in practice one usually chooses  $x_{-1}$  and  $x_0$  as ones which bracket the root, there is no such compulsion.

# Secant Method

- Secondly, in computation of next approximation, older approximation is discarded. Recent most two approximations are used for iteration, irrespective of function values at these end points. So graphically, straight line joining  $(x_{i-1}, f(x_{i-1}))$  and  $(x_i, f(x_i))$  is drawn to generate  $x_{i+1}$ , next approximation. No more, function values sign are checked to ensure that root lies between  $(x_{i-1}, f(x_{i-1}))$  and  $(x_i, f(x_i))$ . Because secant is drawn joining  $(x_{i-1}, f(x_{i-1}))$  and  $(x_i, f(x_i))$ , it is called Secant Method.

# Algorithm

The steps of the Secant method to find the root of an equation  $f(x) = 0$  are

1. Let  $k = 1$ , Evaluate  $f(x_i), f(x_{i-1})$
2. Use an initial guess of the root,  $x_i$ , to estimate the new value of the root,  $x_{i+1}$ , as

$$x_{i+1} = \frac{x_{i-1}f(x_i) - x_i f(x_{i-1})}{f(x_i) - f(x_{i-1})}$$

3. Compare the absolute approximate error  $|c_k - c_{k-1}|$  or absolute relative approximate error  $\frac{|c_k - c_{k-1}|}{|c_k|}$  with the pre-specified relative error tolerance  $\epsilon$ .

where

$c_k$  = estimated root from present iteration

$c_{k-1}$  = estimated root from previous iteration

If  $|c_k - c_{k-1}| < \epsilon$  or  $\frac{|c_k - c_{k-1}|}{|c_k|} < \epsilon$  then exit

Else

$k = k + 1$

go to Step 2

Note one should also check whether the number of iterations is more than the maximum number of iterations allowed. If so, one needs to terminate the algorithm and notify the user about it.

# Advantages and Disadvantages

## Advantages:

1. No constraint of end points of interval to converge
2. Very rapid convergence

## Disadvantages:

1. If initial guess is not sufficiently near the root, it may diverge.


# Example

This polynomial obviously has roots at  $x = 2$  and at  $x = 4$ ; one is a double root, the other a triple root:

$$\begin{aligned} f(x) &= (x - 2)^3(x - 4)^2 \\ &= x^5 - 14x^4 + 76x^3 - 200x^2 + 256x - 128 \end{aligned}$$

- a. Which root can you get with bisection? Which root can't you get?
- b. Repeat part (a) with the secant method.
- c. If you begin with the interval  $[1, 5]$ , which root will you get with  
(1) bisection, (2) the secant method, (3) false position?
- d. Use Newton's method with  $x_0 = 3$ . Does it converge? To which root?

# Example





# Example

This polynomial obviously has roots at  $x = 2$  and at  $x = 4$ ; one is a double root, the other a triple root:

$$\begin{aligned} f(x) &= (x - 2)^3(x - 4)^2 \\ &= x^5 - 14x^4 + 76x^3 - 200x^2 + 256x - 128 \end{aligned}$$

a.

b. Both roots can be reached with secant method

c. If you begin with the interval  $[1, 5]$ , which root will you get with  
(2) the secant method: It will converge to 2

d.

# Example

Eg 1. *Find the real root of the equation  $x^3 - 4x - 9 = 0$  by Secant Method correct.*

*Take  $x_0 = 2, x_1 = 3, \epsilon = 0.01$*

Eg 2. *Find the real root of the equation  $x^3 - x - 1 = 0$  by Secant Method correct. Take  $x_0 = 1, x_1 = 1.5, \epsilon = 0.001$*

Eg 3.  $f(x) = 3x - e^x$  starting from 0,1 and  $\epsilon = 0.01$

# Secant Method

Eg 1. *Find the real root of the equation  $x^3 - 4x - 9 = 0$  by Secant Method correct.*

x	0	1	2	3
f(x)	-9	-12	-9	6

*Take  $x_0 = 2$ ,  $x_1 = 3$ ,  $\epsilon = 0.01$*

# Secant Method

Eg 1. *Find the real root of the equation  $x^3 - 4x - 9 = 0$  by Secant Method correct. Take  $x_0 = 2$ ,  $x_1 = 3$ ,  $\epsilon = 0.001$*

$$f(x) = x^3 - 4x - 9$$

$$f(2) = -9$$

and

$$f(3) = 6$$

$x_0$	$f(x_0)$	$x_1$	$f(x_1)$
2	-9	3	6

# Secant Method

First approximation to the root is

$$x_2 = \frac{(2)(6) - (3)(-9)}{6 - (-9)}$$

$$x_2 = 2.6$$

Now  $f(x_2) = -1.824$

$x_0$	$f(x_0)$	$x_1$	$f(x_1)$	$x_2$	$f(x_2)$	$ x_k - x_{k-1} $
2	-9	3	6	2.6	-1.824	-

# Secant Method

$$x_0 = 3, x_1 = 2.6$$

Second approximation to the root is

$$x_2 = \frac{(3)(-1.824) - (2.6)6}{(-1.824) - 6}$$

$$= 2.6933$$

$$\text{Now } f(x_2) = -0.2372$$

$x_0$	$f(x_0)$	$x_1$	$f(x_1)$	$x_2$	$f(x_2)$	$ x_k - x_{k-1} $
2	-9	3	6	2.6	-1.824	-
3	6	2.6	-1.824	2.6933	-0.2372	0.0933

# Secant Method

$$x_0 = 2.6, x_1 = 2.6933$$

Third approximation to the root is

$$x_2 = \frac{(2.6)(-0.2372) - (2.6933)(-1.824)}{(-0.2372) - (-1.824)}$$
$$= 2.7072$$

$$\text{Now } f(x_2) = 0.012$$

$x_0$	$f(x_0)$	$x_1$	$f(x_1)$	$x_2$	$f(x_2)$	$ x_k - x_{k-1} $
2	-9	3	6	2.6	-1.824	-
3	6	2.6	-1.824	2.6933	-0.2372	0.0933
2.6	-1.824	2.6933	-0.2372	2.7072	0.012	0.0139

# Secant Method

$$x_0 = 2.6933, x_1 = 2.7072$$

Fourth approximation to the root is

$$x_2 = \frac{(2.6933)(0.012) - (2.7072)(-0.2372)}{0.012 - (-0.2372)} = 2.7065$$

$$\epsilon = 0.001, |x_k - x_{k-1}| = 0.0007$$

Hence, the root is 2.7065, correct to three decimal places.

$x_0$	$f(x_0)$	$x_1$	$f(x_1)$	$x_2$	$f(x_2)$	$ x_k - x_{k-1} $
2	-9	3	6	2.6	-1.824	-
3	6	2.6	-1.824	2.6933	-0.2372	0.0933
2.6	-1.824	2.6933	-0.2372	2.7072	0.012	0.0139
2.6933	-0.2372	2.7072	0.012	2.7065	0	0.0007



# Secant Method

Eg 2. Find the real root of the equation  $x^3 - x - 1 = 0$  by Secant Method correct. Take  $x_0 = 1$ ,  $x_1 = 1.5$ ,  $\epsilon = 0.001$

$$f(x) = x^3 - x - 1$$

$$f(1) = -1$$

and

$$f(1.5) = 0.875$$

$x_0$	$f(x_0)$	$x_1$	$f(x_1)$
1	-1	1.5	0.875

# Secant Method

First approximation to the root is

$$x_2 = \frac{(1)(0.875) - (1.5)(-1)}{0.875 - (-1)}$$

$$x_2 = 1.2667$$

Now  $f(x_2) = -0.2344$

$x_0$	$f(x_0)$	$x_1$	$f(x_1)$	$x_2$	$f(x_2)$	$ x_k - x_{k-1} $
1	-1	1.5	0.875	1.2667	-0.2344	-

# Secant Method

$$x_0 = 1.5, x_1 = 1.2667$$

Second approximation to the root is

$$x_2 = \frac{(1.5)(-0.2344) - (1.2667)0.875}{(-0.2344) - 0.875}$$

$$= 1.316$$

$$\text{Now } f(x_2) = -0.037$$

$x_0$	$f(x_0)$	$x_1$	$f(x_1)$	$x_2$	$f(x_2)$	$ x_k - x_{k-1} $
1	-1	1.5	0.875	1.2667	-0.2344	-
1.5	0.875	1.2667	-0.2344	1.316	-0.037	0.0493

# Secant Method

$$x_0 = 1.2667, x_1 = 1.316$$

Third approximation to the root is

$$x_2 = \frac{(1.2667)(-0.037) - (1.316)(-0.2344)}{(-0.037) - (-0.2344)}$$
$$= 1.3252$$

$$\text{Now } f(x_2) = 0.0021$$

$x_0$	$f(x_0)$	$x_1$	$f(x_1)$	$x_2$	$f(x_2)$	$ x_k - x_{k-1} $
1	-1	1.5	0.875	1.2667	-0.2344	-
1.5	0.875	1.2667	-0.2344	1.316	-0.037	0.0493
1.2667	-0.2344	1.316	-0.037	1.3252	0.0021	0.0092

# Secant Method

$$x_0 = 1.316, x_1 = 1.3252$$

Fourth approximation to the root is

$$x_2 = \frac{(1.316)(0.0021) - (1.3252)(-0.037)}{0.0021 - (-0.037)} = 1.3247$$

$$\epsilon = 0.001, |x_k - x_{k-1}| = 0.0005$$

Hence, the root is 1.3247, correct to three decimal places.

$x_0$	$f(x_0)$	$x_1$	$f(x_1)$	$x_2$	$f(x_2)$	$ x_k - x_{k-1} $
1	-1	1.5	0.875	1.2667	-0.2344	-
1.5	0.875	1.2667	-0.2344	1.316	-0.037	0.0493
1.2667	-0.2344	1.316	-0.037	1.3252	0.0021	0.0092
1.316	-0.037	1.3252	0.0021	1.3247	0	0.0005

# Example

- $f(x) = 3x - e^x$  starting from 0,1 and  $\epsilon = 0.01$

$x_0$	$f(x_0)$	$x_1$	$f(x_1)$	$x_2$	$f(x_2)$	$ x_k - x_{k-1} $
0	-1	1	0.2817	0.7802	0.1587	-
1	0.2817	0.7802	0.1587	0.4967	-0.1532	0.2835
0.7802	0.1587	0.4967	-0.1532	0.636	0.019	0.1393
0.4967	-0.1532	0.636	0.019	0.6206	0.0017	0.0154
0.636	0.019	0.6206	0.0017	0.619	0	0.0016