

Problems **1** and **2** are *optional*, as they should be familiar from QFT I. However, if you are not comfortable with manipulating Gamma matrices, I *encourage* you to complete them. Completing them will result in *bonus points*.

1. The Dirac matrices $\gamma^\mu = (\gamma^0, \gamma^j)$ in the chiral (Weyl) representation are defined as

$$\gamma^0 = \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix}, \quad \gamma^j = \begin{pmatrix} 0 & \sigma^j \\ -\sigma^j & 0 \end{pmatrix},$$

where I is the 2×2 identity matrix and σ^j are the Pauli matrices.

- (a) With this representation, confirm that $\{\gamma^\mu, \gamma^\nu\} = 2g^{\mu\nu}$.

Solution: To clarify some of the manipulations in these problems, we introduce I_4 as the 4×4 identity, and let $I \rightarrow I_2$ be the 2×2 identity. Thus, what is to be shown is $\{\gamma^\mu, \gamma^\nu\} = 2g^{\mu\nu} I_4$, given the chiral representation

$$\gamma^0 = \begin{pmatrix} 0 & I_2 \\ I_2 & 0 \end{pmatrix}, \quad \gamma^j = \begin{pmatrix} 0 & \sigma^j \\ -\sigma^j & 0 \end{pmatrix}.$$

Recall the properties of the Pauli matrices, $\{\sigma^j, \sigma^k\} = 2\delta^{jk} I_2$.

$$\begin{aligned} \{\gamma^0, \gamma^0\} &= 2(\gamma^0)^2, \\ &= 2 \begin{pmatrix} 0 & I_2 \\ I_2 & 0 \end{pmatrix} \begin{pmatrix} 0 & I_2 \\ I_2 & 0 \end{pmatrix}, \\ &= 2 \begin{pmatrix} I_2 & 0 \\ 0 & I_2 \end{pmatrix} = 2g^{00} I_4, \\ \{\gamma^0, \gamma^j\} &= \gamma^0 \gamma^j + \gamma^j \gamma^0, \\ &= \begin{pmatrix} 0 & I_2 \\ I_2 & 0 \end{pmatrix} \begin{pmatrix} 0 & \sigma^j \\ -\sigma^j & 0 \end{pmatrix} + \begin{pmatrix} 0 & \sigma^j \\ -\sigma^j & 0 \end{pmatrix} \begin{pmatrix} 0 & I_2 \\ I_2 & 0 \end{pmatrix}, \\ &= \begin{pmatrix} -\sigma^j & 0 \\ 0 & \sigma^j \end{pmatrix} + \begin{pmatrix} \sigma^j & 0 \\ 0 & -\sigma^j \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} = 2g^{0j} I_4, \end{aligned}$$

where we note that $g^{00} = +1$ and $g^{0j} = g^{j0} = 0$. Continuing,

$$\begin{aligned}
 \{\gamma^j, \gamma^0\} &= \gamma^j \gamma^0 + \gamma^0 \gamma^j = \{\gamma^0, \gamma^j\} = 2g^{0j} I_4 = 2g^{j0} I_4, \\
 \{\gamma^j, \gamma^k\} &= \gamma^j \gamma^k + \gamma^k \gamma^j, \\
 &= \begin{pmatrix} 0 & \sigma^j \\ -\sigma^j & 0 \end{pmatrix} \begin{pmatrix} 0 & \sigma^k \\ -\sigma^k & 0 \end{pmatrix} + \begin{pmatrix} 0 & \sigma^k \\ -\sigma^k & 0 \end{pmatrix} \begin{pmatrix} 0 & \sigma^j \\ -\sigma^j & 0 \end{pmatrix}, \\
 &= \begin{pmatrix} -\sigma^j \sigma^k & 0 \\ 0 & -\sigma^j \sigma^k \end{pmatrix} + \begin{pmatrix} -\sigma^k \sigma^j & 0 \\ 0 & -\sigma^k \sigma^j \end{pmatrix}, \\
 &= - \begin{pmatrix} \sigma^j \sigma^k + \sigma^k \sigma^j & 0 \\ 0 & \sigma^j \sigma^k + \sigma^k \sigma^j \end{pmatrix}, \\
 &= - \begin{pmatrix} 2\delta^{jk} I_2 & 0 \\ 0 & 2\delta^{jk} I_2 \end{pmatrix}, \\
 &= -2\delta^{jk} I_4 = 2g^{jk} I_4
 \end{aligned}$$

Therefore, we have shown $\{\gamma^\mu, \gamma^\nu\} = 2g^{\mu\nu} I_4$

- (b) Using the result in (a), show that $\gamma_\mu \gamma^\mu = 4$.

Solution: Contract $\{\gamma^\mu, \gamma^\nu\} = 2g^{\mu\nu} I_4$ with $g_{\mu\nu}$,

$$\begin{aligned}
 g_{\mu\nu} \{\gamma^\mu, \gamma^\nu\} &= 2g_{\mu\nu} g^{\mu\nu} I_4, \\
 \{\gamma_\mu, \gamma^\mu\} &= 2g^\mu{}_\mu I_4, \\
 2\gamma_\mu \gamma^\mu &= 2 \cdot 4 I_4.
 \end{aligned}$$

So, we conclude $\gamma_\mu \gamma^\mu = 4 I_4$.

- (c) Prove that $\gamma_\mu \gamma^\nu \gamma^\mu = -2\gamma^\nu$ without using an explicit matrix representation.

Solution: Using the anticommutator relation, as well as the result from part (b), we find

$$\begin{aligned}
 \gamma_\mu \gamma^\nu \gamma^\mu &= \gamma_\mu (2g^{\mu\nu} I_4 - \gamma^\mu \gamma^\nu), \\
 &= 2\gamma_\nu - \gamma_\mu \gamma^\mu \gamma^\nu, \\
 &= 2\gamma_\nu - 4\gamma^\nu, \\
 &= -2\gamma^\nu.
 \end{aligned}$$

- (d) Similarly, prove that $\gamma_\mu \gamma^\nu \gamma^\rho \gamma^\mu = 4g^{\nu\rho}$.

Solution: Using the anticommutator relation, as well as the result from part (c), we find

$$\begin{aligned}
 \gamma_\mu \gamma^\nu \gamma^\rho \gamma^\mu &= \gamma_\mu \gamma^\nu (2g^{\rho\mu} I_4 - \gamma^\mu \gamma^\rho), \\
 &= 2\gamma^\rho \gamma^\nu - \gamma_\mu \gamma^\nu \gamma^\mu \gamma^\rho, \\
 &= 2\gamma^\rho \gamma^\nu + 2\gamma^\nu \gamma^\rho, \\
 &= 2\{\gamma^\rho, \gamma^\nu\}, \\
 &= 4g^{\nu\rho} I_4.
 \end{aligned}$$

2. Given $\gamma^5 = \gamma_5 = i\gamma^0\gamma^1\gamma^2\gamma^3$, prove the following trace identities:

(a) $\text{tr}(\gamma^\mu \gamma^\nu) = 4g^{\mu\nu}$,

Solution: Taking the trace, we use the cyclic properties of the trace and the anticommutation relations, we have

$$\begin{aligned}
 \text{tr}(\gamma^\mu \gamma^\nu) &= \frac{1}{2} \text{tr}(\gamma^\mu \gamma^\nu + \gamma^\mu \gamma^\nu), \\
 &= \frac{1}{2} [\text{tr}(\gamma^\mu \gamma^\nu) + \text{tr}(\gamma^\mu \gamma^\nu)], \\
 &= \frac{1}{2} \text{tr}(\gamma^\mu \gamma^\nu + \gamma^\nu \gamma^\mu), \\
 &= \frac{1}{2} \text{tr}(\{\gamma^\mu \gamma^\nu\}), \\
 &= \frac{1}{2} \cdot 2g^{\mu\nu} \text{tr}(I_4), \\
 &= 4g^{\mu\nu},
 \end{aligned}$$

where $\text{tr}(I_4) = 4$.

(b) $\text{tr}(\gamma^\mu \gamma^\nu \gamma^\rho \gamma^\sigma) = 4(g^{\mu\nu} g^{\rho\sigma} - g^{\mu\rho} g^{\nu\sigma} + g^{\mu\sigma} g^{\nu\rho}),$

Solution: Here we use the anticommutation relation inside the trace,

$$\begin{aligned}
 \text{tr}(\gamma^\mu \gamma^\nu \gamma^\rho \gamma^\sigma) &= \text{tr}[\gamma^\mu \gamma^\nu (2g^{\rho\sigma} I_4 - \gamma^\sigma \gamma^\rho)], \\
 &= 2g^{\rho\sigma} \text{tr}(\gamma^\mu \gamma^\nu) - \text{tr}(\gamma^\mu \gamma^\nu \gamma^\sigma \gamma^\rho), \\
 &= 8g^{\mu\nu} g^{\rho\sigma} - \text{tr}[\gamma^\mu (2g^{\nu\sigma} I_4 - \gamma^\sigma \gamma^\nu) \gamma^\rho], \\
 &= 8g^{\mu\nu} g^{\rho\sigma} - 2g^{\nu\sigma} \text{tr}(\gamma^\mu \gamma^\rho) + \text{tr}(\gamma^\mu \gamma^\sigma \gamma^\nu \gamma^\rho), \\
 &= 8g^{\mu\nu} g^{\rho\sigma} - 8g^{\nu\sigma} g^{\mu\rho} + \text{tr}[2g^{\mu\sigma} I_4 - \gamma^\sigma \gamma^\mu) \gamma^\nu \gamma^\rho], \\
 &= 8g^{\mu\nu} g^{\rho\sigma} - 8g^{\nu\sigma} g^{\mu\rho} + 2g^{\mu\sigma} \text{tr}(\gamma^\nu \gamma^\rho) - \text{tr}(\gamma^\sigma \gamma^\mu \gamma^\nu \gamma^\rho), \\
 &= 8g^{\mu\nu} g^{\rho\sigma} - 8g^{\nu\sigma} g^{\mu\rho} + 8g^{\mu\sigma} g^{\nu\rho} - \text{tr}(\gamma^\mu \gamma^\nu \gamma^\rho \gamma^\sigma),
 \end{aligned}$$

where in the last line we used the cyclic property of the trace. Then, adding this final trace to the left-hand side, we find

$$\begin{aligned}
 2 \text{tr}(\gamma^\mu \gamma^\nu \gamma^\rho \gamma^\sigma) &= 8g^{\mu\nu} g^{\rho\sigma} - 8g^{\nu\sigma} g^{\mu\rho} + 8g^{\mu\sigma} g^{\nu\rho} \\
 \implies \text{tr}(\gamma^\mu \gamma^\nu \gamma^\rho \gamma^\sigma) &= 4(g^{\mu\nu} g^{\rho\sigma} - g^{\nu\sigma} g^{\mu\rho} + g^{\mu\sigma} g^{\nu\rho})
 \end{aligned}$$

(c) The trace of *any* odd number of gamma matrices is zero.

Solution: We first prove that $\text{tr}(\gamma^\mu) = 0$, which is obvious in the Weyl basis but is true in general. Recall that $(\gamma_5)^2 = \gamma_5 \gamma^5 = 4I_4$. Therefore, the trace can be written as

$$\text{tr}(\gamma^\mu) = \text{tr}(\gamma^\mu I_4) = \text{tr}(\gamma^\mu \gamma^5 \gamma^5) = -\text{tr}(\gamma^5 \gamma^\mu \gamma^5) = -\text{tr}(\gamma^\mu \gamma^5 \gamma^5) = -\text{tr}(\gamma^\mu),$$

where in the fourth equality we used the anticommutation relation $\gamma^\mu \gamma^5 = -\gamma^5 \gamma^\mu$ and in the fifth equality results from the cyclic properties of the trace. Therefore, we conclude

$$\text{tr}(\gamma^\mu) = 0.$$

A generic trace over an odd number of gamma matrices can be written as a trace over $2n + 1$ gamma matrices where $n \in \mathbb{N}$, $\text{tr}(\gamma^{\mu_1} \gamma^{\mu_2} \dots \gamma^{\mu_{2n}} \gamma^{\mu_{2n+1}})$. So, inserting $I_4 = (\gamma^5)^2$ at the end gives,

$$\begin{aligned}
 \text{tr}(\gamma^{\mu_1} \gamma^{\mu_2} \dots \gamma^{\mu_{2n}} \gamma^{\mu_{2n+1}}) &= \text{tr}(\gamma^{\mu_1} \gamma^{\mu_2} \dots \gamma^{\mu_{2n}} \gamma^{\mu_{2n+1}} I_4), \\
 &= \text{tr}(\gamma^{\mu_1} \gamma^{\mu_2} \dots \gamma^{\mu_{2n}} \gamma^{\mu_{2n+1}} \gamma^5 \gamma^5), \\
 &= (-1)^{2n+1} \text{tr}(\gamma^5 \gamma^{\mu_1} \gamma^{\mu_2} \dots \gamma^{\mu_{2n}} \gamma^{\mu_{2n+1}} \gamma^5), \\
 &= -\text{tr}(\gamma^{\mu_1} \gamma^{\mu_2} \dots \gamma^{\mu_{2n}} \gamma^{\mu_{2n+1}} \gamma^5 \gamma^5), \\
 &= -\text{tr}(\gamma^{\mu_1} \gamma^{\mu_2} \dots \gamma^{\mu_{2n}} \gamma^{\mu_{2n+1}}),
 \end{aligned}$$

where the factor $(-1)^{2n+1} = (-1)$ comes from anticommuting γ^5 to the left through all $2n + 1$ gamma matrices. We conclude that $\text{tr}(\gamma^{\mu_1} \gamma^{\mu_2} \dots \gamma^{\mu_{2n}} \gamma^{\mu_{2n+1}}) = 0$

$$(d) \operatorname{tr}(\gamma^5) = \operatorname{tr}(\gamma^5 \gamma^\mu) = \operatorname{tr}(\gamma^5 \gamma^\mu \gamma^\nu) = \operatorname{tr}(\gamma^5 \gamma^\mu \gamma^\nu \gamma^\rho) = 0,$$

Solution: We begin by first proving $\operatorname{tr}(\gamma^5) = 0$. By definition, $\gamma^5 \gamma^\mu = -\gamma^\mu \gamma^5$. Now, taking the trace, and inserting the identity in the form $I_4 = (\gamma^0)^2$,

$$\begin{aligned} \operatorname{tr}(\gamma^5) &= \operatorname{tr}(I_4 \gamma^5), \\ &= \operatorname{tr}(\gamma^0 \gamma^0 \gamma^5), \\ &= -\operatorname{tr}(\gamma^0 \gamma^5 \gamma^0), \\ &= -\operatorname{tr}(\gamma^0 \gamma^0 \gamma^5), \\ &= -\operatorname{tr}(\gamma^5), \end{aligned}$$

where we anticommutated γ^0 to the right going to line 3, and then used the cyclic property of the trace in line 4. Therefore, we conclude $\operatorname{tr}(\gamma^5) = 0$.

We note that since γ^5 is defined as $\gamma^5 = i\gamma^0 \gamma^1 \gamma^2 \gamma^3$, that $\operatorname{tr}(\gamma^5 \gamma^\mu) = \operatorname{tr}(\gamma^5 \gamma^\mu \gamma^\nu \gamma^\rho) = 0$ since this is the trace of an odd number of gamma matrices.

Therefore, the remaining identity to show is $\operatorname{tr}(\gamma^5 \gamma^\mu \gamma^\nu) = 0$. Note that if $\mu = \nu$, then $(\gamma^\mu)^2 = \pm I_4$ where the ‘+’ is for $\mu = 0$, and ‘−’ otherwise. So, if $\mu = \nu$, then $\operatorname{tr}(\gamma^5 \gamma^\mu \gamma^\nu) \rightarrow \pm \operatorname{tr}(\gamma^5 I_4) = 0$ by the first identity proved in this solution. What remains is the case where $\mu \neq \nu$. We insert an identity of the form $I_4 = \pm(\gamma^\rho)^2$, where we are free to choose $\rho \neq \mu$ and $\rho \neq \nu$, so that $\{\gamma^\rho, \gamma^\mu\} = \{\gamma^\rho, \gamma^\nu\} = 0$. Taking the trace,

$$\begin{aligned} \operatorname{tr}(\gamma^5 \gamma^\mu \gamma^\nu) &= \operatorname{tr}(I_4 \gamma^5 \gamma^\mu \gamma^\nu), \\ &= \pm \operatorname{tr}(\gamma^\rho \gamma^\rho \gamma^5 \gamma^\mu \gamma^\nu), \\ &= (\pm 1)(-1)^3 \operatorname{tr}(\gamma^\rho \gamma^5 \gamma^\mu \gamma^\nu \gamma^\rho), \\ &= (\pm 1)(-1)^3 \operatorname{tr}(\gamma^\rho \gamma^\rho \gamma^5 \gamma^\mu \gamma^\nu), \\ &= (-1)^3 \operatorname{tr}(\gamma^5 \gamma^\mu \gamma^\nu) = -\operatorname{tr}(\gamma^5 \gamma^\mu \gamma^\nu), \end{aligned}$$

where in the third line we anticommutated γ^ρ to the right three times, and in the fourth used the cyclic property of the trace. We conclude that $\operatorname{tr}(\gamma^5 \gamma^\mu \gamma^\nu) = 0$ for all μ, ν .

$$(e) \operatorname{tr}(\gamma^5 \gamma^\mu \gamma^\nu \gamma^\rho \gamma^\sigma) = -4i\epsilon^{\mu\nu\rho\sigma}.$$

Solution: Using the anticommutation relation on the last two gamma matrices, we find

$$\begin{aligned} \operatorname{tr}(\gamma^5 \gamma^\mu \gamma^\nu \gamma^\rho \gamma^\sigma) &= \operatorname{tr}[\gamma^5 \gamma^\mu \gamma^\nu (2g^{\rho\sigma} I_4 - \gamma^\sigma \gamma^\rho)], \\ &= 2g^{\rho\sigma} \operatorname{tr}(\gamma^5 \gamma^\mu \gamma^\nu) - \operatorname{tr}(\gamma^5 \gamma^\mu \gamma^\nu \gamma^\sigma \gamma^\rho), \\ &= -\operatorname{tr}(\gamma^5 \gamma^\mu \gamma^\nu \gamma^\sigma \gamma^\rho), \end{aligned}$$

where we used the result from part d that $\operatorname{tr}(\gamma^5 \gamma^\mu \gamma^\nu) = 0$. If $\rho = \sigma$, then $(\gamma^\rho)^2 = \pm I_4$

where the ‘+’ is for $\rho = 0$ and ‘−’ otherwise. Thus, if $\rho = \sigma$, we have $\text{tr}(\gamma^5 \gamma^\mu \gamma^\nu \gamma^\rho \gamma^\sigma) \rightarrow \mp \text{tr}(\gamma^5 \gamma^\mu \gamma^\nu) = 0$. So, we conclude that $\text{tr}(\gamma^5 \gamma^\mu \gamma^\nu \gamma^\rho \gamma^\sigma)$ is antisymmetric in ρ and σ . We can repeat this argument for any pair of indices, ultimately concluding that $\text{tr}(\gamma^5 \gamma^\mu \gamma^\nu \gamma^\rho \gamma^\sigma)$ is completely antisymmetric in all μ, ν, ρ, σ indices. In 4D spacetime, the only Lorentz tensor that is completely antisymmetric is the Levi-Civita, therefore we conclude

$$\text{tr}(\gamma^5 \gamma^\mu \gamma^\nu \gamma^\rho \gamma^\sigma) = A \epsilon^{\mu\nu\rho\sigma},$$

where A is an undetermined constant and $\epsilon^{\mu\nu\rho\sigma}$ is defined such that $\epsilon^{0123} = +1$.

To determine the constant, we can take any particular combination of Lorentz indices. Let us take $(\mu, \nu, \rho, \sigma) = (0, 1, 2, 3)$, so that

$$\text{tr}(\gamma^5 \gamma^0 \gamma^1 \gamma^2 \gamma^3) = A \epsilon^{0123} = A.$$

Using the definition $\gamma^5 = i\gamma^0 \gamma^1 \gamma^2 \gamma^3$, we evaluate the trace,

$$\begin{aligned} \text{tr}(\gamma^5 \gamma^0 \gamma^1 \gamma^2 \gamma^3) &= i \text{tr}(\gamma^0 \gamma^1 \gamma^2 \gamma^3 \gamma^0 \gamma^1 \gamma^2 \gamma^3), \\ &= (-)^3 i \text{tr}(\gamma^0 \gamma^0 \gamma^1 \gamma^2 \gamma^3 \gamma^1 \gamma^2 \gamma^3), \\ &= (-1)^3 (-1)^2 i \text{tr}(\gamma^0 \gamma^0 \gamma^1 \gamma^1 \gamma^2 \gamma^3 \gamma^2 \gamma^3), \\ &= (-1)^3 (-1)^2 (-1) i \text{tr}(\gamma^0 \gamma^0 \gamma^1 \gamma^1 \gamma^2 \gamma^2 \gamma^3 \gamma^3), \\ &= (-1)^3 (-1)^2 (-1) i \text{tr}((+I_4)(-I_4)(-I_4)(-I_4)), \\ &= -i \text{tr}(I_4), \\ &= -i4, \end{aligned}$$

where in the second, third, and fourth lines we anticommutation relations to arrange identical gamma matrices into pairs, and in the fifth line we used that $(\gamma^0)^2 = +I_4$ while $(\gamma^j)^2 = -I_4$. We conclude that $A = -4i$, so that

$$\text{tr}(\gamma^5 \gamma^\mu \gamma^\nu \gamma^\rho \gamma^\sigma) = -4i \epsilon^{\mu\nu\rho\sigma}$$

3. The chiral projectors are defined as

$$P_R = \frac{1}{2}(I_4 + \gamma^5), \quad P_L = \frac{1}{2}(I_4 - \gamma^5),$$

where I_4 is the 4×4 identity matrix. Prove the following properties:

(a) $\gamma^5 P_L = -P_L$, and $\gamma^5 P_R = P_R$,

Solution: Note that $(\gamma^5)^2 = I_4$, thus,

$$\begin{aligned}\gamma^5 P_{L/R} &= \frac{1}{2} \gamma^5 (I_4 \mp \gamma^5), \\ &= \frac{1}{2} (\gamma^5 \mp (\gamma^5)^2), \\ &= \frac{1}{2} (\gamma^5 \mp I_4), \\ &= \mp \frac{1}{2} (I_4 \mp \gamma^5), \\ &= \mp P_{L/R}.\end{aligned}$$

(b) $(P_{L/R})^2 = P_{L/R}$,

Solution: Taking the square of the projectors,

$$\begin{aligned}(P_{L/R})^2 &= \left(\frac{1}{2} (I_4 \mp \gamma^5) \right)^2, \\ &= \frac{1}{4} (I_4 \mp \gamma^5)(I_4 \mp \gamma^5), \\ &= \frac{1}{4} (I_4 \mp \gamma^5 \mp \gamma^5 + (\gamma^5)^2), \\ &= \frac{1}{2} (I_4 \mp \gamma^5) = P_{L/R}\end{aligned}$$

(c) $P_L P_R = P_R P_L = 0$,

Solution: Taking product

$$\begin{aligned}P_{L/R} P_{R/L} &= \frac{1}{2} (I_4 \mp \gamma^5) \cdot \frac{1}{2} (I_4 \pm \gamma^5), \\ &= \frac{1}{2} (I_4 \mp \gamma^5 \pm \gamma^5 - (\gamma^5)^2), \\ &= \frac{1}{2} (I_4 - I_4) = 0.\end{aligned}$$

Therefore, we conclude $P_L P_R = P_R P_L = 0$.

(d) $P_L + P_R = I_4$.

Solution: Taking $P_L + P_R$, we find

$$\begin{aligned} P_L + P_R &= \frac{1}{2}(I_4 - \gamma^5) + \frac{1}{2}(I_4 + \gamma^5), \\ &= \frac{1}{2}(2I_4 - \gamma^5 + \gamma^5), \\ &= I_4. \end{aligned}$$

4. Suppose the charge conjugation operator is defined as $C = i\gamma^2\gamma^0$. Confirm that in the Weyl representation,

(a) $C^{-1} = C^\top = C^\dagger = -C$.

Solution: Given $C = i\gamma^2\gamma^0$, we first check if the matrix C is unitary, $C^\dagger C = CC^\dagger = I_4$. Note the following useful property,

$$\begin{aligned} (\gamma^\mu)^2 &= \frac{1}{2}\{\gamma^\mu, \gamma^\mu\} \text{ (no sum on } \mu), \\ &= \frac{1}{2} \cdot 2g^{\mu\mu} I_4 \text{ (no sum on } \mu), \\ &= g^{\mu\mu} I_4 \text{ (no sum on } \mu), \end{aligned}$$

so $(\gamma^0)^2 = I_4$ and $(\gamma^j)^2 = -I_4$.

$$C^\dagger = (i\gamma^2\gamma^0)^\dagger = -i(\gamma^0)^\dagger(\gamma^2)^\dagger = -i\gamma^0(-\gamma^2) = i\gamma^0\gamma^2 = -i\gamma^2\gamma^0 = -C$$

So,

$$C^\dagger C = (-i\gamma^2\gamma^0)(i\gamma^2\gamma^0) = \gamma^2\gamma^0\gamma^2\gamma^0 = -\gamma^2\gamma^0\gamma^0\gamma^2 = -\gamma^2\gamma^2 = +I_4$$

Since C is unitary, we conclude that $C^\dagger = C^{-1}$.

$$C^\top = (C^\dagger)^* = -C^* = -(i\gamma^2\gamma^0)^* = -(-i)(\gamma^2)^*(\gamma^0)^*.$$

In the Weyl basis, γ^0 is real, so $(\gamma^0)^* = \gamma^0$, and $(\gamma^2)^*$ is

$$(\gamma^2)^* = \begin{pmatrix} 0 & (\sigma^2)^* \\ -(\sigma^2)^* & 0 \end{pmatrix} = -\begin{pmatrix} 0 & \sigma^2 \\ -\sigma^2 & 0 \end{pmatrix} = -\gamma^2,$$

where we used the fact that $(\sigma^2)^* = -\sigma^2$ since the non-zero entries of σ^2 are purely imaginary. Therefore,

$$C^\top = -(-i)(\gamma^2)^*(\gamma^0)^* = i(-\gamma^2)\gamma^0 = -i\gamma^2\gamma^0 = -C.$$

We thus conclude that C is a real matrix, $C^* = C$, and that

$$C^{-1} = C^\top = C^\dagger = -C,$$

as desired.

(b) $C\gamma^\mu C^{-1} = -(\gamma^\mu)^\top$,

Solution: To prove this, first recall that $(\gamma^\mu)^\dagger = \gamma^0 \gamma^\mu \gamma^0$. Since γ^0 is real in the Weyl basis, then we conclude $(\gamma^\mu)^\top = \gamma^0 (\gamma^\mu)^* \gamma^0$, where the $*$ denote complex conjugation. Now, with $C = i\gamma^2 \gamma^0 = -i\gamma^0 \gamma^2$, with $C^{-1} = C^\dagger = -i\gamma^2 \gamma^0 = -C$, multiply $(\gamma^\mu)^\top = \gamma^0 (\gamma^\mu)^* \gamma^0$ on the left with C^{-1} and on the right with C ,

$$\begin{aligned} C^{-1}(\gamma^\mu)^\top C &= C^\dagger (\gamma^0 (\gamma^\mu)^* \gamma^0) C, \\ &= (-i\gamma^2 \gamma^0) (\gamma^0 (\gamma^\mu)^* \gamma^0) (-i\gamma^0 \gamma^2), \\ &= -\gamma^2 (\gamma^\mu)^* \gamma^2, \end{aligned}$$

where in the last line we used $(\gamma^0)^2 = I_4$. For $\mu = 0, 1, 3$, then $(\gamma^\mu)^* = \gamma^\mu$ since they are real in the Weyl basis. Then, $\gamma^\mu \gamma^2 = -\gamma^2 \gamma^\mu$ for $\mu \neq 2$. Since $(\gamma^2)^2 = -I_4$, we find $C^{-1}(\gamma^\mu)^\top C = -\gamma^2 (-\gamma^2 \gamma^\mu) = (-1)^2 (-\gamma^\mu) = -\gamma^\mu$ for $\mu \neq 2$. When $\mu = 2$, then $(\gamma^2)^* = -\gamma^2$. So, $C^{-1}(\gamma^\mu)^\top C = -\gamma^2 (-\gamma^2) \gamma^2 = -\gamma^2$. Therefore, we conclude for all μ , $C^{-1}(\gamma^\mu)^\top C = -\gamma^\mu$. Now, multiply on the left by C , and on the right by C^{-1} ,

$$CC^{-1}(\gamma^\mu)^\top CC^{-1} = -C\gamma^\mu C^{-1}, \implies C\gamma^\mu C^{-1} = -(\gamma^\mu)^\top,$$

which was to be proved.

(c) $C\gamma^5 C^{-1} = (\gamma^5)^\top$,

Solution: Here, let us take the commutator of C and γ^5 ,

$$\begin{aligned} [C, \gamma^5] &= C\gamma^5 - \gamma^5 C, \\ &= i\gamma^2 \gamma^0 \gamma^5 - i\gamma^5 \gamma^2 \gamma^0, \\ &= i\gamma^2 \gamma^0 \gamma^5 - i\gamma^2 \gamma^0 \gamma^5, \\ &= 0 \end{aligned}$$

since $\{\gamma^5, \gamma^\mu\} = 0$. Thus, $C\gamma^5 C^{-1} = \gamma^5$. Note that $(\gamma^5)^\dagger = \gamma^5$, and in the Weyl basis $\gamma^5 = (\gamma^5)^*$. Thus, we conclude $C\gamma^5 C^{-1} = (\gamma^5)^\top$.

5. A Dirac spinor ψ is called a Majorana spinor if it satisfies the condition $\psi = C\bar{\psi}^\top$, and is called a Weyl spinor if it satisfies either $\psi = P_R\psi$ or $\psi = P_L\psi$. Determine whether or not a spinor can be both Majorana and Weyl.

Solution: Let us define $\psi^c \equiv C\bar{\psi}^\top$, $\psi_L \equiv P_L\psi$, and $\psi_R \equiv P_R\psi$. We take the charge conjugation

of a chiral fermion. For example, let us take $(\psi_L)^c = C\bar{\psi}_L^\top$. Since $\bar{\psi} = \psi^\dagger\gamma^0$, we have

$$\begin{aligned} (\psi_L)^c &= C\bar{\psi}_L^\top, \\ &= C(\psi_L^\dagger\gamma^0)^\top, \\ &= C((P_L\psi)^\dagger\gamma^0)^\top, \\ &= C(\psi^\dagger P_L\gamma^0)^\top, \\ &= C(\gamma^0)^\top (P_L)^\top \psi^*, \end{aligned}$$

where we used that $\psi^\dagger = (\psi^\top)^*$ and $P_L^\dagger = P_L$ since $(\gamma^5)^\dagger = \gamma^5$. Now, we note that

$$\begin{aligned} (P_L)^\top &= \frac{1}{2}(I_4 - \gamma^5)^\top, \\ &= \frac{1}{2}(I_4 - (\gamma^5)^\top), \\ &= \frac{1}{2}(I_4 - C^{-1}\gamma^5 C), \\ &= C^{-1} \left[\frac{1}{2}(I_4 - \gamma^5) \right] C, \\ &= C^{-1} P_L C, \end{aligned}$$

where we used $C^{-1}\gamma^5 C = (\gamma^5)^\top$ and $C^{-1}C = I_4$. So, we have

$$\begin{aligned} (\psi_L)^c &= C(\gamma^0)^\top C^{-1} P_L C \psi^*, \\ &= -\gamma^0 P_L C \psi^*, \end{aligned}$$

where we used $C(\gamma^0)^\top C^{-1} = -\gamma^0$. Recall that $\gamma^0\gamma^5 = -\gamma^5\gamma^0$, so $\gamma^0 P_L = P_R\gamma^0$. Thus, we have

$$\begin{aligned} (\psi_L)^c &= -\gamma^0 P_L C \psi^*, \\ &= -P_R\gamma^0 C \psi^*. \end{aligned}$$

Finally, we use again $C^{-1}\gamma^0 C = -(\gamma^0)^\top \implies \gamma^0 C = -C(\gamma^0)^\top$, so that

$$\begin{aligned} (\psi_L)^c &= -P_R\gamma^0 C \psi^*, \\ &= P_R C(\gamma^0)^\top \psi^*, \\ &= P_R C(\psi^\dagger\gamma^0)^\top, \\ &= P_R C\bar{\psi}^\top, \\ &= P_R \psi^c, \\ &= (\psi^c)_R. \end{aligned}$$

We find that $(\psi_L)^c = (\psi^c)_R$. Similar arguments show that $(\psi_R)^c = (\psi^c)_L$. We conclude that a spinor can be both Majorana and Weyl. However, under charge conjugation the chirality flips for a Weyl fermion.

6. Consider a generic $2 \rightarrow n$ reaction $ab \rightarrow c_1 c_2 \dots c_n$ in the *lab frame* or *fixed-target frame*, that is the frame where particle b is at rest and a is the incident beam. Assume $\max(m_a, m_b) < \min(m_{c_1}, m_{c_2}, \dots, m_{c_n})$.

- (a) Show that $s = m_a^2 + m_b^2 + 2m_b\sqrt{m_a^2 + P_{\text{lab}}^2}$, where P_{lab} is the beam momentum.

Solution: By definition, $s = (p_a + p_b)^2 = m_a^2 + m_b^2 + 2p_a \cdot p_b$, where $p_a^2 = m_a^2$ and $p_b^2 = m_b^2$. Since in the lab frame, $\mathbf{p}_b = \mathbf{0}$ and $E_b = m_b$, we have $p_a \cdot p_b = E_a E_b = m_b \sqrt{m_a^2 + |\mathbf{p}_a|^2}$. Since the beam momentum is $|\mathbf{p}_a| = P_{\text{lab}}$, we find the desired result,

$$s = m_a^2 + m_b^2 + 2m_b\sqrt{m_a^2 + P_{\text{lab}}^2}.$$

- (b) Express the beam kinetic energy, $T_a \equiv E_a - m_a$, in terms of s .

Solution: From the previous result, $s = m_a^2 + m_b^2 + 2m_b E_a$ where $E_a = \sqrt{m_a^2 + P_{\text{lab}}^2}$, we have

$$E_a = \frac{s - m_a^2 - m_b^2}{2m_b},$$

in the lab frame. Then, the kinetic energy is

$$T_a = E_a - m_a = \frac{s - m_a^2 - m_b^2}{2m_b} - m_a = \frac{s - (m_a + m_b)^2}{2m_b}.$$

- (c) What is the minimum kinetic energy of the beam with which the reaction can occur?

Solution: Since $\max(m_a, m_b) < \min(m_{c_1}, m_{c_2}, \dots, m_{c_n})$, the minimum s is

$$s^{(\min)} = (m_{c_1} + m_{c_2} + \dots + m_{c_n})^2 = \left(\sum_{j=1}^n m_{c_j} \right)^2.$$

So, the minimum kinetic energy is

$$\begin{aligned} T_a^{(\min)} &= \frac{s^{(\min)} - (m_a + m_b)^2}{2m_b}, \\ &= \frac{1}{2m_b} \left[\left(\sum_{j=1}^n m_{c_j} \right)^2 - (m_a + m_b)^2 \right]. \end{aligned}$$

7. Consider the following Yukawa theory as a simplified model of an interacting proton p , neutron n , and neutral pion π^0 . We assume that the proton and neutron are distinguishable, but mass degenerate. The Lagrange density is given by

$$\mathcal{L} = \sum_f \frac{i}{2} \bar{\psi}_f \not{\partial} \psi_f + \text{h.c.} - \sum_f M \bar{\psi}_f \psi_f + \frac{1}{2} \partial_\mu \varphi \partial^\mu \varphi - \frac{1}{2} m^2 \varphi^2 - \sum_f g \varphi \bar{\psi}_f \gamma^5 \psi_f,$$

where f is fermion index $f = \{n, p\}$, M is the proton and neutron mass, m is the pion mass, and g is the coupling between proton and pion, as well as the neutron and pion.

(a) Consider the elastic reaction

$$n(p, s) + p(k, r) \rightarrow n(p', s') + p(k', r'),$$

where the arguments are the momenta and the subscripts are the spin-state. Write down the $np \rightarrow np$ scattering amplitude to leading order in the coupling g . **Hint:** Only one diagram contributes at $\mathcal{O}(g^2)$. Refer to the summary notes on **Feynman rules - Yukawa Theory**.

Solution: From the Feynman rules, we find

$$\begin{aligned} i\mathcal{M} &= \bar{u}_{s'}(p')(-ig\gamma^5)u_s(p) \frac{i}{(p' - p)^2 - m^2} \bar{u}_{r'}(k')(-ig\gamma^5)u_r(k) + \mathcal{O}(g^4), \\ &= -\frac{ig^2}{t - m^2} [\bar{u}_{s'}(p')\gamma^5 u_s(p)] [\bar{u}_{r'}(k')\gamma^5 u_r(k)] \end{aligned}$$

(b) The *spin-averaged* squared amplitude is defined as

$$\langle |\mathcal{M}|^2 \rangle \equiv \frac{1}{2} \sum_s \frac{1}{2} \sum_r \sum_{s'} \sum_{r'} \left| \mathcal{M}(n_s p_r \rightarrow n_{s'} p_{r'}) \right|^2.$$

Show that at leading order

$$\langle |\mathcal{M}|^2 \rangle = g^4 \frac{t^2}{(t - m^2)^2} + \mathcal{O}(g^6)$$

where s , t , and u are the Mandelstam invariants. **Note:** You are encouraged to use a computer algebra software such as **FeynCalc** (<https://feyncalc.github.io>), which is a **Mathematica** package for symbolic evaluation of Feynman diagrams and algebraic calculations in quantum field theory and elementary particle physics. A useful tutorial can be found [here](https://software.wm.edu). **Mathematica** is free to all students at William & Mary (see <https://software.wm.edu>).

Solution: By definition,

$$\begin{aligned} \langle |\mathcal{M}|^2 \rangle &\equiv \frac{1}{2} \sum_s \frac{1}{2} \sum_r \sum_{s'} \sum_{r'} \left| \mathcal{M} \right|^2, \\ &\equiv \frac{1}{4} \sum_{s, s'} \sum_{r, r'} \mathcal{M}^\dagger \mathcal{M} \end{aligned}$$

Now,

$$\mathcal{M}^\dagger \mathcal{M} = \frac{g^4}{(t - m^2)^2} [\bar{u}_{s'}(p')\gamma^5 u_s(p)]^\dagger [\bar{u}_{r'}(k')\gamma^5 u_r(k)]^\dagger [\bar{u}_{s'}(p')\gamma^5 u_s(p)] [\bar{u}_{r'}(k')\gamma^5 u_r(k)] + \mathcal{O}(g^6)$$

Note that

$$\begin{aligned}
 [\bar{u}_{s'} \gamma_5 u_s]^\dagger &= u_s^\dagger \gamma_5^\dagger \bar{u}_{s'}^\dagger, \\
 &= u_s^\dagger \gamma_5^\dagger (u_{s'}^\dagger \gamma^0)^\dagger, \\
 &= u_s^\dagger \gamma_5 (\gamma^0)^\dagger u_{s'}, \\
 &= u_s^\dagger \gamma_5 \gamma^0 u_{s'}, \\
 &= -u_s^\dagger \gamma^0 \gamma_5 u_{s'}, \\
 &= -\bar{u}_s \gamma_5 u_{s'},
 \end{aligned}$$

where $\gamma_5^\dagger = \gamma_5$, $(\gamma^0)^\dagger = \gamma^0$, and $\{\gamma^0, \gamma^5\} = 0$

$$\begin{aligned}
 \sum_{s,s'} \bar{u}_s(p) \gamma^5 u_{s'}(p') \bar{u}_{s'}(p') \gamma^5 u_s(p) &= \sum_{s,s'} \text{tr} \left(\bar{u}_s(p) \gamma^5 u_{s'}(p') \bar{u}_{s'}(p') \gamma^5 u_s(p) \right), \\
 &= \text{tr} \left(\gamma^5 \sum_{s'} u_{s'}(p') \bar{u}_{s'}(p') \gamma^5 \sum_s u_s(p) \bar{u}_s(p) \right), \\
 &= \text{tr} \left(\gamma^5 (\not{p}' + M) \gamma^5 (\not{p} + M) \right), \\
 &= 4(M^2 - p' \cdot p) = 2t,
 \end{aligned}$$

where $t = (p' - p)^2 = 2M^2 - 2p' \cdot p$. Similarly, we find

$$\begin{aligned}
 \sum_{r,r'} \bar{u}_r(k) \gamma^5 u_{r'}(k') \bar{u}_{r'}(k') \gamma^5 u_r(k) &= \text{tr} \left(\gamma^5 (\not{k}' + M) \gamma^5 (\not{k} + M) \right), \\
 &= 4(M^2 - k' \cdot k) = 2t.
 \end{aligned}$$

Putting all the pieces together, we find

$$\langle |\mathcal{M}|^2 \rangle = \frac{1}{4} \frac{g^4}{(t - m^2)^2} (2t)^2 + \mathcal{O}(g^6) = g^4 \frac{t^2}{(t - m^2)^2} + \mathcal{O}(g^6),$$

as desired.

- (c) Compute the unpolarized differential cross-section $d\sigma/dt$ in terms of the Mandelstam invariants.

Solution: The unpolarized differential cross section for equal mass scattering is

$$\frac{d\sigma}{d\Omega} = \frac{1}{64\pi^2 s} \langle |\mathcal{M}|^2 \rangle.$$

To find $d\sigma/dt$, note that for these kinematics $t = -2|\mathbf{p}|^2(1 - \cos\theta)$, where $|\mathbf{p}|$ is the magnitude of the relative momentum of the neutron in the CM frame. So, $dt = 2|\mathbf{p}|^2 d\cos\theta$,

and $d\Omega = 2\pi d\cos\theta$. Finally, $|\mathbf{p}|^2 = (s - 4M^2)/4$. So, the unpolarized $d\sigma/dt$ is

$$\begin{aligned}\frac{d\sigma}{dt} &= \frac{d\Omega}{dt} \frac{d\sigma}{d\Omega} = \frac{\pi}{|\mathbf{p}|^2} \frac{1}{64\pi^2 s} \langle |\mathcal{M}|^2 \rangle, \\ &= \frac{1}{16\pi s(s - 4M^2)} \langle |\mathcal{M}|^2 \rangle.\end{aligned}$$

Therefore, given that the spin-averaged matrix element squared is

$$\langle |\mathcal{M}|^2 \rangle = g^4 \frac{t^2}{(t - m^2)^2} + \mathcal{O}(g^6),$$

we have

$$\frac{d\sigma}{dt} = \frac{g^4}{16\pi} \frac{1}{s(s - 4M^2)} \frac{t^2}{(t - m^2)^2} + \mathcal{O}(g^6)$$

- (d) Express $d\sigma/d\Omega$ in terms of s and the center-of-momentum frame scattering angle θ .

Solution:

$$\begin{aligned}\frac{d\sigma}{d\Omega} &= \frac{g^4}{64\pi^2 s} \frac{t^2}{(t - m^2)^2} + \mathcal{O}(g^6), \\ &= \frac{g^4}{64\pi^2 s} \frac{4|\mathbf{p}|^4(1 - \cos\theta)^2}{(-2|\mathbf{p}|^2(1 - \cos\theta) - m^2)^2} + \mathcal{O}(g^6), \\ &= \frac{g^4}{64\pi^2 s} \left(\frac{1 - \cos\theta}{\zeta(s) - \cos\theta} \right)^2 + \mathcal{O}(g^6),\end{aligned}$$

where in the last line we defined for convenience,

$$\zeta(s) \equiv 1 + \frac{m^2}{2|\mathbf{p}|^2} = 1 + \frac{2m^2}{s - 4M^2}.$$

- (e) Compute the total cross-section as a function of s .

Solution: The total cross-section is defined as

$$\begin{aligned}\sigma &= \int d\Omega \frac{d\sigma}{d\Omega} = 2\pi \int_{-1}^{+1} d\cos\theta \frac{d\sigma}{d\Omega}, \\ &= \frac{g^4}{32\pi s} \int_{-1}^{+1} d\cos\theta \left(\frac{1 - \cos\theta}{\zeta(s) - \cos\theta} \right)^2 + \mathcal{O}(g^6).\end{aligned}$$

Note that for physical scattering, $\zeta(s) > 1$ for all $s \geq 4M^2$. So, the integral yields

$$\int_{-1}^{+1} d\cos\theta \left(\frac{1 - \cos\theta}{\zeta(s) - \cos\theta} \right)^2 = \frac{2}{\zeta(s) + 1} \left[2\zeta(s) + (\zeta^2(s) - 1) \log \left(\frac{\zeta(s) - 1}{\zeta(s) + 1} \right) \right].$$

So, the total cross-section is

$$\begin{aligned}\sigma &= \frac{g^4}{16\pi s} \frac{1}{\zeta(s) + 1} \left[2\zeta(s) + (\zeta^2(s) - 1) \log \left(\frac{\zeta(s) - 1}{\zeta(s) + 1} \right) \right], \\ &= \left(\frac{g^2}{4\pi} \right)^2 \frac{\pi}{s(\zeta(s) + 1)} \left[2\zeta(s) + (\zeta^2(s) - 1) \log \left(\frac{\zeta(s) - 1}{\zeta(s) + 1} \right) \right], \\ &= \left(\frac{g^2}{4\pi} \right)^2 \frac{\pi}{s} \left[\frac{2\zeta(s)}{\zeta(s) + 1} + (\zeta(s) - 1) \log \left(\frac{\zeta(s) - 1}{\zeta(s) + 1} \right) \right].\end{aligned}$$

- (f) Estimate the magnitude of the pion-nucleon coupling g , as well as the quantity $g^2/4\pi$, from the experimentally observed np total cross-section. **Note:** You do not need to fit the data, however feel free to do so. The Review of Particle Physics contains experimental cross-sections for select processes. See the course webpage for the data file.

Solution: The summary data from the Review of Particle Physics is given in terms of the lab frame beam momentum $P_{\text{lab.}}$, which is related to s via

$$s = 2M^2 + 2M\sqrt{M^2 + P_{\text{lab.}}^2}.$$

So, the total cross-section is given by

$$\sigma(P_{\text{lab.}}) = \left(\frac{g^2}{4\pi} \right)^2 (\hbar c)^2 \mathcal{I} \left(2M^2 + 2M\sqrt{M^2 + P_{\text{lab.}}^2} \right),$$

where $\mathcal{I}(s)$ is a distribution function


$$\mathcal{I}(s) \equiv \frac{\pi}{s} \left[\frac{2\zeta(s)}{\zeta(s) + 1} + (\zeta(s) - 1) \log \left(\frac{\zeta(s) - 1}{\zeta(s) + 1} \right) \right],$$

and $s = 2M^2 + 2M\sqrt{M^2 + P_{\text{lab.}}^2}$. The factor $(\hbar c)^2$ converts the cross-section in natural units to millibarn (mb), with $(\hbar c)^2 \approx 0.389 \text{ GeV}^2 \cdot \text{mb}$.


Note that the first inelastic threshold in this theory is π^0 production, $np \rightarrow np\pi^0$, when $s = (2M + m)^2$. Therefore, we must only include data when $(2M)^2 \leq s < (2M + m)^2$, or in terms of $P_{\text{lab.}}$,

$$\begin{aligned}P_{\text{lab.}} &< \sqrt{\left(\frac{(2M + m)^2 - 2M^2}{2M} \right)^2 - M^2}, \\ &= \frac{1}{2M} \sqrt{m(4M + m)(2M + m)^2}\end{aligned}$$

Since $M \approx 0.940 \text{ GeV}$ and $m \approx 0.140 \text{ GeV}$, then $P_{\text{lab.}} \lesssim 0.790 \text{ GeV}/c$.


 Figure 1: The function \mathcal{I} as a function of $P_{\text{lab.}}$.

We can gain some insight into this theoretical cross-section by plotting the function \mathcal{I} as a function of $P_{\text{lab.}}$. In Fig 1, we see that the function increases as $P_{\text{lab.}}$ increases. This is not the behavior of the data, which can be seen in Fig. 2, which rises dramatically as we approach threshold. Therefore, we suspect that this calculation is incomplete, either in the need for higher orders (which seems problematic from the want of a perturbative expansion if the threshold behavior is so different), or that the theory itself is incomplete (indeed, np scattering contains charge exchange, but even still this theory is not well-defined perturbatively.)


 Figure 2: np cross-section as a function of $P_{\text{lab.}}$.

Nevertheless, we can still obtain a rough estimate for the coupling can then be given by comparing the theoretical cross-section to the measured cross-section at some $P_{\text{lab.}}$.

Since the near threshold region is not captured, let's consider taking a data point near $P_{\text{lab.}}$. For example, point 440 from the RPP table, $P_{\text{lab.}} = 0.696 \text{ GeV}/c$ with $\sigma_{\text{exp.}} = (38.963 \pm 0.169) \text{ mb}$. At this momentum, $s_{\text{exp.}} = 3.97 \text{ GeV}^2$, and $\mathcal{I}(s_{\text{exp.}}) = 0.601 \text{ GeV}^{-2}$. So,

$$\begin{aligned} \left(\frac{g^2}{4\pi} \right)^2 &\approx \frac{\sigma_{\text{exp.}}}{(\hbar c)^2 \mathcal{I}(s_{\text{exp.}})} , \\ &\approx \frac{38.963 \text{ mb}}{(0.389 \text{ GeV}^2 \cdot \text{mb})(0.601 \text{ GeV}^{-2})} , \\ &\approx 166 . \end{aligned}$$

So, $g^2/4\pi \approx 13$, and $g \approx 12.8$. The coupling is extremely large, which clearly indicates that this theory does not admit a valid interpretation as a perturbation series.

FILE_NAME	REACTION	BEAM_MASS	TARGET_MASS	THRESHOLD	FINAL_STATE	MULTIPLICITY	NUMBER_OF_DATA_POINTS	POINT_NUMBER	PLAB(GEV/C)	PLAB_MIN	PLAB_MAX	SIG(MB)	STA_ERR+	STA_ERR-	SY_ER+(PCT)	SY_ER-(PCT)	REFERENCE_FLAG
FORMAT(I5,1X,4F11.5,2F8.4,1X,2F6.1,A)																	

NP_TOTAL.DAT																	
N P --> ANYTHING																	
0.939570	0.938270	0.	0	20360.	100.00	100.00	100.00	100.00	0.00531	0.00123	0.00385	0.00433	0.03043	0.03043	0.03043	0.03043	PR 76, 1744
578				6202.0	11.200	11.200	11.200	11.200	0.03043	0.03043	0.03043	0.03043	0.03043	0.03043	0.03043	0.03043	PR 129, 324
1	0.00385	0.00123	0.00531	20360.	100.00	100.00	100.00	100.00	0.00531	0.00123	0.00385	0.00433	0.03043	0.03043	0.03043	0.03043	P
2	0.03043	0.03043	0.03043	6202.0	11.200	11.200	11.200	11.200	0.03043	0.03043	0.03043	0.03043	0.03043	0.03043	0.03043	0.03043	PR 76, 188
3	0.03873	0.03873	0.03873	4700.0	40.000	40.000	40.000	40.000	0.03873	0.03873	0.03873	0.03873	0.03873	0.03873	0.03873	0.03873	P
4	0.04347	0.04347	0.04347	4228.0	18.000	18.000	18.000	18.000	0.04347	0.04347	0.04347	0.04347	0.04347	0.04347	0.04347	0.04347	PR 94, 389
5	0.04502	0.04502	0.04502	4060.0	30.000	30.000	30.000	30.000	0.04502	0.04502	0.04502	0.04502	0.04502	0.04502	0.04502	0.04502	PR 76, 188
6	0.04973	0.04973	0.04973	3675.0	16.000	16.000	16.000	16.000	0.04973	0.04973	0.04973	0.04973	0.04973	0.04973	0.04973	0.04973	PR 95, 1252
7	0.05000	0.05000	0.05000	3609.5	4.5100	4.5100	4.5100	4.5100	0.05000	0.05000	0.05000	0.05000	0.05000	0.05000	0.05000	0.05000	PR 23, 866
8	0.05020	0.05020	0.05020	3630.0	40.000	40.000	40.000	40.000	0.05020	0.05020	0.05020	0.05020	0.05020	0.05020	0.05020	0.05020	P
9	0.05100	0.05100	0.05100	3542.9	3.4400	3.4400	3.4400	3.4400	0.05100	0.05100	0.05100	0.05100	0.05100	0.05100	0.05100	0.05100	PR 23, 866
10	0.05200	0.05200	0.05200	3472.1	3.8000	3.8000	3.8000	3.8000	0.05200	0.05200	0.05200	0.05200	0.05200	0.05200	0.05200	0.05200	PR 23, 866
11	0.05300	0.05300	0.05300	3398.6	3.7400	3.7400	3.7400	3.7400	0.05300	0.05300	0.05300	0.05300	0.05300	0.05300	0.05300	0.05300	PR 23, 866
12	0.05311	0.05311	0.05311	3447.0	22.000	22.000	22.000	22.000	0.05311	0.05311	0.05311	0.05311	0.05311	0.05311	0.05311	0.05311	PR C3, 1798
13	0.05400	0.05400	0.05400	3326.5	3.6900	3.6900	3.6900	3.6900	0.05400	0.05400	0.05400	0.05400	0.05400	0.05400	0.05400	0.05400	PR 23, 866
14	0.05448	0.05448	0.05448	3330.0	20.000	20.000	20.000	20.000	0.05448	0.05448	0.05448	0.05448	0.05448	0.05448	0.05448	0.05448	P
15	0.05500	0.05500	0.05500	3266.6	4.2100	4.2100	4.2100	4.2100	0.05500	0.05500	0.05500	0.05500	0.05500	0.05500	0.05500	0.05500	PR 23, 866
16	0.05600	0.05600	0.05600	3206.6	3.6000	3.6000	3.6000	3.6000	0.05600	0.05600	0.05600	0.05600	0.05600	0.05600	0.05600	0.05600	PR 23, 866
17	0.05700	0.05700	0.05700	3138.6	4.1000	4.1000	4.1000	4.1000	0.05700	0.05700	0.05700	0.05700	0.05700	0.05700	0.05700	0.05700	PR 23, 866
18	0.05800	0.05800	0.05800	3079.6	3.5100	3.5100	3.5100	3.5100	0.05800	0.05800	0.05800	0.05800	0.05800	0.05800	0.05800	0.05800	PR 23, 866
19	0.05900	0.05900	0.05900	3015.5	3.9900	3.9900	3.9900	3.9900	0.05900	0.05900	0.05900	0.05900	0.05900	0.05900	0.05900	0.05900	PR 23, 866
20	0.05970	0.05970	0.05970	3004.0	24.000	24.000	24.000	24.000	0.05970	0.05970	0.05970	0.05970	0.05970	0.05970	0.05970	0.05970	PR C3, 1798
21	0.06000	0.06000	0.06000	2963.6	3.9500	3.9500	3.9500	3.9500	0.06000	0.06000	0.06000	0.06000	0.06000	0.06000	0.06000	0.06000	PR 23, 866
22	0.06100	0.06100	0.06100	2907.2	3.9000	3.9000	3.9000	3.9000	0.06100	0.06100	0.06100	0.06100	0.06100	0.06100	0.06100	0.06100	PR 23, 866
23	0.06200	0.06200	0.06200	2851.3	3.8600	3.8600	3.8600	3.8600	0.06200	0.06200	0.06200	0.06200	0.06200	0.06200	0.06200	0.06200	PR 23, 866
24	0.06300	0.06300	0.06300	2792.6	3.8000	3.8000	3.8000	3.8000	0.06300	0.06300	0.06300	0.06300	0.06300	0.06300	0.06300	0.06300	PR 23, 866
25	0.06400	0.06400	0.06400	2754.0	4.6200	4.6200	4.6200	4.6200	0.06400	0.06400	0.06400	0.06400	0.06400	0.06400	0.06400	0.06400	PR 23, 866
26	0.06500	0.06500	0.06500	2705.5	3.7400	3.7400	3.7400	3.7400	0.06500	0.06500	0.06500	0.06500	0.06500	0.06500	0.06500	0.06500	PR 23, 866
27	0.06571	0.06571	0.06571	2677.0	22.000	22.000	22.000	22.000	0.06571	0.06571	0.06571	0.06571	0.06571	0.06571	0.06571	0.06571	PR C3, 1798
28	0.06600	0.06600	0.06600	2651.5	3.6900	3.6900	3.6900	3.6900	0.06600	0.06600	0.06600	0.06600	0.06600	0.06600	0.06600	0.06600	PR 23, 866
29	0.06700	0.06700	0.06700	2607.4	4.4800	4.4800	4.4800	4.4800	0.06700	0.06700	0.06700	0.06700	0.06700	0.06700	0.06700	0.06700	PR 23, 866
30	0.06800	0.06800	0.06800	2558.2	3.6200	3.6200	3.6200	3.6200	0.06800	0.06800	0.06800	0.06800	0.06800	0.06800	0.06800	0.06800	PR 23, 866
...																	