

02-680 Module 15

Essentials of Mathematics and Statistics

Aidan Jan

November 18, 2025

Multiple Random Variables

Sometimes, we need to consider interactions between more than one random variable. These situations arise in many real-world and statistical applications. Consider tossing a (fair) coin 3 times, and define two random variables:

- X - the number of heads in the first toss
- Y - the number of heads in all 3 tosses

We want to know the **joint probability** (that is, the probability $p(X = x, Y = y)$)

		Y			
		0	1	2	3
X	0	$\frac{1}{8}$	$\frac{1}{4}$	$\frac{1}{8}$	0
	1	0	$\frac{1}{8}$	$\frac{1}{4}$	$\frac{1}{8}$

Marginal Probabilities

What if we're given a **joint distribution** and want to compute the probabilities of individual variables? This leads us to **marginal probabilities**:

For example, unfair coins. Let

- X = outcome of coin 1
- Y = outcome of coin 2

These are two unfair coins modeled as **random variables**. Suppose we're given the following joint probabilities:

		Y		
		Heads	Tails	
X	Heads	$\frac{1}{10}$	$\frac{2}{10}$	$\frac{3}{10}$
	Tails	$\frac{3}{10}$	$\frac{4}{10}$	$\frac{7}{10}$
		$\frac{4}{10}$	$\frac{6}{10}$	

If we want to determine, say $P(X = \text{Tails})$, it turns out we can sum over all the possibilities of Y :

$$P(X = \text{Tails}) = \sum_{y \in \{\text{Heads}, \text{Tails}\}} p(X = \text{Tails}, Y = y) = \frac{3}{10} + \frac{4}{10} = \frac{7}{10}$$

Doing all the math, both coins are biased toward Tails, the first coin (X) more-so.

What are Marginal Probabilities?

They are individual probabilities of one variable, found by summing over the other:

$$\begin{aligned}P(X = \text{Heads}) &= \frac{1}{10} + \frac{2}{10} = \frac{3}{10} \\P(X = \text{Tails}) &= \frac{3}{10} + \frac{4}{10} = \frac{7}{10} \\P(Y = \text{Heads}) &= \frac{1}{10} + \frac{3}{10} = \frac{4}{10} \\P(Y = \text{Tails}) &= \frac{2}{10} + \frac{4}{10} = \frac{6}{10}\end{aligned}$$

Continuous Random Variables

For continuous variables, we move from summation to integration:

$$p(a \leq X \leq b, c \leq Y \leq d) = \int_a^b \int_c^d f(x, y) \, dy \, dx$$

And to get a **marginal distribution function**

$$f(x) = \int_{-\infty}^{\infty} f(x, y) \, dy$$

Let's say we're drawing points uniformly over the unit square, in this case:

$$f(x, y) = \begin{cases} 1 & 0 \leq x, y \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

Therefore, if we want to find

$$p\left(X \leq \frac{1}{2}, Y \leq \frac{1}{2}\right) = \int_0^{\frac{1}{2}} \int_0^{\frac{1}{2}} 1 \, dy \, dx = \frac{1}{4}$$

Independence of Random Variables

Similar to independence for event probabilities, two random variables X and Y are **independent** if and only if

$$\forall x, y : p(X = x, Y = y) = p(X = x) \cdot p(Y = y)$$

Conditional Distributions

Conditional distribution can be derived from the joint distribution $P(X, Y)$

$$p(X|Y) = \frac{p(X, Y)}{p(Y)}$$