

# The nim game and the mathematical language

R. Ramanujam

Azim Premji University, Bengaluru  
email: [jam@imsc.res.in](mailto:jam@imsc.res.in)

Nature of Math Course, Week 5

February 1, 2022

# Nim

Nim, from the German word **nehmen**, analysed in 1904 by C. L. Bouton of Harvard University.'

- ▶ Two players I and II, move alternately.
- ▶ The game is played with  $m$  piles of counters.
- ▶ When a player moves, she picks a pile and removes some non-zero many counters from that pile.
- ▶ When a player cannot move, he loses (and the other wins).

# Ingredients

Every game has three main ingredients:

- ▶ The set of players, often  $\{I, II\}$ . In general,  $[n] = \{1, 2, \dots, n\}$ .
- ▶ The rules of the game, that specify, at any game position, whose turn it is to move, what moves are applicable, and the resulting new game position after any move.
- ▶ **Outcomes** or winning conditions, that specify at which positions the game is over, and perhaps depending on the course of play, the outcome at those positions.

# Backward induction

Zermelo 1913: In every **finite** extensive form game of perfect information, we can compute whether player  $i$  can win (or not).

- ▶ **Theorem:** Backward induction shows who wins, gives a winning strategy in the case of win / lose games, and an NE for general games.
- ▶ Note that the game arena for any Nim heap is acyclic and hence the unfolding is a finite tree, so BI applies.

# End of theory?

Backward induction completely solves finite extensive form games of perfect information, so we might as well go home.

- ▶ So the Nim game is solved, isn't it ?

# End of theory?

Backward induction completely solves finite extensive form games of perfect information, so we might as well go home.

- ▶ So the Nim game is solved, isn't it ?
- ▶ If we are only interested in **existence** of winning strategies, this suffices. If we also wish to look at the **structure** of strategies, this leaves us quite unsatisfied.
- ▶ Indeed, in the case of Nim, combinatorial analysis offers more.

## Simple cases

Consider the Nim game  $(1, 1, 2)$ , and you are to move. How would you play ?

# Simple cases

Consider the Nim game  $(1, 1, 2)$ , and you are to move. How would you play ?

- ▶ It is easy to see that removing one from the heap with 2 counters is not a good idea.



# Simple cases

Consider the Nim game  $(1, 1, 2)$ , and you are to move. How would you play ?

- ▶ It is easy to see that removing one from the heap with 2 counters is not a good idea.
- ▶ But removing all counters from that heap works.
- ▶ We can call  $(1, 1, 2)$  a winning position (for whoever plays) and  $(1, 1)$  a losing position (for whoever plays).

# Simple cases

Consider the Nim game  $(1, 1, 2)$ , and you are to move. How would you play ?

- ▶ It is easy to see that removing one from the heap with 2 counters is not a good idea.
- ▶ But removing all counters from that heap works.
- ▶ We can call  $(1, 1, 2)$  a winning position (for whoever plays) and  $(1, 1)$  a losing position (for whoever plays).
- ▶ We can see that  $(1, 1, 1)$  is a winning position.

# Simple cases

Consider the Nim game  $(1, 1, 2)$ , and you are to move. How would you play ?

- ▶ It is easy to see that removing one from the heap with 2 counters is not a good idea.
- ▶ But removing all counters from that heap works.
- ▶ We can call  $(1, 1, 2)$  a winning position (for whoever plays) and  $(1, 1)$  a losing position (for whoever plays).
- ▶ We can see that  $(1, 1, 1)$  is a winning position.
- ▶ Can we see that a  $k$ -tuple  $(1, 1, \dots, 1)$  is a winning position iff  $k$  is odd ?

# The copy strategy

Consider the case of two heaps  $(m, n)$ .

- ▶ Suppose  $m = n = 4$ , say. Now, whatever move  $I$  plays on one heap,  $II$  can **copy** that move on the other heap, thus making the heaps equal again. So this is a losing position for player  $I$ .

# The copy strategy

Consider the case of two heaps  $(m, n)$ .

- ▶ Suppose  $m = n = 4$ , say. Now, whatever move  $I$  plays on one heap,  $II$  can **copy** that move on the other heap, thus making the heaps equal again. So this is a losing position for player  $I$ .
- ▶ On the other hand, given heaps of unequal size, player  $I$  can equalize them and present  $II$  with equal heaps (which is losing for  $II$ ).

# The copy strategy

Consider the case of two heaps  $(m, n)$ .

- ▶ Suppose  $m = n = 4$ , say. Now, whatever move  $I$  plays on one heap,  $II$  can **copy** that move on the other heap, thus making the heaps equal again. So this is a losing position for player  $I$ .
- ▶ On the other hand, given heaps of unequal size, player  $I$  can equalize them and present  $II$  with equal heaps (which is losing for  $II$ ).
- ▶ **Lemma:** For all  $m, n \geq 0$ .  $(m, n)$  is winning iff  $m \neq n$ .

# Subgames

With three or more heaps, analysis becomes difficult: what can we say about  $(2, 3, 6)$ , for instance ?

# Subgames

With three or more heaps, analysis becomes difficult: what can we say about  $(2, 3, 6)$ , for instance ?

- ▶ Observe that every finite extensive form game is of the form 0 or

$$g_1 + g_2 + \dots + g_m$$

.

- ▶ 0 can be thought of as the empty game (in which no player can make any move).
- ▶  $g_1, g_2, \dots, g_m$  are subgames.



# Sum of games

Choosing between subgames has an interesting algebraic structure.

- ▶ Suppose  $g = g_1 + g_2 + \dots + g_m$ .
- ▶ Also suppose  $h = h_1 + h_2 + \dots + h_n$ .
- ▶ Then

$$g + h = (g_1 + h) + \dots + (g_m + h) + (g + h_1) + \dots + (g + h_n)$$

.

# Sum of games

Choosing between subgames has an interesting algebraic structure.

- ▶ Suppose  $g = g_1 + g_2 + \dots + g_m$ .
- ▶ Also suppose  $h = h_1 + h_2 + \dots + h_n$ .
- ▶ Then

$$g + h = (g_1 + h) + \dots + (g_m + h) + (g + h_1) + \dots + (g + h_n)$$

.

- ▶ This suggests the notation  $1 + 3 + 6$  for the nim game  $(1, 3, 6)$ .
- ▶ When  $g$  is a subgame of  $h$ , we write  $g \leq h$ .

# Equivalence of games

Once we have game expressions, a natural question arises:  
when do we consider two games  $g$  and  $g'$  to be equivalent ?

# Equivalence of games

Once we have game expressions, a natural question arises: when do we consider two games  $g$  and  $g'$  to be equivalent ? (This is only for the analysis of Nim, we will return to these considerations again later.)

- ▶ **Definition:**  $g_1 \equiv g_2$  if, for all  $h$ ,  $g_1 + h$  is winning (losing) iff  $g_2 + h$  is winning (losing).

# Equivalence of games

Once we have game expressions, a natural question arises: when do we consider two games  $g$  and  $g'$  to be equivalent ? (This is only for the analysis of Nim, we will return to these considerations again later.)

- ▶ **Definition:**  $g_1 \equiv g_2$  if, for all  $h$ ,  $g_1 + h$  is winning (losing) iff  $g_2 + h$  is winning (losing).
- ▶ Sanity check:  $\equiv$  is indeed an equivalence relation.

# Equivalence of games

Once we have game expressions, a natural question arises: when do we consider two games  $g$  and  $g'$  to be equivalent? (This is only for the analysis of Nim, we will return to these considerations again later.)

- ▶ **Definition:**  $g_1 \equiv g_2$  if, for all  $h$ ,  $g_1 + h$  is winning (losing) iff  $g_2 + h$  is winning (losing).
- ▶ Sanity check:  $\equiv$  is indeed an equivalence relation.
- ▶ Sanity check:  $g + 0 = g$ .

# Equivalence of games

Once we have game expressions, a natural question arises: when do we consider two games  $g$  and  $g'$  to be equivalent ? (This is only for the analysis of Nim, we will return to these considerations again later.)

- ▶ **Definition:**  $g_1 \equiv g_2$  if, for all  $h$ ,  $g_1 + h$  is winning (losing) iff  $g_2 + h$  is winning (losing).
- ▶ Sanity check:  $\equiv$  is indeed an equivalence relation.
- ▶ Sanity check:  $g + 0 = g$ .
- ▶ Therefore, if  $g_1 \equiv g_2$  then  $g_1$  is winning iff  $g_2$  is winning. But the converse is not true.

# Equivalence of games

Once we have game expressions, a natural question arises: when do we consider two games  $g$  and  $g'$  to be equivalent ? (This is only for the analysis of Nim, we will return to these considerations again later.)

- ▶ **Definition:**  $g_1 \equiv g_2$  if, for all  $h$ ,  $g_1 + h$  is winning (losing) iff  $g_2 + h$  is winning (losing).
- ▶ Sanity check:  $\equiv$  is indeed an equivalence relation.
- ▶ Sanity check:  $g + 0 = g$ .
- ▶ Therefore, if  $g_1 \equiv g_2$  then  $g_1$  is winning iff  $g_2$  is winning. But the converse is not true.
- ▶ However, all **losing** games are equivalent, to 0.



# The loser's lemma

**Lemma:** If  $g$  is losing then  $g \equiv 0$ .

# The loser's lemma

**Lemma:** If  $g$  is losing then  $g \equiv 0$ .

- Fix a losing game  $g$ .

# The loser's lemma

**Lemma:** If  $g$  is losing then  $g \equiv 0$ .

- ▶ Fix a losing game  $g$ .
- ▶ To prove: for all  $h$ ,  $g + h$  is losing iff  $h$  is losing.

# The loser's lemma

**Lemma:** If  $g$  is losing then  $g \equiv 0$ .

- ▶ Fix a losing game  $g$ .
- ▶ To prove: for all  $h$ ,  $g + h$  is losing iff  $h$  is losing.
- ▶ We prove by induction on  $h$  that whenever  $h$  is losing, so is  $g + h$ .

# The loser's lemma

**Lemma:** If  $g$  is losing then  $g \equiv 0$ .

- ▶ Fix a losing game  $g$ .
- ▶ To prove: for all  $h$ ,  $g + h$  is losing iff  $h$  is losing.
- ▶ We prove by induction on  $h$  that whenever  $h$  is losing, so is  $g + h$ .
- ▶ Assuming this, the other direction is easy:

# The loser's lemma

**Lemma:** If  $g$  is losing then  $g \equiv 0$ .

- ▶ Fix a losing game  $g$ .
- ▶ To prove: for all  $h$ ,  $g + h$  is losing iff  $h$  is losing.
- ▶ We prove by induction on  $h$  that whenever  $h$  is losing, so is  $g + h$ .
- ▶ Assuming this, the other direction is easy: Suppose  $h$  is winning. Then there is a move to  $h'$  that's losing.

# The loser's lemma

**Lemma:** If  $g$  is losing then  $g \equiv 0$ .

- ▶ Fix a losing game  $g$ .
- ▶ To prove: for all  $h$ ,  $g + h$  is losing iff  $h$  is losing.
- ▶ We prove by induction on  $h$  that whenever  $h$  is losing, so is  $g + h$ .
- ▶ Assuming this, the other direction is easy: Suppose  $h$  is winning. Then there is a move to  $h'$  that's losing. By claim above,  $g + h'$  is losing.

# The loser's lemma

**Lemma:** If  $g$  is losing then  $g \equiv 0$ .

- ▶ Fix a losing game  $g$ .
- ▶ To prove: for all  $h$ ,  $g + h$  is losing iff  $h$  is losing.
- ▶ We prove by induction on  $h$  that whenever  $h$  is losing, so is  $g + h$ .
- ▶ Assuming this, the other direction is easy: Suppose  $h$  is winning. Then there is a move to  $h'$  that's losing. By claim above,  $g + h'$  is losing. Hence there is a winning move in  $g + h$ , and hence  $g + h$  is winning.



# The loser's lemma

We prove by induction on  $h$  that whenever  $h$  is losing, so is  $g + h$ .

# The loser's lemma

We prove by induction on  $h$  that whenever  $h$  is losing, so is  $g + h$ .

► **IH1**: for all  $g' \leq g$ , if  $h$  is losing, then so is  $g' + h$ .

# The loser's lemma

We prove by induction on  $h$  that whenever  $h$  is losing, so is  $g + h$ .

- ▶ **IH1**: for all  $g' \leq g$ , if  $h$  is losing, then so is  $g' + h$ .
- ▶ **IH2**: for all  $h' \leq h$ , if  $h'$  is losing, then so is  $g + h'$ .

# The loser's lemma

We prove by induction on  $h$  that whenever  $h$  is losing, so is  $g + h$ .

- ▶ **IH1**: for all  $g' \leq g$ , if  $h$  is losing, then so is  $g' + h$ .
- ▶ **IH2**: for all  $h' \leq h$ , if  $h'$  is losing, then so is  $g + h'$ .
- ▶ Every initial move in  $g + h$  is either in  $g$  or in  $h$ . First consider the latter.

# The loser's lemma

We prove by induction on  $h$  that whenever  $h$  is losing, so is  $g + h$ .

- ▶ **IH1**: for all  $g' \leq g$ , if  $h$  is losing, then so is  $g' + h$ .
- ▶ **IH2**: for all  $h' \leq h$ , if  $h'$  is losing, then so is  $g + h'$ .
- ▶ Every initial move in  $g + h$  is either in  $g$  or in  $h$ . First consider the latter.
- ▶  $h$  is losing, so every move in  $h$  to  $h'$  is winning, hence there exists a move in  $h'$  to  $h''$  which is losing. By IH2,  $g + h''$  is losing.

# The loser's lemma

We prove by induction on  $h$  that whenever  $h$  is losing, so is  $g + h$ .

- ▶ **IH1**: for all  $g' \leq g$ , if  $h$  is losing, then so is  $g' + h$ .
- ▶ **IH2**: for all  $h' \leq h$ , if  $h'$  is losing, then so is  $g + h'$ .
- ▶ Every initial move in  $g + h$  is either in  $g$  or in  $h$ . First consider the latter.
- ▶  $h$  is losing, so every move in  $h$  to  $h'$  is winning, hence there exists a move in  $h'$  to  $h''$  which is losing. By IH2,  $g + h''$  is losing.
- ▶  $g$  is losing, so every move in  $g$  to  $g'$  is winning, hence there exists a move in  $g'$  to  $g''$  which is losing. By IH2,  $g'' + h$  is losing.

# The loser's lemma

We prove by induction on  $h$  that whenever  $h$  is losing, so is  $g + h$ .

- ▶ **IH1:** for all  $g' \leq g$ , if  $h$  is losing, then so is  $g' + h$ .
- ▶ **IH2:** for all  $h' \leq h$ , if  $h'$  is losing, then so is  $g + h'$ .
- ▶ Every initial move in  $g + h$  is either in  $g$  or in  $h$ . First consider the latter.
- ▶  $h$  is losing, so every move in  $h$  to  $h'$  is winning, hence there exists a move in  $h'$  to  $h''$  which is losing. By IH2,  $g + h''$  is losing.
- ▶  $g$  is losing, so every move in  $g$  to  $g'$  is winning, hence there exists a move in  $g'$  to  $g''$  which is losing. By IH2,  $g'' + h$  is losing.
- ▶ Thus every move in  $g + h$  is losing, and we are done.

# Applications

We now have: if  $h$  is losing then for all  $g$ ,  $g + h = g$ .



# Applications

We now have: if  $h$  is losing then for all  $g$ ,  $g + h = g$ .

- ▶ This means that every losing subgame can be ignored !

# Applications

We now have: if  $h$  is losing then for all  $g$ ,  $g + h = g$ .

- ▶ This means that every losing subgame can be ignored !
- ▶ Let us use this knowledge to analyse  $1 + 2 + 3 + 4 + 5$ .

# Applications

We now have: if  $h$  is losing then for all  $g$ ,  $g + h = g$ .

- ▶ This means that every losing subgame can be ignored !
- ▶ Let us use this knowledge to analyse  $1 + 2 + 3 + 4 + 5$ .
- ▶ Consider  $1 + 2 + 3$ . Note that removing an entire heap leads to a winning position. Reducing any heap leads to two equal heaps, which is losing and can be ignored. Thus  $1 + 2 + 3$  is losing.

# Applications

We now have: if  $h$  is losing then for all  $g$ ,  $g + h = g$ .

- ▶ This means that every losing subgame can be ignored !
- ▶ Let us use this knowledge to analyse  $1 + 2 + 3 + 4 + 5$ .
- ▶ Consider  $1 + 2 + 3$ . Note that removing an entire heap leads to a winning position. Reducing any heap leads to two equal heaps, which is losing and can be ignored. Thus  $1 + 2 + 3$  is losing.
- ▶ We know that  $4 + 5$  is winning, so  $1 + 2 + 3 + 4 + 5$  is winning.

# A principle

Can we get some more general mileage than analysing simple Nim heaps ?

- ▶ **Question:** How do you ensure that you do not lose in a Chess game against a Grandmaster ?

# The copycat principle

We can now enunciate an important principle in **bipartisan** games.

# The copycat principle

We can now enunciate an important principle in bipartisan games.

► **Lemma**  $g + g \equiv 0$ .

# The copycat principle

We can now enunciate an important principle in **bipartisan** games.

- ▶ **Lemma**  $g + g \equiv 0$ .
- ▶ Given  $g$ , assume that the claim holds for  $g' \leq g$ .



# The copycat principle

We can now enunciate an important principle in **bipartisan** games.

- ▶ **Lemma**  $g + g \equiv 0$ .
- ▶ Given  $g$ , assume that the claim holds for  $g' \leq g$ .
- ▶ A move in  $g + g$  is of the form  $g' + g$ .

# The copycat principle

We can now enunciate an important principle in **bipartisan** games.

- ▶ **Lemma**  $g + g \equiv 0$ .
- ▶ Given  $g$ , assume that the claim holds for  $g' \leq g$ .
- ▶ A move in  $g + g$  is of the form  $g' + g$ .  
This is winning by the move to the subgame  $g' + g'$  which is losing by IH.

# The copycat principle

We can now enunciate an important principle in **bipartisan** games.

- ▶ **Lemma**  $g + g \equiv 0$ .
- ▶ Given  $g$ , assume that the claim holds for  $g' \leq g$ .
- ▶ A move in  $g + g$  is of the form  $g' + g$ .  
This is winning by the move to the subgame  $g' + g'$  which is losing by IH.  
So  $g + g$  is losing and by loser's lemma, equivalent to 0.

# The algebraic structure

The class of games with addition forms an **abelian group**.

# The algebraic structure

The class of games with addition forms an **abelian group**.

- ▶ Commutativity and associativity are easy.

# The algebraic structure

The class of games with addition forms an **abelian group**.

- ▶ Commutativity and associativity are easy.
- ▶ 0 is the neutral element.

# The algebraic structure

The class of games with addition forms an **abelian group**.

- ▶ Commutativity and associativity are easy.
- ▶ 0 is the neutral element.
- ▶ Every element has an inverse, and as we just saw, has itself as inverse.

# The algebraic structure

The class of games with addition forms an **abelian group**.

- ▶ Commutativity and associativity are easy.
- ▶ 0 is the neutral element.
- ▶ Every element has an inverse, and as we just saw, has itself as inverse. Reminiscent of some very familiar algebraic structures?



# The algebraic structure

The class of games with addition forms an **abelian group**.

- ▶ Commutativity and associativity are easy.
- ▶ 0 is the neutral element.
- ▶ Every element has an inverse, and as we just saw, has itself as inverse. Reminiscent of some very familiar algebraic structures?

What we have seen is a glimpse of the Sprague-Grundy theory of impartial games.

# Games and numbers

John Conway took this much farther.

- ▶ There is a distinguished sub-group of games called **numbers** which can also be multiplied and which form a **field**.
- ▶ This field contains both the real numbers and the ordinal numbers.
- ▶ In fact, Conway's definition generalizes both Dedekind cuts and von Neumann ordinals.
- ▶ A beautiful microcosmos of numbers and games which are infinitesimally close to zero, and ones which are infinitely large.

# Games and numbers

John Conway took this much farther.

- ▶ There is a distinguished sub-group of games called **numbers** which can also be multiplied and which form a **field**.
- ▶ This field contains both the real numbers and the ordinal numbers.
- ▶ In fact, Conway's definition generalizes both Dedekind cuts and von Neumann ordinals.
- ▶ A beautiful microcosmos of numbers and games which are infinitesimally close to zero, and ones which are infinitely large.
- ▶ **Donald Knuth's** novel: **Surreal numbers**: every real number is surrounded by a whole lot of new numbers that lie closer to it than any other 'real' value does.