

Central limit theorems in number theory and graph theory

Kaneenika Sinha

IISc-IISER 20-20 symposium

September 18, 2021

Motivation: key ideas from a paper of Serre (1997)

Motivation: key ideas from a paper of Serre (1997)

- In 1997, Serre proved an asymptotic distribution result for the eigenvalues of the Hecke operator T_p acting on spaces of modular cusp forms of weight k and level N , $N + k \rightarrow \infty$.

Motivation: key ideas from a paper of Serre (1997)

- In 1997, Serre proved an asymptotic distribution result for the eigenvalues of the Hecke operator T_p acting on spaces of modular cusp forms of weight k and level N , $N + k \rightarrow \infty$.
- A “vertical” variant of the Sato-Tate distribution.

Motivation: key ideas from a paper of Serre (1997)

- In 1997, Serre proved an asymptotic distribution result for the eigenvalues of the Hecke operator T_p acting on spaces of modular cusp forms of weight k and level N , $N + k \rightarrow \infty$.
- A “vertical” variant of the Sato-Tate distribution.
- General principles of asymptotic distribution of families in compact intervals of \mathbb{R} .

Motivation: key ideas from a paper of Serre (1997)

- In 1997, Serre proved an asymptotic distribution result for the eigenvalues of the Hecke operator T_p acting on spaces of modular cusp forms of weight k and level N , $N + k \rightarrow \infty$.
- A “vertical” variant of the Sato-Tate distribution.
- General principles of asymptotic distribution of families in compact intervals of \mathbb{R} .
- Thematic similarities between the eigenvalues of Hecke operators and eigenvalues of families of regular graphs.

Equidistribution of families

Equidistribution of families

- Let Ω be a closed interval in \mathbb{R} with a probability measure μ .

Equidistribution of families

- Let Ω be a closed interval in \mathbb{R} with a probability measure μ . We consider families $A_V \subseteq \Omega$ such that $|A_V| \rightarrow \infty$ as $V \rightarrow \infty$. Let $C(\Omega, \mathbb{R})$ denote the set of all continuous functions $f : \Omega \rightarrow \mathbb{R}$.

Equidistribution of families

- Let Ω be a closed interval in \mathbb{R} with a probability measure μ . We consider families $A_V \subseteq \Omega$ such that $|A_V| \rightarrow \infty$ as $V \rightarrow \infty$. Let $C(\Omega, \mathbb{R})$ denote the set of all continuous functions $f : \Omega \rightarrow \mathbb{R}$.
- We say that the families $\{A_V\}$ are μ -equidistributed if

Equidistribution of families

- Let Ω be a closed interval in \mathbb{R} with a probability measure μ . We consider families $A_V \subseteq \Omega$ such that $|A_V| \rightarrow \infty$ as $V \rightarrow \infty$. Let $C(\Omega, \mathbb{R})$ denote the set of all continuous functions $f : \Omega \rightarrow \mathbb{R}$.
- We say that the families $\{A_V\}$ are μ -equidistributed if

$$\lim_{V \rightarrow \infty} \frac{1}{|A_V|} \sum_{\lambda \in A_V} f(\lambda) = \int_{\Omega} f(t) \mu(t) dt \text{ for all } f \in C(\Omega, \mathbb{R}).$$

Equidistribution of families

- Let Ω be a closed interval in \mathbb{R} with a probability measure μ . We consider families $A_V \subseteq \Omega$ such that $|A_V| \rightarrow \infty$ as $V \rightarrow \infty$. Let $C(\Omega, \mathbb{R})$ denote the set of all continuous functions $f : \Omega \rightarrow \mathbb{R}$.
- We say that the families $\{A_V\}$ are μ -equidistributed if

$$\lim_{V \rightarrow \infty} \frac{1}{|A_V|} \sum_{\lambda \in A_V} f(\lambda) = \int_{\Omega} f(t) \mu(t) dt \text{ for all } f \in C(\Omega, \mathbb{R}).$$

- Equivalently, for all intervals $I \subseteq \Omega$,

$$\lim_{V \rightarrow \infty} \frac{|A_V \cap I|}{|A_V|} = \int_I \mu(t) dt.$$

Equidistribution of families

- Let Ω be a closed interval in \mathbb{R} with a probability measure μ . We consider families $A_V \subseteq \Omega$ such that $|A_V| \rightarrow \infty$ as $V \rightarrow \infty$. Let $C(\Omega, \mathbb{R})$ denote the set of all continuous functions $f : \Omega \rightarrow \mathbb{R}$.
- We say that the families $\{A_V\}$ are μ -equidistributed if

$$\lim_{V \rightarrow \infty} \frac{1}{|A_V|} \sum_{\lambda \in A_V} f(\lambda) = \int_{\Omega} f(t) \mu(t) dt \text{ for all } f \in C(\Omega, \mathbb{R}).$$

- Equivalently, for all intervals $I \subseteq \Omega$,

$$\lim_{V \rightarrow \infty} \frac{|A_V \cap I|}{|A_V|} = \int_I \mu(t) dt.$$

- Let $\{P_n(t)\}$, $P_n(t)$ is a degree n polynomial. For all $n \geq 0$,

$$\lim_{V \rightarrow \infty} \frac{1}{|A_V|} \sum_{\lambda \in A_V} P_n(\lambda) = \int_{\Omega} P_n(t) \mu(t) dt.$$

Chebyshev polynomials

Chebyshev polynomials

- We may also choose

$$\{X_0(y), X_1(y), \dots, X_n(y), \dots\}.$$

Chebyshev polynomials

- We may also choose

$$\{X_0(y), X_1(y), \dots, X_n(y), \dots\}.$$

Here, $X_m(y)$ is the m -th Chebyshev polynomial given by

$$X_n(y) = \frac{\sin(n+1)\pi\theta}{\sin\pi\theta}, \quad y = 2\cos\pi\theta, \quad \theta \in [0, 1].$$

Chebyshev polynomials

- We may also choose

$$\{X_0(y), X_1(y), \dots, X_n(y), \dots\}.$$

Here, $X_m(y)$ is the m -th Chebyshev polynomial given by

$$X_n(y) = \frac{\sin(n+1)\pi\theta}{\sin\pi\theta}, \quad y = 2\cos\pi\theta, \quad \theta \in [0, 1].$$

- Note that

$$X_n(y) = \sum_{j=0}^{\lfloor n/2 \rfloor} (-1)^j \binom{n-j}{j} y^{n-2j}.$$

Chebyshev polynomials

- We may also choose

$$\{X_0(y), X_1(y), \dots, X_n(y), \dots\}.$$

Here, $X_m(y)$ is the m -th Chebyshev polynomial given by

$$X_n(y) = \frac{\sin(n+1)\pi\theta}{\sin\pi\theta}, \quad y = 2\cos\pi\theta, \quad \theta \in [0, 1].$$

- Note that

$$X_n(y) = \sum_{j=0}^{\lfloor n/2 \rfloor} (-1)^j \binom{n-j}{j} y^{n-2j}.$$

- We denote $Z_n(x) = X_n(x) - X_{n-2}(x)$, $n \geq 2$.

Modular cusp forms and Hecke operators

Modular cusp forms and Hecke operators

- Let k and N be positive integers with k even.

Modular cusp forms and Hecke operators

- Let k and N be positive integers with k even.
- Let $S_k(N)$ denote the space of primitive cusp forms of weight k with respect to $\Gamma_0(N)$.

Modular cusp forms and Hecke operators

- Let k and N be positive integers with k even.
- Let $S_k(N)$ denote the space of primitive cusp forms of weight k with respect to $\Gamma_0(N)$.
- Let $s_k(N)$ denote the dimension of $S_k(N)$.

Modular cusp forms and Hecke operators

- Let k and N be positive integers with k even.
- Let $S_k(N)$ denote the space of primitive cusp forms of weight k with respect to $\Gamma_0(N)$.
- Let $s_k(N)$ denote the dimension of $S_k(N)$.

$$s_k(N) \rightarrow \infty \text{ as } k + N \rightarrow \infty.$$

Modular cusp forms and Hecke operators

- Let k and N be positive integers with k even.
- Let $S_k(N)$ denote the space of primitive cusp forms of weight k with respect to $\Gamma_0(N)$.
- Let $s_k(N)$ denote the dimension of $S_k(N)$.

$$s_k(N) \rightarrow \infty \text{ as } k + N \rightarrow \infty.$$

- For $n \geq 1$, let $T_n : S_k(N) \rightarrow S_k(N)$ denote the n -th Hecke operator.

- By the Ramanujan-Deligne bound, for a prime p such that $(p, N) = 1$, the eigenvalues of T_p lie in the interval

$$\left[-2p^{(k-1)/2}, 2p^{(k-1)/2} \right].$$

- By the Ramanujan-Deligne bound, for a prime p such that $(p, N) = 1$, the eigenvalues of T_p lie in the interval

$$\left[-2p^{(k-1)/2}, 2p^{(k-1)/2}\right].$$

- We consider the normalized operator

$$T'_n := \frac{T_n}{n^{(k-1)/2}}.$$

- By the Ramanujan-Deligne bound, for a prime p such that $(p, N) = 1$, the eigenvalues of T_p lie in the interval

$$\left[-2p^{(k-1)/2}, 2p^{(k-1)/2}\right].$$

- We consider the normalized operator

$$T'_n := \frac{T_n}{n^{(k-1)/2}}.$$

The eigenvalues of T'_p lie in $[-2, 2]$.

- By the Ramanujan-Deligne bound, for a prime p such that $(p, N) = 1$, the eigenvalues of T_p lie in the interval

$$\left[-2p^{(k-1)/2}, 2p^{(k-1)/2}\right].$$

- We consider the normalized operator

$$T'_n := \frac{T_n}{n^{(k-1)/2}}.$$

The eigenvalues of T'_p lie in $[-2, 2]$.

- We denote the set of Hecke newforms in $S_k(N)$ by $\mathcal{F}_k(N)$.

- By the Ramanujan-Deligne bound, for a prime p such that $(p, N) = 1$, the eigenvalues of T_p lie in the interval

$$\left[-2p^{(k-1)/2}, 2p^{(k-1)/2}\right].$$

- We consider the normalized operator

$$T'_n := \frac{T_n}{n^{(k-1)/2}}.$$

The eigenvalues of T'_p lie in $[-2, 2]$.

- We denote the set of Hecke newforms in $S_k(N)$ by $\mathcal{F}_k(N)$.
Then $s_k(N) := |\mathcal{F}_k(N)|$.

- By the Ramanujan-Deligne bound, for a prime p such that $(p, N) = 1$, the eigenvalues of T_p lie in the interval

$$\left[-2p^{(k-1)/2}, 2p^{(k-1)/2} \right].$$

- We consider the normalized operator

$$T'_n := \frac{T_n}{n^{(k-1)/2}}.$$

The eigenvalues of T'_p lie in $[-2, 2]$.

- We denote the set of Hecke newforms in $S_k(N)$ by $\mathcal{F}_k(N)$. Then $s_k(N) := |\mathcal{F}_k(N)|$.
- Any Hecke newform $f(z) \in \mathcal{F}_k(N)$ has a Fourier expansion

$$f(z) = \sum_{n=1}^{\infty} n^{\frac{k-1}{2}} a_f(n) q^n,$$

where $a_f(1) = 1$ and

$$\frac{T_n(f(z))}{n^{\frac{k-1}{2}}} = a_f(n)f(z), \quad n \geq 1.$$

Distribution of Hecke eigenvalues

Distribution of Hecke eigenvalues

Theorem (Sarnak, 1984 and Serre, 1997)

Let $A_p(N, k)$ denote the family of eigenvalues of T'_p acting on $S_k(N)$.

Distribution of Hecke eigenvalues

Theorem (Sarnak, 1984 and Serre, 1997)

Let $A_p(N, k)$ denote the family of eigenvalues of T'_p acting on $S_k(N)$. For an interval $I \subseteq [-2, 2]$,

$$N_I(p, N, k) := |A_p(N, k) \cap I| = \sum_{\lambda \in A_p(N, k)} \chi_I(\lambda).$$

Distribution of Hecke eigenvalues

Theorem (Sarnak, 1984 and Serre, 1997)

Let $A_p(N, k)$ denote the family of eigenvalues of T'_p acting on $S_k(N)$. For an interval $I \subseteq [-2, 2]$,

$$N_I(p, N, k) := |A_p(N, k) \cap I| = \sum_{\lambda \in A_p(N, k)} \chi_I(\lambda).$$

$$\lim_{\substack{k+N \rightarrow \infty \\ p \nmid N}} \frac{N_I(p, N, k)}{s_k(N)} = \int_I \nu_p(t) dt,$$

where

$$\nu_p(t) = \frac{p+1}{\pi} \frac{\sqrt{1 - \frac{t^2}{4}}}{(p^{1/2} + p^{-1/2})^2 - t^2}.$$

- We choose an interval $I \subset [-2, 2]$.

- We choose an interval $I \subset [-2, 2]$.
- Denote $\mu(I) := \int_I \mu(t) dt = \frac{1}{\pi} \int_I \sqrt{1 - \frac{t^2}{4}} dt$.

- We choose an interval $I \subset [-2, 2]$.
- Denote $\mu(I) := \int_I \mu(t) dt = \frac{1}{\pi} \int_I \sqrt{1 - \frac{t^2}{4}} dt$.
- Define $Y_p : \mathcal{F}_k(N) \rightarrow \{0, 1\}$ as

$$Y_p(f) = \chi_I(a_f(p)).$$

- We choose an interval $I \subset [-2, 2]$.
- Denote $\mu(I) := \int_I \mu(t) dt = \frac{1}{\pi} \int_I \sqrt{1 - \frac{t^2}{4}} dt$.
- Define $Y_p : \mathcal{F}_k(N) \rightarrow \{0, 1\}$ as

$$Y_p(f) = \chi_I(a_f(p)).$$

- $\mu_p = \mathbb{E}[Y_p] = \frac{1}{s_k(N)} \sum_{f \in \mathcal{F}_k(N)} \chi_I(a_f(p)).$

- We choose an interval $I \subset [-2, 2]$.
- Denote $\mu(I) := \int_I \mu(t) dt = \frac{1}{\pi} \int_I \sqrt{1 - \frac{t^2}{4}} dt$.
- Define $Y_p : \mathcal{F}_k(N) \rightarrow \{0, 1\}$ as

$$Y_p(f) = \chi_I(a_f(p)).$$

- $\mu_p = \mathbb{E}[Y_p] = \frac{1}{s_k(N)} \sum_{f \in \mathcal{F}_k(N)} \chi_I(a_f(p))$.
- $\sigma_p^2 = \frac{1}{s_k(N)} \sum_{f \in \mathcal{F}_k(N)} (Y_p(f) - \mu_p)^2$.

- We choose an interval $I \subset [-2, 2]$.
- Denote $\mu(I) := \int_I \mu(t) dt = \frac{1}{\pi} \int_I \sqrt{1 - \frac{t^2}{4}} dt$.
- Define $Y_p : \mathcal{F}_k(N) \rightarrow \{0, 1\}$ as

$$Y_p(f) = \chi_I(a_f(p)).$$

- $\mu_p = \mathbb{E}[Y_p] = \frac{1}{s_k(N)} \sum_{f \in \mathcal{F}_k(N)} \chi_I(a_f(p))$.
- $\sigma_p^2 = \frac{1}{s_k(N)} \sum_{f \in \mathcal{F}_k(N)} (Y_p(f) - \mu_p)^2$.
- Is there a central limit-type theorem that predicts the distribution of

$$\frac{\sum_{p \leq x} (Y_p - \mu_p)}{\sqrt{\sum_{p \leq x} \sigma_p^2}}.$$

Theorem (Conrey, Duke Farmer, 1997, Nagoshi, 2006)

Let $k = k(x)$ such that $\frac{\log k}{\log x} \rightarrow \infty$ as $x \rightarrow \infty$.

Theorem (Conrey, Duke Farmer, 1997, Nagoshi, 2006)

Let $k = k(x)$ such that $\frac{\log k}{\log x} \rightarrow \infty$ as $x \rightarrow \infty$. Then, as $x \rightarrow \infty$,

$$\sum_{p \leq x} \mu_p := \sum_{p \leq x} \frac{1}{s_k(N)} \sum_{f \in \mathcal{F}_k(N)} Y_p(f) \sim \pi(x) \mu(I).$$

Theorem (Prabhu, S, 2017)

Let $k = k(x)$ such that $\frac{\log k}{\sqrt{x} \log x} \rightarrow \infty$ as $x \rightarrow \infty$.

Theorem (Conrey, Duke Farmer, 1997, Nagoshi, 2006)

Let $k = k(x)$ such that $\frac{\log k}{\log x} \rightarrow \infty$ as $x \rightarrow \infty$. Then, as $x \rightarrow \infty$,

$$\sum_{p \leq x} \mu_p := \sum_{p \leq x} \frac{1}{s_k(N)} \sum_{f \in \mathcal{F}_k(N)} Y_p(f) \sim \pi(x) \mu(I).$$

Theorem (Prabhu, S, 2017)

Let $k = k(x)$ such that $\frac{\log k}{\sqrt{x} \log x} \rightarrow \infty$ as $x \rightarrow \infty$.

$$\sum_{p \leq x} \sigma_p^2 := \sum_{p \leq x} \frac{1}{s_k(N)} \sum_{f \in \mathcal{F}_k(N)} (Y_p(f) - \mu_p)^2 \sim \pi(x) [\mu(I) - \mu(I)^2].$$

Theorem (Ctd.)

For any integer $r \geq 0$,

$$\begin{aligned} \lim_{x \rightarrow \infty} \frac{1}{s_k(1)} \sum_{f \in \mathcal{F}_k(1)} \left(\frac{\sum_{p \leq x} \chi_I(a_f(p)) - \pi(x)\mu(I)}{\sqrt{\pi(x)[\mu(I) - \mu(I)^2]}} \right)^r \\ = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} t^r e^{-t^2/2} dt. \end{aligned}$$

Theorem (Ctd.)

For any integer $r \geq 0$,

$$\begin{aligned} \lim_{x \rightarrow \infty} \frac{1}{s_k(1)} \sum_{f \in \mathcal{F}_k(1)} \left(\frac{\sum_{p \leq x} \chi_I(a_f(p)) - \pi(x)\mu(I)}{\sqrt{\pi(x)[\mu(I) - \mu(I)^2]}} \right)^r \\ = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} t^r e^{-t^2/2} dt. \end{aligned}$$

Theorem (Ctd.)

For any integer $r \geq 0$,

$$\begin{aligned} \lim_{x \rightarrow \infty} \frac{1}{s_k(1)} \sum_{f \in \mathcal{F}_k(1)} \left(\frac{\sum_{p \leq x} \chi_I(a_f(p)) - \pi(x)\mu(I)}{\sqrt{\pi(x)[\mu(I) - \mu(I)^2]}} \right)^r \\ = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} t^r e^{-t^2/2} dt. \end{aligned}$$

That is, $\frac{\sum_{p \leq x} (Y_p - \mu_p)}{\sqrt{\sum_{p \leq x} \sigma_p^2}}$ converges to the normal distribution as $x \rightarrow \infty$.

Theorem (Ctd.)

For any integer $r \geq 0$,

$$\lim_{x \rightarrow \infty} \frac{1}{s_k(1)} \sum_{f \in \mathcal{F}_k(1)} \left(\frac{\sum_{p \leq x} \chi_I(a_f(p)) - \pi(x)\mu(I)}{\sqrt{\pi(x)[\mu(I) - \mu(I)^2]}} \right)^r \\ = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} t^r e^{-t^2/2} dt.$$

That is, $\frac{\sum_{p \leq x} (Y_p - \mu_p)}{\sqrt{\sum_{p \leq x} \sigma_p^2}}$ converges to the normal distribution as $x \rightarrow \infty$. Our motivation was a result of Nagoshi, which showed that

$$\frac{\sum_{p \leq x} a_f(p)}{\sqrt{\pi(x)}}$$

converges to the normal distribution as $x \rightarrow \infty$ (as $\log k / \log x \rightarrow \infty$).

Theorem (Baier-Prabhu-S, 2019)

Let $G(t) = \sum_{m \geq 0} U(m) X_{2m}(t)$, $|U(m)| \ll e^{-\alpha m^\omega}$. Define $Y_p : \mathcal{F}_k(1) \rightarrow \mathbb{R}$ given by

$$Y_p(f) = G(a_f(p)).$$

Theorem (Baier-Prabhu-S, 2019)

Let $G(t) = \sum_{m \geq 0} U(m) X_{2m}(t)$, $|U(m)| \ll e^{-\alpha m^\omega}$. Define $Y_p : \mathcal{F}_k(1) \rightarrow \mathbb{R}$ given by

$$Y_p(f) = G(a_f(p)).$$

Let $k = k(x) \geq 2$ such that

$$\frac{(\log k)}{(\log x)^{1+1/\omega}} \rightarrow \infty \text{ as } x \rightarrow \infty.$$

Theorem (Baier-Prabhu-S, 2019)

Let $G(t) = \sum_{m \geq 0} U(m) X_{2m}(t)$, $|U(m)| \ll e^{-\alpha m^\omega}$. Define $Y_p : \mathcal{F}_k(1) \rightarrow \mathbb{R}$ given by

$$Y_p(f) = G(a_f(p)).$$

Let $k = k(x) \geq 2$ such that

$$\frac{(\log k)}{(\log x)^{1+1/\omega}} \rightarrow \infty \text{ as } x \rightarrow \infty.$$

Then, as $x \rightarrow \infty$,

- $\sum_{p \leq x} \mathbb{E}[Y_p] \sim \pi(x) \int_0^1 G(t) \mu(t) dt,$

Theorem (Baier-Prabhu-S, 2019)

Let $G(t) = \sum_{m \geq 0} U(m) X_{2m}(t)$, $|U(m)| \ll e^{-\alpha m^\omega}$. Define $Y_p : \mathcal{F}_k(1) \rightarrow \mathbb{R}$ given by

$$Y_p(f) = G(a_f(p)).$$

Let $k = k(x) \geq 2$ such that

$$\frac{(\log k)}{(\log x)^{1+1/\omega}} \rightarrow \infty \text{ as } x \rightarrow \infty.$$

Then, as $x \rightarrow \infty$,

- $\sum_{p \leq x} \mathbb{E}[Y_p] \sim \pi(x) \int_0^1 G(t) \mu(t) dt$,
- $\sum_{p \leq x} \sigma_p^2 \sim \pi(x) V_G$, where

$$V_G = \int_0^1 G(t)^2 \mu(t) dt - \left(\int_0^1 G(t) \mu(t) dt \right)^2.$$

Theorem (Contd.)

- For any integer $r \geq 0$,

$$\frac{1}{s_k(1)} \sum_{f \in \mathcal{F}_k(1)} \left(\frac{\sum_{p \leq x} Y_p(f) - \pi(x) \int_0^1 G(t) \mu(t) dt}{\sqrt{\pi(x) V_G}} \right)^r \\ \sim \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} t^r e^{-t^2/2} dt.$$

Eigenvalues of regular graphs

Eigenvalues of regular graphs

- Let $\mathcal{G}(m, d)$ be the collection of d -regular, simple, undirected graphs on m vertices ($d \geq 3$).

Eigenvalues of regular graphs

- Let $\mathcal{G}(m, d)$ be the collection of d -regular, simple, undirected graphs on m vertices ($d \geq 3$).
- Suppose we label the vertices of $X \in \mathcal{G}(m, d)$ as v_1, v_2, \dots, v_m . Its adjacency matrix $\mathcal{A} = \mathcal{A}(X) \in \{0, 1\}^{m \times m}$ is defined by setting $\mathcal{A}_{ij} = 1$ if there is an edge $(v_i, v_j) \in \mathcal{E}$, and $\mathcal{A}_{ij} = 0$ otherwise.

Eigenvalues of regular graphs

- Let $\mathcal{G}(m, d)$ be the collection of d -regular, simple, undirected graphs on m vertices ($d \geq 3$).
- Suppose we label the vertices of $X \in \mathcal{G}(m, d)$ as v_1, v_2, \dots, v_m . Its adjacency matrix $\mathcal{A} = \mathcal{A}(X) \in \{0, 1\}^{m \times m}$ is defined by setting $\mathcal{A}_{ij} = 1$ if there is an edge $(v_i, v_j) \in \mathcal{E}$, and $\mathcal{A}_{ij} = 0$ otherwise.
- \mathcal{A} is symmetric and therefore has only real eigenvalues. These eigenvalues are called the eigenvalues of X .

Eigenvalues of regular graphs

- Let $\mathcal{G}(m, d)$ be the collection of d -regular, simple, undirected graphs on m vertices ($d \geq 3$).
- Suppose we label the vertices of $X \in \mathcal{G}(m, d)$ as v_1, v_2, \dots, v_m . Its adjacency matrix $\mathcal{A} = \mathcal{A}(X) \in \{0, 1\}^{m \times m}$ is defined by setting $\mathcal{A}_{ij} = 1$ if there is an edge $(v_i, v_j) \in \mathcal{E}$, and $\mathcal{A}_{ij} = 0$ otherwise.
- \mathcal{A} is symmetric and therefore has only real eigenvalues. These eigenvalues are called the eigenvalues of X .
- Any eigenvalue λ_i of X with $|\lambda_i| = d$ is called a trivial eigenvalue of X . We define

$$\lambda(X) := \max_{\lambda_i(X) \neq \pm d} |\lambda_i(X)|.$$

Eigenvalues of regular graphs

- Let $\mathcal{G}(m, d)$ be the collection of d -regular, simple, undirected graphs on m vertices ($d \geq 3$).
- Suppose we label the vertices of $X \in \mathcal{G}(m, d)$ as v_1, v_2, \dots, v_m . Its adjacency matrix $\mathcal{A} = \mathcal{A}(X) \in \{0, 1\}^{m \times m}$ is defined by setting $\mathcal{A}_{ij} = 1$ if there is an edge $(v_i, v_j) \in \mathcal{E}$, and $\mathcal{A}_{ij} = 0$ otherwise.
- \mathcal{A} is symmetric and therefore has only real eigenvalues. These eigenvalues are called the eigenvalues of X .
- Any eigenvalue λ_i of X with $|\lambda_i| = d$ is called a trivial eigenvalue of X . We define

$$\lambda(X) := \max_{\lambda_i(X) \neq \pm d} |\lambda_i(X)|.$$

- Let $A(X)$ denote the family of eigenvalues of $\frac{1}{\sqrt{d-1}}\mathcal{A}$. Clearly, $A(X)$ is contained in the interval $\left[-\frac{d}{\sqrt{d-1}}, \frac{d}{\sqrt{d-1}}\right]$.

- Let $A(X)$ denote the family of eigenvalues of $\frac{1}{\sqrt{d-1}}\mathcal{A}$. Clearly, $A(X)$ is contained in the interval $\left[-\frac{d}{\sqrt{d-1}}, \frac{d}{\sqrt{d-1}}\right]$.
- Let $\Lambda(x) := \frac{\lambda(X)}{\sqrt{d-1}}$.

- Let $A(X)$ denote the family of eigenvalues of $\frac{1}{\sqrt{d-1}}\mathcal{A}$. Clearly, $A(X)$ is contained in the interval $\left[-\frac{d}{\sqrt{d-1}}, \frac{d}{\sqrt{d-1}}\right]$.
- Let $\Lambda(x) := \frac{\lambda(x)}{\sqrt{d-1}}$.
- Let $d \geq 3$ be fixed. By a 2008 result of Friedman, for X chosen uniformly at random from $\mathcal{G}(m, d)$ and for any $\epsilon > 0$,

$$P[\Lambda(X) > 2 + \epsilon] \rightarrow 0 \text{ as } m \rightarrow \infty.$$

- Let $A(X)$ denote the family of eigenvalues of $\frac{1}{\sqrt{d-1}}\mathcal{A}$. Clearly, $A(X)$ is contained in the interval $\left[-\frac{d}{\sqrt{d-1}}, \frac{d}{\sqrt{d-1}}\right]$.
- Let $\Lambda(x) := \frac{\lambda(x)}{\sqrt{d-1}}$.
- Let $d \geq 3$ be fixed. By a 2008 result of Friedman, for X chosen uniformly at random from $\mathcal{G}(m, d)$ and for any $\epsilon > 0$,

$$P[\Lambda(X) > 2 + \epsilon] \rightarrow 0 \text{ as } m \rightarrow \infty.$$

- Let us choose a sequence $\{X_m\}$ of graphs such that $X_m \in \mathcal{G}(m, d)$. How are the families $A(X_m)$ distributed as $m \rightarrow \infty$?

- Let $A(X)$ denote the family of eigenvalues of $\frac{1}{\sqrt{d-1}}\mathcal{A}$. Clearly, $A(X)$ is contained in the interval $\left[-\frac{d}{\sqrt{d-1}}, \frac{d}{\sqrt{d-1}}\right]$.
- Let $\Lambda(x) := \frac{\lambda(x)}{\sqrt{d-1}}$.
- Let $d \geq 3$ be fixed. By a 2008 result of Friedman, for X chosen uniformly at random from $\mathcal{G}(m, d)$ and for any $\epsilon > 0$,

$$P[\Lambda(X) > 2 + \epsilon] \rightarrow 0 \text{ as } m \rightarrow \infty.$$

- Let us choose a sequence $\{X_m\}$ of graphs such that $X_m \in \mathcal{G}(m, d)$. How are the families $A(X_m)$ distributed as $m \rightarrow \infty$?
- Let $d = d(m) \rightarrow \infty$ as $m \rightarrow \infty$. How are the families $A(X_m)$ distributed as $m \rightarrow \infty$?

- Let $A(X)$ denote the family of eigenvalues of $\frac{1}{\sqrt{d-1}}\mathcal{A}$. Clearly, $A(X)$ is contained in the interval $\left[-\frac{d}{\sqrt{d-1}}, \frac{d}{\sqrt{d-1}}\right]$.
- Let $\Lambda(x) := \frac{\lambda(x)}{\sqrt{d-1}}$.
- Let $d \geq 3$ be fixed. By a 2008 result of Friedman, for X chosen uniformly at random from $\mathcal{G}(m, d)$ and for any $\epsilon > 0$,

$$P[\Lambda(X) > 2 + \epsilon] \rightarrow 0 \text{ as } m \rightarrow \infty.$$

- Let us choose a sequence $\{X_m\}$ of graphs such that $X_m \in \mathcal{G}(m, d)$. How are the families $A(X_m)$ distributed as $m \rightarrow \infty$?
- Let $d = d(m) \rightarrow \infty$ as $m \rightarrow \infty$. How are the families $A(X_m)$ distributed as $m \rightarrow \infty$?

Theorem (McKay, 1980)

Let $d \geq 3$ be fixed and let $X_m \in \mathcal{G}(m, d)$ such that $\frac{c_n(X_m)}{m} \rightarrow 0$ as $m \rightarrow \infty$.

Theorem (McKay, 1980)

Let $d \geq 3$ be fixed and let $X_m \in \mathcal{G}(m, d)$ such that $\frac{c_n(X_m)}{m} \rightarrow 0$ as $m \rightarrow \infty$. Here, c_n denotes the number of cycles of length n . Then, for any interval $I \subset [-2, 2]$,

$$\lim_{m \rightarrow \infty} \frac{|A(X) \cap I|}{m} = \int_I \kappa_d(t) dt,$$

Theorem (McKay, 1980)

Let $d \geq 3$ be fixed and let $X_m \in \mathcal{G}(m, d)$ such that $\frac{c_n(X_m)}{m} \rightarrow 0$ as $m \rightarrow \infty$. Here, c_n denotes the number of cycles of length n . Then, for any interval $I \subset [-2, 2]$,

$$\lim_{m \rightarrow \infty} \frac{|A(X) \cap I|}{m} = \int_I \kappa_d(t) dt,$$

$$\kappa_d(t) = \frac{d}{\pi \left(\frac{d^2}{d-1} - t^2 \right)} \sqrt{1 - \frac{t^2}{4}}.$$

Theorem (McKay, 1980)

Let $d \geq 3$ be fixed and let $X_m \in \mathcal{G}(m, d)$ such that $\frac{c_n(X_m)}{m} \rightarrow 0$ as $m \rightarrow \infty$. Here, c_n denotes the number of cycles of length n . Then, for any interval $I \subset [-2, 2]$,

$$\lim_{m \rightarrow \infty} \frac{|A(X) \cap I|}{m} = \int_I \kappa_d(t) dt,$$

$$\kappa_d(t) = \frac{d}{\pi \left(\frac{d^2}{d-1} - t^2 \right)} \sqrt{1 - \frac{t^2}{4}}.$$

Equivalently, for any continuous function

$$f : \left[-\frac{d}{\sqrt{d-1}}, \frac{d}{\sqrt{d-1}} \right] \rightarrow \mathbb{R},$$

Theorem (McKay, 1980)

Let $d \geq 3$ be fixed and let $X_m \in \mathcal{G}(m, d)$ such that $\frac{c_n(X_m)}{m} \rightarrow 0$ as $m \rightarrow \infty$. Here, c_n denotes the number of cycles of length n . Then, for any interval $I \subset [-2, 2]$,

$$\lim_{m \rightarrow \infty} \frac{|A(X) \cap I|}{m} = \int_I \kappa_d(t) dt,$$

$$\kappa_d(t) = \frac{d}{\pi \left(\frac{d^2}{d-1} - t^2 \right)} \sqrt{1 - \frac{t^2}{4}}.$$

Equivalently, for any continuous function

$$f : \left[-\frac{d}{\sqrt{d-1}}, \frac{d}{\sqrt{d-1}} \right] \rightarrow \mathbb{R},$$

$$\lim_{m \rightarrow \infty} \frac{1}{m} \sum_{\lambda \in A(X)} f(\lambda) = \int_{-2}^2 f(t) \kappa_d(t) dt.$$

Theorem (McKay, 1980)

Let $d \geq 3$ be fixed and let $X_m \in \mathcal{G}(m, d)$ such that $\frac{c_n(X_m)}{m} \rightarrow 0$ as $m \rightarrow \infty$. Here, c_n denotes the number of cycles of length n . Then, for any interval $I \subset [-2, 2]$,

$$\lim_{m \rightarrow \infty} \frac{|A(X) \cap I|}{m} = \int_I \kappa_d(t) dt,$$

$$\kappa_d(t) = \frac{d}{\pi \left(\frac{d^2}{d-1} - t^2 \right)} \sqrt{1 - \frac{t^2}{4}}.$$

Equivalently, for any continuous function

$$f : \left[-\frac{d}{\sqrt{d-1}}, \frac{d}{\sqrt{d-1}} \right] \rightarrow \mathbb{R},$$

$$\lim_{m \rightarrow \infty} \frac{1}{m} \sum_{\lambda \in A(X)} f(\lambda) = \int_{-2}^2 f(t) \kappa_d(t) dt.$$

Note: if $d = p + 1$, we recover $\nu_p(t)$.

Eigenvalue statistics

Eigenvalue statistics

- For some function $f : \mathbb{R} \rightarrow \mathbb{R}$, we may define a random variable $B_f : \mathcal{G}(m, d) \rightarrow \mathbb{R}$ by

$$B_f(Y) := \sum_{\lambda \in A(Y)} f(\lambda).$$

Eigenvalue statistics

- For some function $f : \mathbb{R} \rightarrow \mathbb{R}$, we may define a random variable $B_f : \mathcal{G}(m, d) \rightarrow \mathbb{R}$ by

$$B_f(Y) := \sum_{\lambda \in A(Y)} f(\lambda).$$

- A theorem of Dumitriu and Pal implies that if $f \in C(\mathbb{R})$, and $d(m) \rightarrow \infty$, $d_m - 1 = m^{\epsilon_m}$ for some $\epsilon_m = o(1)$,

then

$$\mathbb{E}[B_f] := \frac{1}{|\mathcal{G}(m, d(m))|} \sum_{Y \in \mathcal{G}(m, d(m))} B_f(Y) \sim m \int_{-2}^2 f(t) \mu(t) dt.$$

Eigenvalue statistics

- For some function $f : \mathbb{R} \rightarrow \mathbb{R}$, we may define a random variable $B_f : \mathcal{G}(m, d) \rightarrow \mathbb{R}$ by

$$B_f(Y) := \sum_{\lambda \in A(Y)} f(\lambda).$$

- A theorem of Dumitriu and Pal implies that if $f \in C(\mathbb{R})$, and

$$d(m) \rightarrow \infty, \quad d_m - 1 = m^{\epsilon_m} \text{ for some } \epsilon_m = o(1),$$

then

$$\mathbb{E}[B_f] := \frac{1}{|\mathcal{G}(m, d(m))|} \sum_{Y \in \mathcal{G}(m, d(m))} B_f(Y) \sim m \int_{-2}^2 f(t) \mu(t) dt.$$

- Moreover, for an interval $I \subset [-2, 2]$,

$$\mathbb{E}[B_{\chi_I}] \sim m \int_I \mu(t).$$

Central limit theorem

Central limit theorem

Theorem (Johnson, 2015)

Suppose f is an entire function of finite order. In this case, f can be expressed as

$$f(x) = \sum_{k=0}^{\infty} a_k X_k(x).$$

Central limit theorem

Theorem (Johnson, 2015)

Suppose f is an entire function of finite order. In this case, f can be expressed as

$$f(x) = \sum_{k=0}^{\infty} a_k X_k(x).$$

Let $d = d(m) \rightarrow \infty$ such that

$$\frac{\log \log m}{\log d} \rightarrow \infty \text{ as } m \rightarrow \infty.$$

Central limit theorem

Theorem (Johnson, 2015)

Suppose f is an entire function of finite order. In this case, f can be expressed as

$$f(x) = \sum_{k=0}^{\infty} a_k X_k(x).$$

Let $d = d(m) \rightarrow \infty$ such that

$$\frac{\log \log m}{\log d} \rightarrow \infty \text{ as } m \rightarrow \infty.$$

Then,

$$V[B_f] := \mathbb{E} \left[(B_f - \mathbb{E}[B_f])^2 \right] \sim m \frac{1}{2} \sum_{k=3}^{\infty} k a_k^2 \text{ as } m \rightarrow \infty.$$

Theorem (Ctd.)

For every $r \geq 0$,

$$\lim_{m \rightarrow \infty} \frac{1}{|\mathcal{G}(m, d(m))|} \sum_{Y \in \mathcal{G}(m, d(m))} \left(\frac{\left(B_f(Y) - m \int_{-2}^2 f(t) \mu(t) dt \right)}{\sqrt{V[B_f]}} \right)^r$$

Theorem (Ctd.)

For every $r \geq 0$,

$$\lim_{m \rightarrow \infty} \frac{1}{|\mathcal{G}(m, d(m))|} \sum_{Y \in \mathcal{G}(m, d(m))} \left(\frac{\left(B_f(Y) - m \int_{-2}^2 f(t) \mu(t) dt \right)}{\sqrt{V[B_f]}} \right)^r \\ = \begin{cases} 0 & \text{if } r \text{ is odd} \\ \frac{r!}{(r/2)! 2^{r/2}} & \text{if } r \text{ is even.} \end{cases}$$

Can we obtain asymptotics $\mathbb{E}[B_f]$, $V[B_f]$ and a central limit theorem under “looser” growth conditions for d in terms of m (possibly for a restricted class of test functions)?

Smooth analogue (Joint work with M. Podder)

- One takes a real-valued, even test function $g \in C^\infty(\mathbb{R})$.

Smooth analogue (Joint work with M. Podder)

- One takes a real-valued, even test function $g \in C^\infty(\mathbb{R})$.
- One then considers, for $L \geq 1$,

$$G_L(\theta) := \sum_{m \in \mathbb{Z}} g(L(\theta + m)).$$

Smooth analogue (Joint work with M. Podder)

- One takes a real-valued, even test function $g \in C^\infty(\mathbb{R})$.
- One then considers, for $L \geq 1$,

$$G_L(\theta) := \sum_{m \in \mathbb{Z}} g(L(\theta + m)).$$

- Suppose the Fourier transform \widehat{g} has compact support, without loss of generality, in $[-1, 1]$.

Smooth analogue (Joint work with M. Podder)

- One takes a real-valued, even test function $g \in C^\infty(\mathbb{R})$.
- One then considers, for $L \geq 1$,

$$G_L(\theta) := \sum_{m \in \mathbb{Z}} g(L(\theta + m)).$$

- Suppose the Fourier transform \hat{g} has compact support, without loss of generality, in $[-1, 1]$.
- $G_L(\theta)$ is a periodic function with Fourier expansion

$$G_L(\theta) = \frac{1}{L} \sum_{|m| \leq L} \hat{g}\left(\frac{m}{L}\right) e(m\theta), \quad e(x) = e^{2\pi i x}.$$

- In fact,

$$H_L(t) := G_L(\theta) = \sum_{0 \leq m \leq L} U(m) X_{2m}(t), \quad t = 2 \cos \pi \theta, \text{ where}$$

$$U(m) = \frac{1}{L} \left(\hat{g}\left(\frac{m}{L}\right) - \hat{g}\left(\frac{m+1}{L}\right) \right).$$

Theorem (Podder, S)

Let $L \geq 1$. Define $B_{H_L} : \mathcal{G}(m, d) \rightarrow \mathbb{R}$ by

$$B_{H_L}(Y) := \sum_{\lambda \in A(Y)} H_L(\lambda).$$

Theorem (Podder, S)

Let $L \geq 1$. Define $B_{H_L} : \mathcal{G}(m, d) \rightarrow \mathbb{R}$ by

$$B_{H_L}(Y) := \sum_{\lambda \in A(Y)} H_L(\lambda).$$

Let $d = d(m) \rightarrow \infty$ such that

$$\frac{\log m}{\log d} \rightarrow \infty \text{ as } m \rightarrow \infty.$$

Theorem (Podder, S)

Let $L \geq 1$. Define $B_{H_L} : \mathcal{G}(m, d) \rightarrow \mathbb{R}$ by

$$B_{H_L}(Y) := \sum_{\lambda \in A(Y)} H_L(\lambda).$$

Let $d = d(m) \rightarrow \infty$ such that

$$\frac{\log m}{\log d} \rightarrow \infty \text{ as } m \rightarrow \infty.$$

Then,

$$\mathbb{E}[B_{H_L}] \sim m \int_{-2}^2 H_L(t) \mu(t) dt, \text{ as } m \rightarrow \infty.$$

Theorem (Podder, S)

Let $L \geq 1$. Define $B_{H_L} : \mathcal{G}(m, d) \rightarrow \mathbb{R}$ by

$$B_{H_L}(Y) := \sum_{\lambda \in A(Y)} H_L(\lambda).$$

Let $d = d(m) \rightarrow \infty$ such that

$$\frac{\log m}{\log d} \rightarrow \infty \text{ as } m \rightarrow \infty.$$

Then,

$$\mathbb{E}[B_{H_L}] \sim m \int_{-2}^2 H_L(t) \mu(t) dt, \text{ as } m \rightarrow \infty.$$

If $d \leq m^{1/3}$ and $m \geq (2L)^{10}$, then

Theorem (Podder, S)

Let $L \geq 1$. Define $B_{H_L} : \mathcal{G}(m, d) \rightarrow \mathbb{R}$ by

$$B_{H_L}(Y) := \sum_{\lambda \in A(Y)} H_L(\lambda).$$

Let $d = d(m) \rightarrow \infty$ such that

$$\frac{\log m}{\log d} \rightarrow \infty \text{ as } m \rightarrow \infty.$$

Then,

$$\mathbb{E}[B_{H_L}] \sim m \int_{-2}^2 H_L(t) \mu(t) dt, \text{ as } m \rightarrow \infty.$$

If $d \leq m^{1/3}$ and $m \geq (2L)^{10}$, then

$$\mathbb{E}[B_{H_L}] = m \int_{-2}^2 H_L(t) \mu(t) dt + O\left(\frac{m \log d}{\log m}\right).$$

Theorem (Podder, S)

Let $L \geq 1$. Define $B_{H_L} : \mathcal{G}(m, d) \rightarrow \mathbb{R}$ by

$$B_{H_L}(Y) := \sum_{\lambda \in A(Y)} H_L(\lambda).$$

Let $d = d(m) \rightarrow \infty$ such that

$$\frac{\log m}{\log d} \rightarrow \infty \text{ as } m \rightarrow \infty.$$

Then,

$$\mathbb{E}[B_{H_L}] \sim m \int_{-2}^2 H_L(t) \mu(t) dt, \text{ as } m \rightarrow \infty.$$

If $d \leq m^{1/3}$ and $m \geq (2L)^{10}$, then

$$\mathbb{E}[B_{H_L}] = m \int_{-2}^2 H_L(t) \mu(t) dt + O\left(\frac{m \log d}{\log m}\right).$$

What next?

We should look at the asymptotics of the average

$$\mathbb{E} \left[\left(\sum_{\lambda \in A(X)} H_L(\lambda) - m \int_0^1 H_L(t) \mu(t) dt \right)^2 \right]$$

What next?

We should look at the asymptotics of the average

$$\mathbb{E} \left[\left(\sum_{\lambda \in A(X)} H_L(\lambda) - m \int_0^1 H_L(t) \mu(t) dt \right)^2 \right]$$

Can we find suitable growth conditions for d under which the above $\sim V_{H_L}$ and

$$\mathbb{E} \left[\left(\frac{B_{H_L}(Y) - m \int_{-2}^2 H_L(t) \mu(t) dt}{\sqrt{V_{H_L}}} \right)^r \right]$$

converges to Gaussian moments as $m \rightarrow \infty$?

Trace formula

Trace formula

Proposition

Let $Y \in \mathcal{G}(m, d)$, $d \geq 3$. Let $A(Y)$ denote the family of eigenvalues of \mathcal{T}' (with respect to Y). For a positive integer n , let $f_n(Y)$ denote the number of closed non-backtracking walks (CNBW)'s of length n in Y and let $C_n(Y)$ denote the number of circuits of length n in Y .

Proposition

Let $Y \in \mathcal{G}(m, d)$, $d \geq 3$. Let $A(Y)$ denote the family of eigenvalues of \mathcal{T}' (with respect to Y). For a positive integer n , let $f_n(Y)$ denote the number of closed non-backtracking walks (CNBW)'s of length n in Y and let $C_n(Y)$ denote the number of circuits of length n in Y . By a circuit, we mean a non-backtracking closed walk in which the first and last edges are distinct.

Trace formula

Proposition

Let $Y \in \mathcal{G}(m, d)$, $d \geq 3$. Let $A(Y)$ denote the family of eigenvalues of \mathcal{T}' (with respect to Y). For a positive integer n , let $f_n(Y)$ denote the number of closed non-backtracking walks (CNBW)'s of length n in Y and let $C_n(Y)$ denote the number of circuits of length n in Y . By a circuit, we mean a non-backtracking closed walk in which the first and last edges are distinct. Then, for all $n \geq 1$,

$$\sum_{\lambda \in A(Y)} X_n(\lambda) = (d-1)^{-n/2} \sum_{0 \leq r \leq \frac{n}{2}} f_{n-2r}.$$

Trace formula

Proposition

Let $Y \in \mathcal{G}(m, d)$, $d \geq 3$. Let $A(Y)$ denote the family of eigenvalues of \mathcal{T}' (with respect to Y). For a positive integer n , let $f_n(Y)$ denote the number of closed non-backtracking walks (CNBW)'s of length n in Y and let $C_n(Y)$ denote the number of circuits of length n in Y . By a circuit, we mean a non-backtracking closed walk in which the first and last edges are distinct. Then, for all $n \geq 1$,

$$\sum_{\lambda \in A(Y)} X_n(\lambda) = (d-1)^{-n/2} \sum_{0 \leq r \leq \frac{n}{2}} f_{n-2r}.$$

Also, for all $n \geq 2$, denoting $Z_n(x) = X_n(x) - X_{n-2}(x)$,

$$\sum_{\lambda \in A(Y)} Z_n(\lambda) = \frac{C_n(Y)}{(d-1)^{n/2}} - \begin{cases} \frac{(d-2)m}{(d-1)^{n/2}} & \text{if } n \text{ is even,} \\ 0 & \text{if } n \text{ is odd.} \end{cases}$$

Trace formula

Proposition

Let $Y \in \mathcal{G}(m, d)$, $d \geq 3$. Let $A(Y)$ denote the family of eigenvalues of \mathcal{T}' (with respect to Y). For a positive integer n , let $f_n(Y)$ denote the number of closed non-backtracking walks (CNBW)'s of length n in Y and let $C_n(Y)$ denote the number of circuits of length n in Y . By a circuit, we mean a non-backtracking closed walk in which the first and last edges are distinct. Then, for all $n \geq 1$,

$$\sum_{\lambda \in A(Y)} X_n(\lambda) = (d-1)^{-n/2} \sum_{0 \leq r \leq \frac{n}{2}} f_{n-2r}.$$

Also, for all $n \geq 2$, denoting $Z_n(x) = X_n(x) - X_{n-2}(x)$,

$$\sum_{\lambda \in A(Y)} Z_n(\lambda) = \frac{C_n(Y)}{(d-1)^{n/2}} - \begin{cases} \frac{(d-2)m}{(d-1)^{n/2}} & \text{if } n \text{ is even,} \\ 0 & \text{if } n \text{ is odd.} \end{cases}$$

Trace formula

Proposition

Let $Y \in \mathcal{G}(m, d)$, $d \geq 3$. Let $A(Y)$ denote the family of eigenvalues of \mathcal{T}' (with respect to Y). For a positive integer n , let $f_n(Y)$ denote the number of closed non-backtracking walks (CNBW)'s of length n in Y and let $C_n(Y)$ denote the number of circuits of length n in Y . By a circuit, we mean a non-backtracking closed walk in which the first and last edges are distinct. Then, for all $n \geq 1$,

$$\sum_{\lambda \in A(Y)} X_n(\lambda) = (d-1)^{-n/2} \sum_{0 \leq r \leq \frac{n}{2}} f_{n-2r}.$$

Also, for all $n \geq 2$, denoting $Z_n(x) = X_n(x) - X_{n-2}(x)$,

$$\sum_{\lambda \in A(Y)} Z_n(\lambda) = \frac{C_n(Y)}{(d-1)^{n/2}} - \begin{cases} \frac{(d-2)m}{(d-1)^{n/2}} & \text{if } n \text{ is even,} \\ 0 & \text{if } n \text{ is odd.} \end{cases}$$