

Strong contrasting diffusivity in general oscillating domains: Homogenization of optimal control problems

Abu Sufian


Joint work with Prof. A. K. Nandakumaran

Department of Mathematics
Indian Institute of Science
Bangaluru-12


IISc-IISER Pune Twenty-20 Symposium

September 18, 2021

Oscillating domain




$$\varepsilon = \frac{1}{5}$$




$$\varepsilon = \frac{1}{10}$$


Oscillating domain




$$\varepsilon = \frac{1}{5}$$



$$\varepsilon = \frac{1}{10}$$



$$\varepsilon = \frac{1}{50}$$



$$\varepsilon = \frac{1}{100}$$

Introduction: Strong contrasting diffusivity

- Partial differential equations (PDEs) with strong contrasting diffusivity are appeared in several context such as: modeling of several multi-scale physical problems such as the double porosity model, effective properties of composite material with soft and hard core, effective conductivity of composites made of materials having high and low conductivities, etc.

Introduction: Strong contrasting diffusivity

- Partial differential equations (PDEs) with strong contrasting diffusivity are appeared in several context such as: modeling of several multi-scale physical problems such as the double porosity model, effective properties of composite material with soft and hard core, effective conductivity of composites made of materials having high and low conductivities, etc.




Figure 1: Composite material

- G. P. Panasenko, N. S. Bakhvalov, A. Yu. Kosarev , H. Charef, M. Bellieudand, G. Bouchitté had studied several problem on this kind of composites.

- G. P. Panasenko, N. S. Bakhvalov, A. Yu. Kosarev , H. Charef, M. Bellieudand, G. Bouchitté had studied several problem on this kind of composites.
- Recently homogenization of a hyperbolic equation is investigated by A. K. Nandakumaran and Ali Sili [[Diff. and Int. Equ . \(2016\)](#)].

- G. P. Panasenko, N. S. Bakhvalov, A. Yu. Kosarev , H. Charef, M. Bellieudand, G. Bouchitté had studied several problem on this kind of composites.
- Recently homogenization of a hyperbolic equation is investigated by A. K. Nandakumaran and Ali Sili [[Diff. and Int. Equ . \(2016\)](#)].
- A. Gaudiello and A. Sili [[SIAM J. Math. Anal.\(2015\)](#)] had studied a variational problem in pillar type oscillating domain where reference pillar is made by two- materials with opposite behavior.

- G. P. Panasenko, N. S. Bakhvalov, A. Yu. Kosarev , H. Charef, M. Bellieudand, G. Bouchitté had studied several problem on this kind of composites.
- Recently homogenization of a hyperbolic equation is investigated by A. K. Nandakumaran and Ali Sili [[Diff. and Int. Equ . \(2016\)](#)].
- A. Gaudiello and A. Sili [[SIAM J. Math. Anal.\(2015\)](#)] had studied a variational problem in pillar type oscillating domain where reference pillar is made by two- materials with opposite behavior.

To be more precised, they have considered domain like the following;




Figure 2: Pillar type oscillating domain

- Here we are allowing the reference cell to be very general as long as the cross section of the reference cell in x_1 direction is connected and having certain properties.

- Here we are allowing the reference cell to be very general as long as the cross section of the reference cell in x_1 direction is connected and having certain properties. For example,




Figure 3: Typical example of reference cells




Figure 4: Reference cell




Figure 5: Oscillating domain

For $\varepsilon = \frac{1}{m}$ where $m \in \mathbb{Z}^+$, (in fact, one can take any $\varepsilon \rightarrow 0$) define

$$\blacksquare C_\varepsilon = \bigcup_{k=0}^{m-1} \{(x_1, x_2) : x_1 \in (k\varepsilon + \varepsilon Y_c(x_2)), x_2 \in (0, 1)\},$$

For $\varepsilon = \frac{1}{m}$ where $m \in \mathbb{Z}^+$, (in fact, one can take any $\varepsilon \rightarrow 0$) define

$$\blacksquare C_\varepsilon = \bigcup_{k=0}^{m-1} \{(x_1, x_2) : x_1 \in (k\varepsilon + \varepsilon Y_c(x_2)), x_2 \in (0, 1)\},$$

$$\blacksquare I_\varepsilon = \bigcup_{k=0}^{m-1} \{(x_1, x_2) : x_1 \in (k\varepsilon + \varepsilon Y_1(x_2)), x_2 \in (0, 1)\}.$$

For $\varepsilon = \frac{1}{m}$ where $m \in \mathbb{Z}^+$, (in fact, one can take any $\varepsilon \rightarrow 0$) define

- $C_\varepsilon = \bigcup_{k=0}^{m-1} \{(x_1, x_2) : x_1 \in (k\varepsilon + \varepsilon Y_C(x_2)), x_2 \in (0, 1)\},$
- $I_\varepsilon = \bigcup_{k=0}^{m-1} \{(x_1, x_2) : x_1 \in (k\varepsilon + \varepsilon Y_I(x_2)), x_2 \in (0, 1)\}.$
- $\Omega_\varepsilon^+ = (\overline{I_\varepsilon \cup C_\varepsilon})^o, \Omega^- = (0, 1) \times (0, -1).$

For $\varepsilon = \frac{1}{m}$ where $m \in \mathbb{Z}^+$, (in fact, one can take any $\varepsilon \rightarrow 0$) define

$$\blacksquare C_\varepsilon = \bigcup_{k=0}^{m-1} \{(x_1, x_2) : x_1 \in (k\varepsilon + \varepsilon Y_C(x_2)), x_2 \in (0, 1)\},$$

$$\blacksquare I_\varepsilon = \bigcup_{k=0}^{m-1} \{(x_1, x_2) : x_1 \in (k\varepsilon + \varepsilon Y_I(x_2)), x_2 \in (0, 1)\}.$$

$$\blacksquare \Omega_\varepsilon^+ = (\overline{I_\varepsilon \cup C_\varepsilon})^o, \Omega^- = (0, 1) \times (0, -1).$$

$$\blacksquare \text{The oscillating domain, } \Omega_\varepsilon = \left(\overline{\Omega_\varepsilon^+ \cup \Omega^-}\right)^o.$$

For $\varepsilon = \frac{1}{m}$ where $m \in \mathbb{Z}^+$, (in fact, one can take any $\varepsilon \rightarrow 0$) define

$$\blacksquare C_\varepsilon = \bigcup_{k=0}^{m-1} \{(x_1, x_2) : x_1 \in (k\varepsilon + \varepsilon Y_C(x_2)), x_2 \in (0, 1)\},$$

$$\blacksquare I_\varepsilon = \bigcup_{k=0}^{m-1} \{(x_1, x_2) : x_1 \in (k\varepsilon + \varepsilon Y_I(x_2)), x_2 \in (0, 1)\}.$$

$$\blacksquare \Omega_\varepsilon^+ = (\overline{I_\varepsilon \cup C_\varepsilon})^o, \Omega^- = (0, 1) \times (0, -1).$$

$$\blacksquare \text{The oscillating domain, } \Omega_\varepsilon = \left(\overline{\Omega_\varepsilon^+ \cup \Omega^-}\right)^o.$$

$$\blacksquare \Omega^+ = (0, 1) \times (0, 1).$$

For $\varepsilon = \frac{1}{m}$ where $m \in \mathbb{Z}^+$, (in fact, one can take any $\varepsilon \rightarrow 0$) define

$$\blacksquare C_\varepsilon = \bigcup_{k=0}^{m-1} \{(x_1, x_2) : x_1 \in (k\varepsilon + \varepsilon Y_C(x_2)), x_2 \in (0, 1)\},$$

$$\blacksquare I_\varepsilon = \bigcup_{k=0}^{m-1} \{(x_1, x_2) : x_1 \in (k\varepsilon + \varepsilon Y_I(x_2)), x_2 \in (0, 1)\}.$$

$$\blacksquare \Omega_\varepsilon^+ = (\overline{I_\varepsilon \cup C_\varepsilon})^\circ, \Omega^- = (0, 1) \times (0, -1).$$

$$\blacksquare \text{The oscillating domain, } \Omega_\varepsilon = \left(\overline{\Omega_\varepsilon^+ \cup \Omega^-}\right)^\circ.$$

$$\blacksquare \Omega^+ = (0, 1) \times (0, 1). \text{The limit domain is defined as } \Omega = (\overline{\Omega^+ \cup \Omega^-})^\circ.$$

For $\varepsilon = \frac{1}{m}$ where $m \in \mathbb{Z}^+$, (in fact, one can take any $\varepsilon \rightarrow 0$) define

$$\blacksquare C_\varepsilon = \bigcup_{k=0}^{m-1} \{(x_1, x_2) : x_1 \in (k\varepsilon + \varepsilon Y_C(x_2)), x_2 \in (0, 1)\},$$

$$\blacksquare I_\varepsilon = \bigcup_{k=0}^{m-1} \{(x_1, x_2) : x_1 \in (k\varepsilon + \varepsilon Y_I(x_2)), x_2 \in (0, 1)\}.$$

$$\blacksquare \Omega_\varepsilon^+ = (\overline{I_\varepsilon \cup C_\varepsilon})^\circ, \Omega^- = (0, 1) \times (0, -1).$$

$$\blacksquare \text{The oscillating domain, } \Omega_\varepsilon = \left(\overline{\Omega_\varepsilon^+ \cup \Omega^-}\right)^\circ.$$

$$\blacksquare \Omega^+ = (0, 1) \times (0, 1). \text{The limit domain is defined as } \Omega = (\overline{\Omega^+ \cup \Omega^-})^\circ.$$

$$\blacksquare \text{The interface between } \Omega^+ \text{ and } \Omega^- \text{ is denoted by } \gamma, \text{ which is given by } \gamma = \{(x_1, 0) : x_1 \in (0, 1)\}.$$

- We want to consider the following ε dependent variational problem,

$$\left\{ \begin{array}{l} \text{find } u_\varepsilon \in H^1(\Omega_\varepsilon) \text{ such that} \\ \int_{\Omega_\varepsilon} \left(\chi_{\Omega^-} + \chi_{C_\varepsilon} + \varepsilon^2 \chi_{I_\varepsilon} \right) \nabla u_\varepsilon \nabla \phi + \int_{\Omega_\varepsilon} u_\varepsilon \phi = \int_{\Omega_\varepsilon} f \phi, \end{array} \right. \quad (1)$$

for all $\phi \in H^1(\Omega_\varepsilon)$, where $f \in L^2(\Omega)$.

- We want to consider the following ε dependent variational problem,

$$\left\{ \begin{array}{l} \text{find } u_\varepsilon \in H^1(\Omega_\varepsilon) \text{ such that} \\ \int_{\Omega_\varepsilon} (\chi_{\Omega^-} + \chi_{C_\varepsilon} + \varepsilon^2 \chi_{I_\varepsilon}) \nabla u_\varepsilon \nabla \phi + \int_{\Omega_\varepsilon} u_\varepsilon \phi = \int_{\Omega_\varepsilon} f \phi, \end{array} \right. \quad (1)$$

for all $\phi \in H^1(\Omega_\varepsilon)$, where $f \in L^2(\Omega)$. The Lax-Milgram theorem ensures the existence and the uniqueness of the solution u_ε of the problem (1).

- We want to consider the following ε dependent variational problem,

$$\left\{ \begin{array}{l} \text{find } u_\varepsilon \in H^1(\Omega_\varepsilon) \text{ such that} \\ \int_{\Omega_\varepsilon} \left(\chi_{\Omega^-} + \chi_{C_\varepsilon} + \varepsilon^2 \chi_{I_\varepsilon} \right) \nabla u_\varepsilon \nabla \phi + \int_{\Omega_\varepsilon} u_\varepsilon \phi = \int_{\Omega_\varepsilon} f \phi, \end{array} \right. \quad (1)$$

for all $\phi \in H^1(\Omega_\varepsilon)$, where $f \in L^2(\Omega)$. The Lax-Milgram theorem ensures the existence and the uniqueness of the solution u_ε of the problem (1).

- Our aim is to analyze the asymptotic behavior of the above variational form as the oscillating parameter $\varepsilon \rightarrow 0$.

The unfolded domain corresponding to the upper part Ω_ε^+ is given by

$$\blacksquare \Omega^u = \{(x_1, x_2, y_1) : (x_1, x_2) \in \Omega^+, y_1 \in Y(x_2)\}.$$

The unfolded domain corresponding to the upper part Ω_ε^+ is given by

- $\Omega^u = \{(x_1, x_2, y_1) : (x_1, x_2) \in \Omega^+, y_1 \in Y(x_2)\}.$
- $\Omega_C^u = \{(x_1, x_2, y_1) : (x_1, x_2) \in \Omega^+, y_1 \in Y_C(x_2)\}.$

Unfolding operator and properties

The unfolded domain corresponding to the upper part Ω_ε^+ is given by

- $\Omega^u = \{(x_1, x_2, y_1) : (x_1, x_2) \in \Omega^+, y_1 \in Y(x_2)\}.$
- $\Omega_c^u = \{(x_1, x_2, y_1) : (x_1, x_2) \in \Omega^+, y_1 \in Y_c(x_2)\}.$

Definition.


(The unfolding operator) Let $\phi^\varepsilon : \Omega^u \rightarrow \Omega_\varepsilon^+$ be defined as $\phi^\varepsilon(x_1, x_2, y_1) = (\varepsilon \left[\frac{x_1}{\varepsilon} \right] + \varepsilon y_1, x_2)$. The ε -unfolding of a function $u : \Omega_\varepsilon^+ \rightarrow \mathbb{R}$ is the function $u \circ \phi^\varepsilon : \Omega^u \rightarrow \mathbb{R}$. The operator which maps every function $u : \Omega_\varepsilon^+ \rightarrow \mathbb{R}$ to its ε -unfolding is called the unfolding operator. Let the unfolding operator is denoted by T^ε , that is,

$$T^\varepsilon : \{u : \Omega_\varepsilon^+ \rightarrow \mathbb{R}\} \rightarrow \{T^\varepsilon u : \Omega^u \rightarrow \mathbb{R}\}$$

is defined by

$$T^\varepsilon u(x_1, x_2, y_1) = u\left(\varepsilon \left[\frac{x_1}{\varepsilon} \right] + \varepsilon y_1, x_2\right) \text{ for all } (x_1, x_2, y_1) \in \Omega^u.$$

Limit domain



Limit function spaces

For any function ϕ defined on Ω , we may write $\phi = \phi^+ \chi_{\Omega^+} + \phi^- \chi_{\Omega^-} = (\phi^+, \phi^-)$ throughout the presentation.

Limit function spaces

For any function ϕ defined on Ω , we may write

$\phi = \phi^+ \chi_{\Omega^+} + \phi^- \chi_{\Omega^-} = (\phi^+, \phi^-)$ throughout the presentation.

■ Define

$H(\Omega) = \{\phi : \phi^+ \in L^2((0,1); H^1(0,1)), \phi^- \in H^1(\Omega^-), \phi^+ = \phi^- \text{ on } \gamma\}$ with the following norm

$$\|\phi\|_{H(\Omega)} = \|\phi^-\|_{H^1(\Omega^-)} + \|\phi^+\|_{L^2(\Omega^+)} + \left\| \frac{\partial \phi^+}{\partial x_2} \right\|_{L^2(\Omega^+)}.$$

For any function ϕ defined on Ω , we may write

$\phi = \phi^+ \chi_{\Omega^+} + \phi^- \chi_{\Omega^-} = (\phi^+, \phi^-)$ throughout the presentation.

■ Define

$H(\Omega) = \{\phi : \phi^+ \in L^2((0,1); H^1(0,1)), \phi^- \in H^1(\Omega^-), \phi^+ = \phi^- \text{ on } \gamma\}$ with the following norm

$$\|\phi\|_{H(\Omega)} = \|\phi^-\|_{H^1(\Omega^-)} + \|\phi^+\|_{L^2(\Omega^+)} + \left\| \frac{\partial \phi^+}{\partial x_2} \right\|_{L^2(\Omega^+)}.$$

■ For any $x_2 \in (0,1)$, define $V^{x_2} = \{w \in H^1(Y(x_2)) : w = 0 \text{ in } Y_c(x_2)\}$ with the following norm

$$\|w\|_{V^{x_2}} = \|w\|_{L^2(Y(x_2))} + \left\| \frac{\partial w}{\partial y_1} \right\|_{L^2(Y(x_2))}.$$

Limit function spaces

For any function ϕ defined on Ω , we may write

$\phi = \phi^+ \chi_{\Omega^+} + \phi^- \chi_{\Omega^-} = (\phi^+, \phi^-)$ throughout the presentation.

■ Define

$H(\Omega) = \{\phi : \phi^+ \in L^2((0,1); H^1(0,1)), \phi^- \in H^1(\Omega^-), \phi^+ = \phi^- \text{ on } \gamma\}$ with the following norm

$$\|\phi\|_{H(\Omega)} = \|\phi^-\|_{H^1(\Omega^-)} + \|\phi^+\|_{L^2(\Omega^+)} + \left\| \frac{\partial \phi^+}{\partial x_2} \right\|_{L^2(\Omega^+)}.$$

■ For any $x_2 \in (0,1)$, define $V^{x_2} = \{w \in H^1(Y(x_2)) : w = 0 \text{ in } Y_c(x_2)\}$ with the following norm

$$\|w\|_{V^{x_2}} = \|w\|_{L^2(Y(x_2))} + \left\| \frac{\partial w}{\partial y_1} \right\|_{L^2(Y(x_2))}.$$

■ $V(\Omega) = \left\{ \psi \in L^2(\Omega^u) : \psi = 0 \text{ in } \Omega_c^u, \frac{\partial \psi}{\partial y_1} \in L^2(\Omega^u) \right\}$ with the following norm

$$\|\psi\|_{V(\Omega)} = \|\psi\|_{L^2(\Omega^u)} + \left\| \frac{\partial \psi}{\partial y_1} \right\|_{L^2(\Omega^u)}.$$

Limit problem

The limit variational problem :find $u = (u^+, u^-) \in H(\Omega)$ such that

$$\left\{ \begin{array}{l} \int_{\Omega^+} |Y_c(x_2)| \frac{\partial u^+}{\partial x_2} \frac{\partial \phi}{\partial x_2} + \int_{\Omega^+} \alpha(x) u^+ \phi + \int_{\Omega^-} u^- \phi \\ \qquad \qquad \qquad + \int_{\Omega^-} \nabla u^- \nabla \phi = \int_{\Omega^+} \alpha(x) f \phi + \int_{\Omega^-} f \phi, \end{array} \right.$$

for all $\phi \in H(\Omega)$,

Limit problem

The limit variational problem :find $u = (u^+, u^-) \in H(\Omega)$ such that

$$\left\{ \begin{array}{l} \int_{\Omega^+} |Y_c(x_2)| \frac{\partial u^+}{\partial x_2} \frac{\partial \phi}{\partial x_2} + \int_{\Omega^+} \alpha(x) u^+ \phi + \int_{\Omega^-} u^- \phi \\ \quad + \int_{\Omega^-} \nabla u^- \nabla \phi = \int_{\Omega^+} \alpha(x) f \phi + \int_{\Omega^-} f \phi, \end{array} \right.$$

for all $\phi \in H(\Omega)$, here $\alpha(x) = \left(|Y(x_2)| - \int_{Y_1(x_2)} \xi dy_1 \right)$,

Limit problem

The limit variational problem :find $u = (u^+, u^-) \in H(\Omega)$ such that

$$\left\{ \begin{array}{l} \int_{\Omega^+} |Y_c(x_2)| \frac{\partial u^+}{\partial x_2} \frac{\partial \phi}{\partial x_2} + \int_{\Omega^+} \alpha(x) u^+ \phi + \int_{\Omega^-} u^- \phi \\ \quad + \int_{\Omega^-} \nabla u^- \nabla \phi = \int_{\Omega^+} \alpha(x) f \phi + \int_{\Omega^-} f \phi, \end{array} \right.$$

for all $\phi \in H(\Omega)$, here $\alpha(x) = \left(|Y(x_2)| - \int_{Y_I(x_2)} \xi dy_1 \right)$, where

$$\left\{ \int_{Y(x_2)} \frac{\partial \xi(x_2, y_1)}{\partial y_1} \frac{\partial w(y_1)}{\partial y_1} + \int_{Y(x_2)} \xi(x_2, y_1) w(y_1) = \int_{Y(x_2)} w(y_1), \text{ for all } w \in V^{x_2}. \right.$$

Limit problem

The limit variational problem :find $u = (u^+, u^-) \in H(\Omega)$ such that

$$\left\{ \begin{array}{l} \int_{\Omega^+} |Y_c(x_2)| \frac{\partial u^+}{\partial x_2} \frac{\partial \phi}{\partial x_2} + \int_{\Omega^+} \alpha(x) u^+ \phi + \int_{\Omega^-} u^- \phi \\ \quad + \int_{\Omega^-} \nabla u^- \nabla \phi = \int_{\Omega^+} \alpha(x) f \phi + \int_{\Omega^-} f \phi, \end{array} \right.$$

for all $\phi \in H(\Omega)$, here $\alpha(x) = \left(|Y(x_2)| - \int_{Y_1(x_2)} \xi dy_1 \right)$, where

$$\left\{ \int_{Y(x_2)} \frac{\partial \xi(x_2, y_1)}{\partial y_1} \frac{\partial w(y_1)}{\partial y_1} + \int_{Y(x_2)} \xi(x_2, y_1) w(y_1) = \int_{Y(x_2)} w(y_1), \text{ for all } w \in V^{x_2}. \right.$$

- Now we will show $\alpha(x) = \left(|Y(x_2)| - \int_{Y_1(x_2)} \xi \right) > 0$. By taking $w = \xi$ in the cell problem, we get

Limit problem

The limit variational problem :find $u = (u^+, u^-) \in H(\Omega)$ such that

$$\left\{ \begin{array}{l} \int_{\Omega^+} |Y_c(x_2)| \frac{\partial u^+}{\partial x_2} \frac{\partial \phi}{\partial x_2} + \int_{\Omega^+} \alpha(x) u^+ \phi + \int_{\Omega^-} u^- \phi \\ \quad + \int_{\Omega^-} \nabla u^- \nabla \phi = \int_{\Omega^+} \alpha(x) f \phi + \int_{\Omega^-} f \phi, \end{array} \right.$$

for all $\phi \in H(\Omega)$, here $\alpha(x) = \left(|Y(x_2)| - \int_{Y_1(x_2)} \xi dy_1 \right)$, where

$$\left\{ \begin{array}{l} \int_{Y(x_2)} \frac{\partial \xi(x_2, y_1)}{\partial y_1} \frac{\partial w(y_1)}{\partial y_1} + \int_{Y(x_2)} \xi(x_2, y_1) w(y_1) = \int_{Y(x_2)} w(y_1), \text{ for all } w \in V^{x_2}. \end{array} \right.$$

- Now we will show $\alpha(x) = \left(|Y(x_2)| - \int_{Y_1(x_2)} \xi \right) > 0$. By taking $w = \xi$ in the cell problem, we get

$$\int_{Y(x_2)} \left(\left| \frac{\partial \xi}{\partial y_1} \right|^2 + \xi^2 \right) = \int_{Y(x_2)} \xi$$

Limit problem

The limit variational problem :find $u = (u^+, u^-) \in H(\Omega)$ such that

$$\left\{ \begin{array}{l} \int_{\Omega^+} |Y_c(x_2)| \frac{\partial u^+}{\partial x_2} \frac{\partial \phi}{\partial x_2} + \int_{\Omega^+} \alpha(x) u^+ \phi + \int_{\Omega^-} u^- \phi \\ \quad + \int_{\Omega^-} \nabla u^- \nabla \phi = \int_{\Omega^+} \alpha(x) f \phi + \int_{\Omega^-} f \phi, \end{array} \right.$$

for all $\phi \in H(\Omega)$, here $\alpha(x) = \left(|Y(x_2)| - \int_{Y_1(x_2)} \xi dy_1 \right)$, where

$$\left\{ \begin{array}{l} \int_{Y(x_2)} \frac{\partial \xi(x_2, y_1)}{\partial y_1} \frac{\partial w(y_1)}{\partial y_1} + \int_{Y(x_2)} \xi(x_2, y_1) w(y_1) = \int_{Y(x_2)} w(y_1), \text{ for all } w \in V^{x_2}. \end{array} \right.$$

- Now we will show $\alpha(x) = \left(|Y(x_2)| - \int_{Y_1(x_2)} \xi \right) > 0$. By taking $w = \xi$ in the cell problem, we get

$$\int_{Y(x_2)} \left(\left| \frac{\partial \xi}{\partial y_1} \right|^2 + \xi^2 \right) = \int_{Y(x_2)} \xi \Rightarrow \|\xi\|_{L^2(Y_1(x_2))} \leq |Y_1(x_2)|^{\frac{1}{2}}$$

Limit problem

The limit variational problem :find $u = (u^+, u^-) \in H(\Omega)$ such that

$$\left\{ \begin{array}{l} \int_{\Omega^+} |Y_C(x_2)| \frac{\partial u^+}{\partial x_2} \frac{\partial \phi}{\partial x_2} + \int_{\Omega^+} \alpha(x) u^+ \phi + \int_{\Omega^-} u^- \phi \\ \quad + \int_{\Omega^-} \nabla u^- \nabla \phi = \int_{\Omega^+} \alpha(x) f \phi + \int_{\Omega^-} f \phi, \end{array} \right.$$

for all $\phi \in H(\Omega)$, here $\alpha(x) = \left(|Y(x_2)| - \int_{Y_I(x_2)} \xi dy_1 \right)$, where

$$\left\{ \begin{array}{l} \int_{Y(x_2)} \frac{\partial \xi(x_2, y_1)}{\partial y_1} \frac{\partial w(y_1)}{\partial y_1} + \int_{Y(x_2)} \xi(x_2, y_1) w(y_1) = \int_{Y(x_2)} w(y_1), \text{ for all } w \in V^{x_2}. \end{array} \right.$$

- Now we will show $\alpha(x) = \left(|Y(x_2)| - \int_{Y_I(x_2)} \xi \right) > 0$. By taking $w = \xi$ in the cell problem, we get

$$\int_{Y(x_2)} \left(\left| \frac{\partial \xi}{\partial y_1} \right|^2 + \xi^2 \right) = \int_{Y(x_2)} \xi \Rightarrow \|\xi\|_{L^2(Y_I(x_2))} \leq |Y_I(x_2)|^{\frac{1}{2}}$$

$$\begin{aligned} \left(|Y(x_2)| - \int_{Y_I(x_2)} \xi \right) &\geq (|Y(x_2)| - |Y_I(x_2)|)^{1/2} \|\xi\|_{L^2(Y_I(x_2))} \\ &\geq |Y(x_2)| - |Y_I(x_2)| = |Y_C(x_2)| > \delta. \end{aligned}$$

Theorem (Nandakumaran-Sufian, JDE-2021).

For every $\varepsilon > 0$, let u_ε be the unique solution to the considered variational problem. Let $H(\Omega)$ and V^{x_2} be defined as earlier and $u = (u^+, u^-) \in H(\Omega)$ be the unique solution of the limit variational form.

Theorem (Nandakumaran-Sufian, JDE-2021).

For every $\varepsilon > 0$, let u_ε be the unique solution to the considered variational problem. Let $H(\Omega)$ and V^{x_2} be defined as earlier and $u = (u^+, u^-) \in H(\Omega)$ be the unique solution of the limit variational form. Then

$$\begin{cases} u_\varepsilon^- \rightharpoonup u^- \text{ weakly in } H^1(\Omega^-), \\ \begin{cases} \widetilde{u_\varepsilon^+} \rightharpoonup |Y(x_2)|u^+ + \int_{Y_I(x_2)} (f - u^+) \xi(x_2, y_1) dy_1 \\ \chi_{c_\varepsilon}^+ \frac{\partial \widetilde{u_\varepsilon^+}}{\partial x_1} \rightharpoonup 0, \quad \chi_{c_\varepsilon}^+ \frac{\partial \widetilde{u_\varepsilon^+}}{\partial x_2} \rightharpoonup |Y_c(x_2)| \frac{\partial u^+}{\partial x_2} \\ \varepsilon \chi_{I_\varepsilon}^+ \frac{\partial \widetilde{u_\varepsilon^+}}{\partial x_1} \rightharpoonup (f - u^+) \int_{Y_I(x_2)} \frac{\partial \xi}{\partial y_1} dy_1, \quad \varepsilon \chi_{I_\varepsilon}^+ \frac{\partial \widetilde{u_\varepsilon^+}}{\partial x_2} \rightharpoonup 0 \\ \text{weakly in } L^2(\Omega^+) \end{cases} \end{cases}$$

as $\varepsilon \rightarrow 0$.

For $\theta_\varepsilon \in L^2(C_\varepsilon)$ consider the cost functional

$$J_\varepsilon(u_\varepsilon, \theta_\varepsilon) = \frac{1}{2} \int_{\Omega_\varepsilon} |u_\varepsilon - u_d|^2 + \frac{\beta}{2} \int_{C_\varepsilon} |\theta_\varepsilon|^2$$

For $\theta_\varepsilon \in L^2(C_\varepsilon)$ consider the cost functional

$$J_\varepsilon(u_\varepsilon, \theta_\varepsilon) = \frac{1}{2} \int_{\Omega_\varepsilon} |u_\varepsilon - u_d|^2 + \frac{\beta}{2} \int_{C_\varepsilon} |\theta_\varepsilon|^2$$

where u_ε is the unique solution of the following variational problem: for $f \in L^2(\Omega)$

$$\left\{ \begin{array}{l} \text{find } u_\varepsilon \in H^1(\Omega_\varepsilon) \text{ such that} \\ \int_{\Omega_\varepsilon} (\chi_{\Omega^-} + \chi_{C_\varepsilon} + \varepsilon^2 \chi_{I_\varepsilon}) \nabla u_\varepsilon \nabla \phi + u_\varepsilon \phi = \int_{\Omega_\varepsilon} f \phi + \int_{\Omega_\varepsilon} \chi_{C_\varepsilon} \theta_\varepsilon \phi, \\ \text{for all } \phi \in H^1(\Omega_\varepsilon). \end{array} \right.$$

For $\theta_\varepsilon \in L^2(C_\varepsilon)$ consider the cost functional

$$J_\varepsilon(u_\varepsilon, \theta_\varepsilon) = \frac{1}{2} \int_{\Omega_\varepsilon} |u_\varepsilon - u_d|^2 + \frac{\beta}{2} \int_{C_\varepsilon} |\theta_\varepsilon|^2$$

where u_ε is the unique solution of the following variational problem: for $f \in L^2(\Omega)$

$$\left\{ \begin{array}{l} \text{find } u_\varepsilon \in H^1(\Omega_\varepsilon) \text{ such that} \\ \int_{\Omega_\varepsilon} (\chi_{\Omega^-} + \chi_{C_\varepsilon} + \varepsilon^2 \chi_{I_\varepsilon}) \nabla u_\varepsilon \nabla \phi + u_\varepsilon \phi = \int_{\Omega_\varepsilon} f \phi + \int_{\Omega_\varepsilon} \chi_{C_\varepsilon} \theta_\varepsilon \phi, \\ \text{for all } \phi \in H^1(\Omega_\varepsilon). \end{array} \right.$$

The optimal control problem is to find $(\bar{u}_\varepsilon, \bar{\theta}_\varepsilon) \in H^1(\Omega_\varepsilon) \times L^2(C_\varepsilon)$ such that

$$J_\varepsilon(\bar{u}_\varepsilon, \bar{\theta}_\varepsilon) = \inf \{J_\varepsilon(u_\varepsilon, \theta_\varepsilon)\}. \quad (2)$$

- We will use the characterization of optimal control $\bar{\theta}_\varepsilon$ by introducing the adjoint state \bar{v}_ε which is the solution of the following variational form

$$\left\{ \begin{array}{l} \text{find } \bar{v}_\varepsilon \in H^1(\Omega_\varepsilon) \text{ such that} \\ \int_{\Omega_\varepsilon} \left(\chi_{\Omega^-} + \chi_{C_\varepsilon} + \varepsilon^2 \chi_{I_\varepsilon} \right) \nabla \bar{v}_\varepsilon \nabla \phi + \bar{v}_\varepsilon \phi = \int_{\Omega_\varepsilon} (\bar{u}_\varepsilon - u_d) \phi, \end{array} \right. \quad (3)$$

for all $\phi \in H^1(\Omega_\varepsilon)$.

- We will use the characterization of optimal control $\bar{\theta}_\varepsilon$ by introducing the adjoint state \bar{v}_ε which is the solution of the following variational form

$$\left\{ \begin{array}{l} \text{find } \bar{v}_\varepsilon \in H^1(\Omega_\varepsilon) \text{ such that} \\ \int_{\Omega_\varepsilon} \left(\chi_{\Omega^-} + \chi_{C_\varepsilon} + \varepsilon^2 \chi_{I_\varepsilon} \right) \nabla \bar{v}_\varepsilon \nabla \phi + \bar{v}_\varepsilon \phi = \int_{\Omega_\varepsilon} (\bar{u}_\varepsilon - u_d) \phi, \end{array} \right. \quad (3)$$

for all $\phi \in H^1(\Omega_\varepsilon)$.

Theorem.

Let $(\bar{u}_\varepsilon, \bar{\theta}_\varepsilon)$ be the optimal solution to the optimal control problem (2) and \bar{v}_ε be the unique solution of (3). Then $\bar{\theta}_\varepsilon$ is characterized by

$$\bar{\theta}_\varepsilon = -\chi_{C_\varepsilon} \frac{1}{\beta} \bar{v}_\varepsilon. \quad (4)$$

Cost functional: For $\theta \in L^2(\Omega^+)$

$$J(u, \theta) = \frac{1}{2} \int_{\Omega^+} \int_{Y(x_2)} |(1 - \xi)u^+ + f\xi - u_d|^2 + \frac{1}{2} \int_{\Omega^-} |u^- - u_d|^2 \\ + \frac{\beta}{2} \int_{\Omega^+} |Y_C(x_2)| |\theta|^2$$

Cost functional: For $\theta \in L^2(\Omega^+)$

$$J(u, \theta) = \frac{1}{2} \int_{\Omega^+} \int_{Y(x_2)} |(1 - \xi)u^+ + f\xi - u_d|^2 + \frac{1}{2} \int_{\Omega^-} |u^- - u_d|^2 \\ + \frac{\beta}{2} \int_{\Omega^+} |Y_c(x_2)| |\theta|^2$$

Limit state equation:

$$\left\{ \begin{array}{l} \text{find } u \in H(\Omega), \text{ such that,} \\ \int_{\Omega^+} |Y_c(x_2)| \frac{\partial u^+}{\partial x_2} \frac{\partial \phi}{\partial x_2} + \int_{\Omega^+} \alpha(x) u^+ \phi + \int_{\Omega^-} u \phi + \int_{\Omega^-} \nabla u^- \nabla \phi \\ \qquad \qquad \qquad = \int_{\Omega^+} \alpha(x) f \phi + \int_{\Omega^-} f \phi + \int_{\Omega^+} |Y_c(x_2)| \theta \phi, \\ \text{for all } \phi \in H(\Omega). \end{array} \right.$$

Cost functional: For $\theta \in L^2(\Omega^+)$

$$J(u, \theta) = \frac{1}{2} \int_{\Omega^+} \int_{Y(x_2)} |(1 - \xi)u^+ + f\xi - u_d|^2 + \frac{1}{2} \int_{\Omega^-} |u^- - u_d|^2 \\ + \frac{\beta}{2} \int_{\Omega^+} |Y_c(x_2)| |\theta|^2$$

Limit state equation:

$$\left\{ \begin{array}{l} \text{find } u \in H(\Omega), \text{ such that,} \\ \int_{\Omega^+} |Y_c(x_2)| \frac{\partial u^+}{\partial x_2} \frac{\partial \phi}{\partial x_2} + \int_{\Omega^+} \alpha(x) u^+ \phi + \int_{\Omega^-} u \phi + \int_{\Omega^-} \nabla u^- \nabla \phi \\ \qquad \qquad \qquad = \int_{\Omega^+} \alpha(x) f \phi + \int_{\Omega^-} f \phi + \int_{\Omega^+} |Y_c(x_2)| \theta \phi, \\ \text{for all } \phi \in H(\Omega). \end{array} \right.$$

■ $J(\bar{u}, \bar{\theta}) = \inf \{J(u, \theta)\}$

Adjoint equation:

$$\left\{ \begin{array}{l} \int_{\Omega^+} |Y_c(x_2)| \frac{\partial \bar{v}^+}{\partial x_2} \frac{\partial \phi}{\partial x_2} + \int_{\Omega^+} \alpha(x) \bar{v}^+ \phi + \int_{\Omega^-} \bar{v}^- \phi + \int_{\Omega^-} (\nabla \bar{v}^- \nabla \phi \\ = \int_{\Omega^+} \left[\left(\int_{Y(x_2)} (1 - \xi)^2 dy_1 \right) \bar{u}^+ - \alpha(x) u_d + \left(\int_{Y_1(x_2)} (\xi - \xi^2) dy_1 \right) f \right] \phi \\ \quad \quad \quad + \int_{\Omega^-} (\bar{u}^- - u_d) \phi. \end{array} \right.$$

Adjoint equation:

$$\left\{ \begin{array}{l} \int_{\Omega^+} |Y_c(x_2)| \frac{\partial \bar{v}^+}{\partial x_2} \frac{\partial \phi}{\partial x_2} + \int_{\Omega^+} \alpha(x) \bar{v}^+ \phi + \int_{\Omega^-} \bar{v}^- \phi + \int_{\Omega^-} (\nabla \bar{v}^- \nabla \phi \\ = \int_{\Omega^+} \left[\left(\int_{Y(x_2)} (1 - \xi)^2 dy_1 \right) \bar{u}^+ - \alpha(x) u_d + \left(\int_{Y_1(x_2)} (\xi - \xi^2) dy_1 \right) f \right] \phi \\ \quad \quad \quad + \int_{\Omega^-} (\bar{u}^- - u_d) \phi. \end{array} \right.$$

- Optimal control is given by $\bar{\theta} = -\frac{1}{\beta} \bar{v}^+$

For $\theta_\varepsilon \in L^2(I_\varepsilon)$, consider the following L^2 -cost functional

$$J_\varepsilon(u_\varepsilon, \theta_\varepsilon) = \frac{1}{2} \int_{\Omega_\varepsilon} |u_\varepsilon - u_d|^2 + \frac{\beta}{2} \int_{I_\varepsilon} |\theta_\varepsilon|^2,$$

For $\theta_\varepsilon \in L^2(I_\varepsilon)$, consider the following L^2 -cost functional

$$J_\varepsilon(u_\varepsilon, \theta_\varepsilon) = \frac{1}{2} \int_{\Omega_\varepsilon} |u_\varepsilon - u_d|^2 + \frac{\beta}{2} \int_{I_\varepsilon} |\theta_\varepsilon|^2,$$

where u_ε is the unique solution of the following variational problem: for $f \in L^2(\Omega)$

$$\left\{ \begin{array}{l} \text{find } u_\varepsilon \in H^1(\Omega_\varepsilon) \text{ such that} \\ \int_{\Omega_\varepsilon} \left(\chi_{\Omega^-} + \chi_{C_\varepsilon} + \varepsilon^2 \chi_{I_\varepsilon} \right) \nabla u_\varepsilon \nabla \phi + u_\varepsilon \phi = \int_{\Omega_\varepsilon} f \phi + \int_{\Omega_\varepsilon} \chi_{I_\varepsilon} \theta_\varepsilon \phi, \end{array} \right. \quad (5)$$

for all $\phi \in H^1(\Omega_\varepsilon)$.

For $\theta_\varepsilon \in L^2(I_\varepsilon)$, consider the following L^2 -cost functional

$$J_\varepsilon(u_\varepsilon, \theta_\varepsilon) = \frac{1}{2} \int_{\Omega_\varepsilon} |u_\varepsilon - u_d|^2 + \frac{\beta}{2} \int_{I_\varepsilon} |\theta_\varepsilon|^2,$$

where u_ε is the unique solution of the following variational problem: for $f \in L^2(\Omega)$

$$\begin{cases} \text{find } u_\varepsilon \in H^1(\Omega_\varepsilon) \text{ such that} \\ \int_{\Omega_\varepsilon} (\chi_{\Omega^-} + \chi_{C_\varepsilon} + \varepsilon^2 \chi_{I_\varepsilon}) \nabla u_\varepsilon \nabla \phi + u_\varepsilon \phi = \int_{\Omega_\varepsilon} f \phi + \int_{\Omega_\varepsilon} \chi_{I_\varepsilon} \theta_\varepsilon \phi, \end{cases} \quad (5)$$

for all $\phi \in H^1(\Omega_\varepsilon)$. The optimal control problem is to find $(\bar{u}_\varepsilon, \bar{\theta}_\varepsilon) \in H^1(\Omega_\varepsilon) \times L^2(I_\varepsilon)$ such that

$$J_\varepsilon(\bar{u}_\varepsilon, \bar{\theta}_\varepsilon) = \inf \{ J_\varepsilon(u_\varepsilon, \theta_\varepsilon) : (u_\varepsilon, \theta_\varepsilon) \text{ satisfies (5)} \}. \quad (6)$$

Theorem (Characterization).

Let $(\bar{u}_\epsilon, \bar{\theta}_\epsilon)$ be the optimal solution to the optimal control problem (6) and \bar{v}_ϵ be the unique solution of the adjoint state. Then $\bar{\theta}_\epsilon$ can be written as $\bar{\theta}_\epsilon = -\chi_{I_\epsilon} \frac{1}{\beta} \bar{v}_\epsilon$.

Reduced cost functional: The L^2 -cost functional reduces to

$$\begin{aligned} J(u, \mathbf{u}_{11}, \theta, \theta_1) = & \frac{1}{2} \int_{\Omega^+} \int_{Y(x_2)} ((1 - \xi)u^+ + \xi f + \mathbf{u}_{11} - u_d)^2 \\ & + \int_{\Omega^-} (u^- - u_d)^2 + \frac{\beta}{2} \int_{\Omega^+} \int_{Y(x_2)} (\theta + \theta_1)^2 \end{aligned}$$

Reduced cost functional: The L^2 -cost functional reduces to

$$J(u, \mathbf{u}_{11}, \theta, \theta_1) = \frac{1}{2} \int_{\Omega^+} \int_{Y(x_2)} ((1 - \xi)u^+ + \xi f + \mathbf{u}_{11} - u_d)^2 \\ + \int_{\Omega^-} (u^- - u_d)^2 + \frac{\beta}{2} \int_{\Omega^+} \int_{Y(x_2)} (\theta + \theta_1)^2$$

Reduced state equation: The state $(u, u_{11}) \in H(\Omega) \times V(\Omega)$ satisfies the following system,

$$\left\{ \begin{array}{l} \int_{\Omega^+} |Y_c(x_2)| \frac{\partial u^+}{\partial x_2} \frac{\partial \phi^+}{\partial x_2} + \int_{\Omega^+} \alpha(x) u^+ \phi^+ + \int_{\Omega^-} \nabla u^- \nabla \phi^- + \int_{\Omega^-} u^- \phi \\ \quad = \int_{\Omega^+} \int_{Y(x_2)} ((1 - \xi)f + (1 - \xi)(\theta + \theta_1)) \phi^+ + \int_{\Omega^-} f \phi^-, \\ \int_{\Omega^u} \frac{\partial u_{11}}{\partial y_1} \frac{\partial \phi_1}{\partial y_1} + \int_{\Omega^u} u_{11} \phi_1 = \int_{\Omega^u} (\theta + \theta_1) \phi_1, \end{array} \right.$$

Reduced cost functional: The L^2 -cost functional reduces to

$$J(u, \mathbf{u}_{11}, \theta, \theta_1) = \frac{1}{2} \int_{\Omega^+} \int_{Y(x_2)} ((1 - \xi)u^+ + \xi f + \mathbf{u}_{11} - u_d)^2 \\ + \int_{\Omega^-} (u^- - u_d)^2 + \frac{\beta}{2} \int_{\Omega^+} \int_{Y(x_2)} (\theta + \theta_1)^2$$

Reduced state equation: The state $(u, u_{11}) \in H(\Omega) \times V(\Omega)$ satisfies the following system,

$$\left\{ \begin{array}{l} \int_{\Omega^+} |Y_c(x_2)| \frac{\partial u^+}{\partial x_2} \frac{\partial \phi^+}{\partial x_2} + \int_{\Omega^+} \alpha(x) u^+ \phi^+ + \int_{\Omega^-} \nabla u^- \nabla \phi^- + \int_{\Omega^-} u^- \phi \\ \quad = \int_{\Omega^+} \int_{Y(x_2)} ((1 - \xi)f + (1 - \xi)(\theta + \theta_1)) \phi^+ + \int_{\Omega^-} f \phi^-, \\ \int_{\Omega^u} \frac{\partial u_{11}}{\partial y_1} \frac{\partial \phi_1}{\partial y_1} + \int_{\Omega^u} u_{11} \phi_1 = \int_{\Omega^u} (\theta + \theta_1) \phi_1, \end{array} \right.$$

$$\blacksquare J(\bar{u}, \bar{u}_{11}, \bar{\theta}, \bar{\theta}_1) = \inf \{J(u, u_{11}, \theta, \theta_1)\}$$

- **Reduced adjoint state:** The adjoint state $(\bar{v}, \bar{v}_{11}) \in H(\Omega) \times V(\Omega)$ satisfies the following system,

$$\left\{ \begin{array}{l} \int_{\Omega^+} |Y_c(x_2)| \frac{\partial \bar{v}^+}{\partial x_2} \frac{\partial \phi^+}{\partial x_2} + \int_{\Omega^+} \alpha(x) \bar{v}^+ \phi^+ + \int_{\Omega^-} \nabla v^- \nabla \phi^- + \int_{\Omega^-} u^- \phi^- \\ \quad = \int_{\Omega^-} (\bar{u}^- - u_d) \phi^- + \int_{\Omega^+} \int_{Y(x_2)} \left[(1 - \xi)^2 \bar{u}^+ + \xi(1 - \xi)f \right] \phi^+ \\ \quad \quad + \int_{\Omega^+} \int_{Y(x_2)} [(1 - \xi) \bar{u}_{11} - (1 - \xi)u_d] \phi^+, \\ \int_{\Omega^u} \frac{\partial \bar{v}_{11}}{\partial y_1} \frac{\partial \phi_1}{\partial y_1} + \int_{\Omega^u} \bar{v}_{11} \phi_1 = \int_{\Omega^u} [(1 - \xi) \bar{u}^+ + \xi f + \bar{u}_{11} - u_d] \phi_1, \end{array} \right.$$

for all $(\phi, \phi_1) \in H(\Omega) \times V(\Omega)$.

- **Reduced adjoint state:** The adjoint state $(\bar{v}, \bar{v}_{11}) \in H(\Omega) \times V(\Omega)$ satisfies the following system,






$$\left\{ \begin{array}{l} \int_{\Omega^+} |Y_c(x_2)| \frac{\partial \bar{v}^+}{\partial x_2} \frac{\partial \phi^+}{\partial x_2} + \int_{\Omega^+} \alpha(x) \bar{v}^+ \phi^+ + \int_{\Omega^-} \nabla v^- \nabla \phi^- + \int_{\Omega^-} u^- \phi^- \\ \quad = \int_{\Omega^-} (\bar{u}^- - u_d) \phi^- + \int_{\Omega^+} \int_{Y(x_2)} \left[(1 - \xi)^2 \bar{u}^+ + \xi(1 - \xi)f \right] \phi^+ \\ \quad \quad + \int_{\Omega^+} \int_{Y(x_2)} [(1 - \xi) \bar{u}_{11} - (1 - \xi)u_d] \phi^+, \\ \int_{\Omega^u} \frac{\partial \bar{v}_{11}}{\partial y_1} \frac{\partial \phi_1}{\partial y_1} + \int_{\Omega^u} \bar{v}_{11} \phi_1 = \int_{\Omega^u} [(1 - \xi) \bar{u}^+ + \xi f + \bar{u}_{11} - u_d] \phi_1, \end{array} \right.$$

for all $(\phi, \phi_1) \in H(\Omega) \times V(\Omega)$.

- The optimal control is given by $\bar{\theta} + \bar{\theta}_1 = -\frac{1}{\beta} [(1 - \xi) \bar{v}^+ + \bar{v}_{11}]$ in Ω_1^u .

In the above variational problem we have considered the contrasting diffusive coefficients as 1 and ε^2 . In fact, we can consider the coefficient of the form $O(1)$ and α_ε^2 , where $\alpha_\varepsilon \rightarrow 0$ as $\varepsilon \rightarrow 0$. According to the limit $k = \lim_{\varepsilon \rightarrow 0} \frac{\alpha_\varepsilon}{\varepsilon}$, we will get three different limit problems for, $k = 0, k = \infty$ and $k \in (0, \infty)$.

In the above variational problem we have considered the contrasting diffusive coefficients as 1 and ε^2 . In fact, we can consider the coefficient of the form $O(1)$ and α_ε^2 , where $\alpha_\varepsilon \rightarrow 0$ as $\varepsilon \rightarrow 0$. According to the limit $k = \lim_{\varepsilon \rightarrow 0} \frac{\alpha_\varepsilon}{\varepsilon}$, we will get three different limit problems for, $k = 0, k = \infty$ and $k \in (0, \infty)$. What we have studied is essentially the case, where $k \in (0, \infty)$, that is with $\alpha_\varepsilon = \varepsilon$ and hence the exact proof can be reproduced with minor changes. The coefficient of the second order term in the cell problem will be k^2 instead of 1. The other two cases can also be handled with minor modifications. Here we have presented the case when $k = 1$, that is $\alpha_\varepsilon = \varepsilon$.

-  S. AIYAPPAN, A. K. NANDAKUMARAN, AND RAVI PRAKASH, “Generalization of unfolding operator for highly oscillating smooth boundary domains and homogenization”, *Calculus of Variations and Partial Differential Equations* 57.3 (2018): 1-30.
-  A. GAUDIELLO, A. SILI, “Homogenization of highly oscillating boundaries with strongly contrasting diffusivity”, *SIAM J. Math. Anal.* 47 (3) (2015) 16711692, <https://doi.org/10.1137/140987225>.
-  N.S. BAKHVALOV, G.P. PANASENKO, “Averaging Processes in Periodic Media”, Nauka, Moscow, 1984.
-  A.YU. KOSAREV, “Asymptotic behavior of averaged characteristics of periodic elastic media with highly varying properties”, *Dokl. Akad. Nauk SSSR* 267 (1) (1982) 3842.
-  A. K. NANDAKUMARAN, AND ABU SUFIAN, “Strong contrasting diffusivity in general oscillating domains: Homogenization of optimal control problems”, *Journal of Differential Equations* 291 (2021): 57-89.

Thank you for your attention!