

Gersten's conjecture for Milnor K-theory

Rakesh Pawar

Postdoc, IISER Pune

18th September 2021

IISc-IISER Pune Twenty-20 Symposium

Milnor invented the K-theory of fields (denoted by K_*^M) in 1970, where he proposed a close connection between quadratic forms, his K-theory and Galois cohomology.

Let F be a field. Define

$$K_*^M(F) := \frac{\bigoplus_{n=0}^{\infty} (F^*)^{\otimes n}}{I}$$

Milnor invented the K-theory of fields (denoted by K_*^M) in 1970, where he proposed a close connection between quadratic forms, his K-theory and Galois cohomology.

Let F be a field. Define

$$K_*^M(F) := \frac{\bigoplus_{n=0}^{\infty} (F^*)^{\otimes n}}{I}$$

Here $I = \langle a \otimes b \mid a, b \in F^* \text{ and } a + b = 1 \rangle$ denotes the two-sided homogeneous ideal generated by the symbols.

Milnor invented the K-theory of fields (denoted by K_*^M) in 1970, where he proposed a close connection between quadratic forms, his K-theory and Galois cohomology.

Let F be a field. Define

$$K_*^M(F) := \frac{\bigoplus_{n=0}^{\infty} (F^*)^{\otimes n}}{I}$$

Here $I = \langle a \otimes b \mid a, b \in F^* \text{ and } a + b = 1 \rangle$ denotes the two-sided homogeneous ideal generated by the symbols.

The class of $a_1 \otimes \cdots \otimes a_n$ is denoted by $\{a_1, \cdots, a_n\}$.

Milnor invented the K-theory of fields (denoted by K_*^M) in 1970, where he proposed a close connection between quadratic forms, his K-theory and Galois cohomology.

Let F be a field. Define

$$K_*^M(F) := \frac{\bigoplus_{n=0}^{\infty} (F^*)^{\otimes n}}{I}$$

Here $I = \langle a \otimes b \mid a, b \in F^* \text{ and } a + b = 1 \rangle$ denotes the two-sided homogeneous ideal generated by the symbols.

The class of $a_1 \otimes \cdots \otimes a_n$ is denoted by $\{a_1, \dots, a_n\}$.

The following relations hold.

- 1 For $x \in K_n^M(F)$ and $y \in K_m^M(F)$, $xy = (-1)^{nm}yx$.

Milnor invented the K-theory of fields (denoted by K_*^M) in 1970, where he proposed a close connection between quadratic forms, his K-theory and Galois cohomology.

Let F be a field. Define

$$K_*^M(F) := \frac{\bigoplus_{n=0}^{\infty} (F^*)^{\otimes n}}{I}$$

Here $I = \langle a \otimes b \mid a, b \in F^* \text{ and } a + b = 1 \rangle$ denotes the two-sided homogeneous ideal generated by the symbols.

The class of $a_1 \otimes \cdots \otimes a_n$ is denoted by $\{a_1, \dots, a_n\}$.

The following relations hold.

- 1 For $x \in K_n^M(F)$ and $y \in K_m^M(F)$, $xy = (-1)^{nm}yx$.
- 2 (Steinberg relation) $\{a, 1 - a\} = 0$ for $a \in F^*$.

Milnor invented the K-theory of fields (denoted by K_*^M) in 1970, where he proposed a close connection between quadratic forms, his K-theory and Galois cohomology.

Let F be a field. Define

$$K_*^M(F) := \frac{\bigoplus_{n=0}^{\infty} (F^*)^{\otimes n}}{I}$$

Here $I = \langle a \otimes b \mid a, b \in F^* \text{ and } a + b = 1 \rangle$ denotes the two-sided homogeneous ideal generated by the symbols.

The class of $a_1 \otimes \cdots \otimes a_n$ is denoted by $\{a_1, \dots, a_n\}$.

The following relations hold.

- 1 For $x \in K_n^M(F)$ and $y \in K_m^M(F)$, $xy = (-1)^{nm}yx$.
- 2 (Steinberg relation) $\{a, 1 - a\} = 0$ for $a \in F^*$.
- 3 If $a_1, \dots, a_n \in F^*$ and if $a_1 + \cdots + a_n$ is either 0 or 1, then $\{a_1, \dots, a_n\} = 0$ in $K_n^M(F)$.

Let (F, ν) be a discrete valuation field, with the valuation ν . Let \mathcal{O} be the valuation ring.

Let (F, ν) be a discrete valuation field, with the valuation ν . Let \mathcal{O} be the valuation ring.

Let π be a uniformizer for the valuation ν with the residue field κ .

Let (F, ν) be a discrete valuation field, with the valuation ν . Let \mathcal{O} be the valuation ring.

Let π be a uniformizer for the valuation ν with the residue field κ . There is a residue map

$$\partial^\pi : K_n^M(F) \rightarrow K_{n-1}^M(\kappa)$$

Let (F, ν) be a discrete valuation field, with the valuation ν . Let \mathcal{O} be the valuation ring.

Let π be a uniformizer for the valuation ν with the residue field κ . There is a residue map

$$\partial^\pi : K_n^M(F) \rightarrow K_{n-1}^M(\kappa)$$

such that

1 for $u_i \in \mathcal{O}^*$, $\partial^\pi(\{u_1, \dots, u_n\}) = 0$

Let (F, ν) be a discrete valuation field, with the valuation ν . Let \mathcal{O} be the valuation ring.

Let π be a uniformizer for the valuation ν with the residue field κ . There is a residue map

$$\partial^\pi : K_n^M(F) \rightarrow K_{n-1}^M(\kappa)$$

such that

- 1 for $u_i \in \mathcal{O}^*$, $\partial^\pi(\{u_1, \dots, u_n\}) = 0$
- 2 for $u_i \in \mathcal{O}^*$, $\partial^\pi(\{\pi, u_2, \dots, u_n\}) = \{\bar{u}_2, \dots, \bar{u}_n\}$.

Let (F, ν) be a discrete valuation field, with the valuation ν . Let \mathcal{O} be the valuation ring.

Let π be a uniformizer for the valuation ν with the residue field κ . There is a residue map

$$\partial^\pi : K_n^M(F) \rightarrow K_{n-1}^M(\kappa)$$

such that

- 1 for $u_i \in \mathcal{O}^*$, $\partial^\pi(\{u_1, \dots, u_n\}) = 0$
- 2 for $u_i \in \mathcal{O}^*$, $\partial^\pi(\{\pi, u_2, \dots, u_n\}) = \{\bar{u}_2, \dots, \bar{u}_n\}$.

This is used to define the differential map in the Gersten's complex for Milnor K- theory.

Some computations:

- For a finite field F , $K_n^M(F) = 0$ for $n > 1$.
- For a number field F , $K_n^M(F) = (\mathbb{Z}/2\mathbb{Z})^{\oplus r_1}$ for $n > 2$, where r_1 is the number of real embeddings $F \hookrightarrow \mathbb{R}$.
- For a local field F , with finite residue field, $K_n^M(F)$ is uniquely divisible for $n > 2$ and

$$K_2^M(F) = \mu_\infty(F) \oplus \text{div}$$

where $\mu_\infty(F)$ is the group of roots of unity in F and div is uniquely divisible.

Theorem (Milnor)

There is a (split) exact sequence

$$0 \rightarrow K_{n+1}^M(F) \rightarrow K_{n+1}^M(F(T)) \xrightarrow{\partial} \bigoplus_{p(T)} K_n^M(F[T]/(p(T))) \rightarrow 0$$

The sum is over the monic irreducible polynomials $p(T)$.

Theorem (Milnor)

There is a (split) exact sequence

$$0 \rightarrow K_{n+1}^M(F) \rightarrow K_{n+1}^M(F(T)) \xrightarrow{\partial} \bigoplus_{p(T)} K_n^M(F[T]/(p(T))) \rightarrow 0$$

The sum is over the monic irreducible polynomials $p(T)$.

For a finite field extension $F \subset E$, there is a Norm map

$$Nm_{E/F} : K_n^M(E) \rightarrow K_n^M(F).$$

Theorem (Milnor)

There is a (split) exact sequence

$$0 \rightarrow K_{n+1}^M(F) \rightarrow K_{n+1}^M(F(T)) \xrightarrow{\partial} \bigoplus_{p(T)} K_n^M(F[T]/(p(T))) \rightarrow 0$$

The sum is over the monic irreducible polynomials $p(T)$.

For a finite field extension $F \subset E$, there is a Norm map

$$Nm_{E/F} : K_n^M(E) \rightarrow K_n^M(F).$$

Write $E = F[T]/(p(T))$ for a monic irreducible polynomial $p(T)$.

$$\begin{array}{ccc}
 & & s \\
 & \swarrow & \searrow \\
 K_{n+1}^M(F(T)) & \xrightarrow{\partial} & \bigoplus_{p(T)} K_n^M(F[T]/(p(T))) \\
 \downarrow \partial^{-1/T} & & \swarrow Nm_{E/F} \\
 K_n^M(F) & &
 \end{array}$$

Theorem (Voevodsky et al.)

Let F be a field of characteristic $\neq 2$. The maps below are isomorphisms for all $n \geq 0$.

$$\{a_1, \dots, a_n\} \longrightarrow (a_1) \cup \dots \cup (a_n)$$

$$\begin{array}{ccc}
 \{a\} & & K_n^M(F)/2 \\
 \downarrow & & \downarrow \text{Pfister forms} \\
 \langle 1, -a \rangle & & I_F^n / I_F^{n+1}
 \end{array}
 \qquad
 \begin{array}{ccc}
 & \xrightarrow{\text{Galois symbol}} & H^n(F, \mathbb{Z}/2(n))
 \end{array}$$

$H^n(F, \mathbb{Z}/2(n))$ denotes the weight n -Motivic cohomology of F with coefficients in $\mathbb{Z}/2$.

I_F is the fundamental ideal in the Witt ring of F of quadratic forms of even rank. I_F^n denotes its n^{th} power.

(Kato 1986) Let X be an excellent scheme of dimension d . Gersten's/Rost's cycle complex for X is defined as follows.

$$0 \rightarrow \bigoplus_{\eta \in X^{(0)}} K_n^M(\kappa(\eta)) \xrightarrow{d} \bigoplus_{x \in X^{(1)}} K_{n-1}^M(\kappa(x)) \xrightarrow{d} \cdots \xrightarrow{d} \bigoplus_{x \in X^{(d)}} K_{n-d}^M(\kappa(x)) \rightarrow 0$$

where the differential d is defined using the residue maps and the norm maps defined earlier.

Let us denote the cohomology of the complex by $A^p(X, K_n^M)$.

Let A be a regular local ring of dimension d .

Gersten's conjecture

Let A be a regular local ring of dimension d . The Gersten's conjecture states that the complex

$$0 \rightarrow K_n^M(\text{Frac}(A)) \xrightarrow{d} \bigoplus_{\mathfrak{p} \in \text{Spec } A^{(1)}} K_{n-1}^M(\kappa(\mathfrak{p})) \xrightarrow{d} \cdots \xrightarrow{d} \bigoplus_{\mathfrak{p} \in \text{Spec } A^{(d)}} K_{n-d}^M(\kappa(\mathfrak{p})) \rightarrow 0$$

is exact in degrees ≥ 1 , for each $n \geq 0$. Here $\text{Spec } A^{(p)}$ denotes the set of codimension p points of $\text{Spec } A$.

Let A be a regular local ring of dimension d . The Gersten's conjecture states that the complex

$$0 \rightarrow K_n^M(\text{Frac}(A)) \xrightarrow{d} \bigoplus_{\mathfrak{p} \in \text{Spec } A^{(1)}} K_{n-1}^M(\kappa(\mathfrak{p})) \xrightarrow{d} \cdots \xrightarrow{d} \bigoplus_{\mathfrak{p} \in \text{Spec } A^{(d)}} K_{n-d}^M(\kappa(\mathfrak{p})) \rightarrow 0$$

is exact in degrees ≥ 1 , for each $n \geq 0$. Here $\text{Spec } A^{(p)}$ denotes the set of codimension p points of $\text{Spec } A$.

The conjecture above in its most general form is largely open till now.

For A containing a field, Gersten's conjecture has been verified by M. Kerz (2009, 2010).

For A containing a field, Gersten's conjecture has been verified by M. Kerz (2009, 2010).

More recently, M. Lüders proved the following relative version.

For A containing a field, Gersten's conjecture has been verified by M. Kerz (2009, 2010).

More recently, M. Lüders proved the following relative version.

Theorem (Lüders 2020)

Let X be the spectrum of a smooth local ring over an excellent dvr, then the complex

$$0 \rightarrow K_n^M(\text{Frac}(X)) \xrightarrow{d} \bigoplus_{x \in X^{(1)}} K_{n-1}^M(\kappa(x)) \xrightarrow{d} \cdots \xrightarrow{d} \bigoplus_{x \in X^{(d)}} K_{n-d}^M(\kappa(x)) \rightarrow 0$$

is exact in degrees > 0 .

For A containing a field, Gersten's conjecture has been verified by M. Kerz (2009, 2010).

More recently, M. Lüders proved the following relative version.

Theorem (Lüders 2020)

Let X be the spectrum of a smooth local ring over an excellent dvr, then the complex

$$0 \rightarrow K_n^M(\text{Frac}(X)) \xrightarrow{d} \bigoplus_{x \in X^{(1)}} K_{n-1}^M(\kappa(x)) \xrightarrow{d} \cdots \xrightarrow{d} \bigoplus_{x \in X^{(d)}} K_{n-d}^M(\kappa(x)) \rightarrow 0$$

is exact in degrees > 0 .

Here for a dvr R , X is the Spectrum of a local ring of a smooth scheme over $\text{Spec } R$ at a point.

For A containing a field, Gersten's conjecture has been verified by M. Kerz (2009, 2010).

More recently, M. Lüders proved the following relative version.

Theorem (Lüders 2020)

Let X be the spectrum of a smooth local ring over an excellent dvr, then the complex

$$0 \rightarrow K_n^M(\text{Frac}(X)) \xrightarrow{d} \bigoplus_{x \in X^{(1)}} K_{n-1}^M(\kappa(x)) \xrightarrow{d} \cdots \xrightarrow{d} \bigoplus_{x \in X^{(d)}} K_{n-d}^M(\kappa(x)) \rightarrow 0$$

is exact in degrees > 0 .

Here for a dvr R , X is the Spectrum of a local ring of a smooth scheme over $\text{Spec } R$ at a point. This uses the works of M. Rost on "Cycle modules" and a Chinese remainder theorem trick.

We generalize the Lüders theorem to the case where the base S is the Spectrum of a regular Henselian domain and X is the Henselization of a smooth S -scheme at a point.

We generalize the Lüdgers theorem to the case where the base S is the Spectrum of a regular Henselian domain and X is the Henselization of a smooth S -scheme at a point. More concretely, we have

Theorem (———— 2021)

Let $S = \operatorname{Spec} R$ for R a regular local Henselian domain of Krull dimension ≥ 1 . Let X be an essentially smooth Henselian local S -scheme of dimension d , then

$$A^p(X, K_n^M) = 0$$

for $p \geq \dim S \geq 1$ and $n \geq 0$.

We generalize the Lüdters theorem to the case where the base S is the Spectrum of a regular Henselian domain and X is the Henselization of a smooth S -scheme at a point. More concretely, we have

Theorem (——— 2021)

Let $S = \operatorname{Spec} R$ for R a regular local Henselian domain of Krull dimension ≥ 1 . Let X be an essentially smooth Henselian local S -scheme of dimension d , then

$$A^p(X, K_n^M) = 0$$

for $p \geq \dim S \geq 1$ and $n \geq 0$. More explicitly,

$$0 \rightarrow K_n^M(\operatorname{Frac}(X)) \xrightarrow{d} \bigoplus_{x \in X^{(1)}} K_{n-1}^M(\kappa(x)) \xrightarrow{d} \cdots \xrightarrow{d} \bigoplus_{x \in X^{(d)}} K_{n-d}^M(\kappa(x)) \rightarrow 0$$

is exact in degrees $p \geq \dim S$.

We generalize the Lüdters theorem to the case where the base S is the Spectrum of a regular Henselian domain and X is the Henselization of a smooth S -scheme at a point. More concretely, we have

Theorem (——— 2021)

Let $S = \operatorname{Spec} R$ for R a regular local Henselian domain of Krull dimension ≥ 1 . Let X be an essentially smooth Henselian local S -scheme of dimension d , then

$$A^p(X, K_n^M) = 0$$

for $p \geq \dim S \geq 1$ and $n \geq 0$. More explicitly,

$$0 \rightarrow K_n^M(\operatorname{Frac}(X)) \xrightarrow{d} \bigoplus_{x \in X^{(1)}} K_{n-1}^M(\kappa(x)) \xrightarrow{d} \cdots \xrightarrow{d} \bigoplus_{x \in X^{(d)}} K_{n-d}^M(\kappa(x)) \rightarrow 0$$

is exact in degrees $p \geq \dim S$.

Here for a regular local Henselian domain R , X is the Spectrum of a local ring of a smooth scheme over $\operatorname{Spec} R$ at a point.

Theorem (Lüders 2020)

Let X be the spectrum of a smooth local ring over an excellent dvr R , then the complex

$$0 \rightarrow K_n^M(\text{Frac}(X)) \xrightarrow{d} \bigoplus_{x \in X^{(1)}} K_{n-1}^M(\kappa(x)) \xrightarrow{d} \cdots \xrightarrow{d} \bigoplus_{x \in X^{(d)}} K_{n-d}^M(\kappa(x)) \rightarrow 0$$

is exact in degrees > 0 .

Sketch of the proof of Lüders' theorem


Theorem (Lüders 2020)


Let X be the spectrum of a smooth local ring over an excellent dvr R , then the complex

$$0 \rightarrow K_n^M(\text{Frac}(X)) \xrightarrow{d} \bigoplus_{x \in X^{(1)}} K_{n-1}^M(\kappa(x)) \xrightarrow{d} \cdots \xrightarrow{d} \bigoplus_{x \in X^{(d)}} K_{n-d}^M(\kappa(x)) \rightarrow 0$$

is exact in degrees > 0 .

Let $S = \text{Spec } R$. $X = \text{Spec } \mathcal{O}_{V,v}$, for a smooth S -scheme V . Given a Zariski nbd U of v in V , we can think of U as follows.





There are two types of irreducible closed subschemes of positive codimension. Ones which are **vertical** and others which are **horizontal**.

$$A^p(X, K_n^M) = \varinjlim_{(U, u)} A^p(U, K_n^M)$$

where the colimit runs over (U, u) such that U is a Zariski neighbourhood of v in V .

$$A^p(X, K_n^M) = \varinjlim_{(U, u)} A^p(U, K_n^M)$$

where the colimit runs over (U, u) such that U is a Zariski neighbourhood of v in V . Now we see that

$$A^p(U, K_n^M) = \varinjlim_{Y \subset U} A_Y^p(U, K_n^M)$$

where the colimit runs over all closed subschemes Y of U .

$$A^p(X, K_n^M) = \varinjlim_{(U, u)} A^p(U, K_n^M)$$

where the colimit runs over (U, u) such that U is a Zariski neighbourhood of v in V . Now we see that

$$A^p(U, K_n^M) = \varinjlim_{Y \subset U} A_Y^p(U, K_n^M)$$

where the colimit runs over all closed subschemes Y of U .

Idea:

- Every cohomology class in degree $p > 0$ is represented by a class which is supported on closed subschemes which are flat over the base S (horizontal).

$$A^p(X, K_n^M) = \varinjlim_{(U, u)} A^p(U, K_n^M)$$

where the colimit runs over (U, u) such that U is a Zariski neighbourhood of v in V . Now we see that

$$A^p(U, K_n^M) = \varinjlim_{Y \subset U} A_Y^p(U, K_n^M)$$


where the colimit runs over all closed subschemes Y of U .

Idea:


- Every cohomology class in degree $p > 0$ is represented by a class which is supported on closed subschemes which are flat over the base S (horizontal).
- Given a closed subscheme Y in U flat (horizontal) over S , there is a smaller Zariski neighbourhood U' of v in U such that the map

$$A_{Y \times_U U'}^p(U', K_n^M) \rightarrow A^p(U', K_n^M)$$


is 0. (This uses a technical result, Gillet-Levine's version of **Geometric Presentation lemma**).



Let $y \in U^{(p)}$ for $p > 0$. Let $\{\bar{\alpha}_1, \bar{\alpha}_2, \dots, \bar{\alpha}_{n-p}\} \in K_{n-p}^M(\kappa(y))$.




Let $y \in U^{(p)}$ for $p > 0$. Let $\{\bar{\alpha}_1, \bar{\alpha}_2, \dots, \bar{\alpha}_{n-p}\} \in K_{n-p}^M(\kappa(y))$.
 (Saito-Sato) Find an integral closed subscheme Z of codimension $p - 1$ containing $\overline{\{y\}}$ such that Z is regular at y and $Z \cap V_\eta \neq \emptyset$.



Let $y \in U^{(p)}$ for $p > 0$. Let $\{\bar{\alpha}_1, \bar{\alpha}_2, \dots, \bar{\alpha}_{n-p}\} \in K_{n-p}^M(\kappa(y))$.

(Saito-Sato) Find a closed subscheme Z of codimension $p - 1$ containing $\overline{\{y\}}$ such that Z is regular at y and $Z \cap U_\eta \neq \emptyset$.

Now codimension 1 points of Z are y and the points w such that $\overline{\{w\}}$ are **horizontal**.



Now using Chinese remainder theorem, one constructs a Milnor K-theory symbol β on $\kappa(Z)$ such that

$$d(\beta) = \{\bar{\alpha}_1, \bar{\alpha}_2, \dots, \bar{\alpha}_{n-p}\} + \sum_{\{w_i\}} \gamma_i.$$

This finishes the argument for the first step.

Lüders, M., On the relative Gersten conjecture for Milnor K-theory in the smooth case, arXiv:2010.02622 (2020).

Pawar, R., A remark on relative Gersten's complex for Milnor K-theory, arXiv:2105.06962.

Thank you!