

# On the growth of cuspidal cohomology of $GL_4$ by symmetric cube transfer

Sudipa Mondal

(Based on joint work with Dr. Chandrasheel Bhagwat)

IISER Pune

(IISC-IISERP 20-20 symposium)

September 19, 2021

# Contents

- 1 Introduction
- 2 Notations
- 3 Cuspidal cohomology of  $GL_4$
- 4 References

# 1 Introduction

## 2 Notations

## 3 Cuspidal cohomology of $GL_4$

## 4 References

# Introduction

- Let  $\mathbb{E} = \mathbb{Q}(\sqrt{-d})$  be an imaginary quadratic extension of  $\mathbb{Q}$ .

# Introduction

- Let  $\mathbb{E} = \mathbb{Q}(\sqrt{-d})$  be an imaginary quadratic extension of  $\mathbb{Q}$ .
- Consider the group of idèles  $\mathbb{A}_{\mathbb{E}}^{\times}$ .

# Introduction

- Let  $\mathbb{E} = \mathbb{Q}(\sqrt{-d})$  be an imaginary quadratic extension of  $\mathbb{Q}$ .
- Consider the group of idèles  $\mathbb{A}_{\mathbb{E}}^{\times}$ .
- Using automorphic induction, every character of  $\mathbb{A}_{\mathbb{E}}^{\times}$  yields an automorphic form  $\phi$  for  $\mathrm{GL}_2(\mathbb{A}_{\mathbb{Q}})$ .

# Introduction

- Let  $\mathbb{E} = \mathbb{Q}(\sqrt{-d})$  be an imaginary quadratic extension of  $\mathbb{Q}$ .
- Consider the group of idèles  $\mathbb{A}_{\mathbb{E}}^{\times}$ .
- Using automorphic induction, every character of  $\mathbb{A}_{\mathbb{E}}^{\times}$  yields an automorphic form  $\phi$  for  $\mathrm{GL}_2(\mathbb{A}_{\mathbb{Q}})$ .
- It is a mass form if  $\mathbb{E}$  is real quadratic field and a holomorphic modular form if  $\mathbb{E}$  is imaginary quadratic field.

- Moreover,  $\phi$  is a cusp form if it does not factor through the norm map.



- Moreover,  $\phi$  is a cusp form if it does not factor through the norm map.
- The relation between the automorphic representations of  $\mathrm{GL}_2(\mathbb{A}_{\mathbb{Q}})$  and the cuspidal cohomology of  $\mathrm{GL}_2$  can be described in terms of the sheaf  $\tilde{M}_{\mu}$  associated to the highest weight representation  $M_{\mu}$  of  $\mathrm{GL}_2(\mathbb{R})$ .

- Moreover,  $\phi$  is a cusp form if it does not factor through the norm map.
- The relation between the automorphic representations of  $\mathrm{GL}_2(\mathbb{A}_{\mathbb{Q}})$  and the cuspidal cohomology of  $\mathrm{GL}_2$  can be described in terms of the sheaf  $\tilde{M}_{\mu}$  associated to the highest weight representation  $M_{\mu}$  of  $\mathrm{GL}_2(\mathbb{R})$ .
- Also, for a fixed weight and level structure of  $\mathrm{GL}_2(\mathbb{A}_{\mathbb{Q}})$ , the space of cusp forms has finite dimension.

- Moreover,  $\phi$  is a cusp form if it does not factor through the norm map.
- The relation between the automorphic representations of  $\mathrm{GL}_2(\mathbb{A}_{\mathbb{Q}})$  and the cuspidal cohomology of  $\mathrm{GL}_2$  can be described in terms of the sheaf  $\tilde{M}_{\mu}$  associated to the highest weight representation  $M_{\mu}$  of  $\mathrm{GL}_2(\mathbb{R})$ .
- Also, for a fixed weight and level structure of  $\mathrm{GL}_2(\mathbb{A}_{\mathbb{Q}})$ , the space of cusp forms has finite dimension.
- One can ask, how much of cuspidal cohomology of  $\mathrm{GL}_2(\mathbb{A}_{\mathbb{Q}})$  is obtained by automorphic induction.

- C. Ambi in [AM] has estimated the dimension for a fixed weight and varying level structure.

- C. Ambi in [AM] has estimated the dimension for a fixed weight and varying level structure.
- Let  $C_k(N)$  denote the set of normalised cusp eigenforms of Hecke operators of  $\Gamma_1(N)$  of weight  $k$  obtained by automorphic induction of Größencharacters of imaginary quadratic extensions.

- C. Ambi in [AM] has estimated the dimension for a fixed weight and varying level structure.
- Let  $C_k(N)$  denote the set of normalised cusp eigenforms of Hecke operators of  $\Gamma_1(N)$  of weight  $k$  obtained by automorphic induction of Größencharacters of imaginary quadratic extensions.

### Theorem 1 (Ambi)

Let  $k \geq 1, \epsilon \in (0, 1)$ . Then,

$$|C_k(N)| \ll_{k, \epsilon} N \cdot N'^{(1+\epsilon)} \quad \text{as } N \longrightarrow \infty$$

where  $N'$  is the product of all distinct prime factors of  $N$ .

- Consider a cuspidal automorphic representation  $\pi$  of  $\mathrm{GL}_2(\mathbb{A}_{\mathbb{Q}})$ .

- Consider a cuspidal automorphic representation  $\pi$  of  $\mathrm{GL}_2(\mathbb{A}_{\mathbb{Q}})$ .
- Then  $\mathrm{sym}^2(\pi)$  is an automorphic representation of  $\mathrm{GL}_3(\mathbb{A}_{\mathbb{Q}})$ .



- Consider a cuspidal automorphic representation  $\pi$  of  $\mathrm{GL}_2(\mathbb{A}_{\mathbb{Q}})$ .
- Then  $\mathrm{sym}^2(\pi)$  is an automorphic representation of  $\mathrm{GL}_3(\mathbb{A}_{\mathbb{Q}})$ .
- Similarly, one can ask, how much of cuspidal cohomology of  $\mathrm{GL}_3(\mathbb{A}_{\mathbb{Q}})$  is obtained by symmetric square transfer.

- Consider a cuspidal automorphic representation  $\pi$  of  $\mathrm{GL}_2(\mathbb{A}_{\mathbb{Q}})$ .
- Then  $\mathrm{sym}^2(\pi)$  is an automorphic representation of  $\mathrm{GL}_3(\mathbb{A}_{\mathbb{Q}})$ .
- Similarly, one can ask, how much of cuspidal cohomology of  $\mathrm{GL}_3(\mathbb{A}_{\mathbb{Q}})$  is obtained by symmetric square transfer.
- Again in [AM], C. Ambi estimated the above number by fixing the weight and varying the level structure.

We want to generalize this for symmetric cube transfer.

We want to generalize this for symmetric cube transfer.

- Let  $\pi$  be a cuspidal automorphic representation of  $\mathrm{GL}_2(\mathbb{A}_{\mathbb{Q}})$ .

We want to generalize this for symmetric cube transfer.

- Let  $\pi$  be a cuspidal automorphic representation of  $\mathrm{GL}_2(\mathbb{A}_{\mathbb{Q}})$ .
- Then  $\mathrm{sym}^3(\pi)$  is a representation of  $\mathrm{GL}_4(\mathbb{A}_{\mathbb{Q}})$ .

We want to generalize this for symmetric cube transfer.

- Let  $\pi$  be a cuspidal automorphic representation of  $\mathrm{GL}_2(\mathbb{A}_{\mathbb{Q}})$ .
- Then  $\mathrm{sym}^3(\pi)$  is a representation of  $\mathrm{GL}_4(\mathbb{A}_{\mathbb{Q}})$ .
- Due to Kim and Shahidi [Theorem 6.1, [KS]],  $\mathrm{sym}^3(\pi)$  is an automorphic representation of  $\mathrm{GL}_4(\mathbb{A}_{\mathbb{Q}})$ .

We want to generalize this for symmetric cube transfer.

- Let  $\pi$  be a cuspidal automorphic representation of  $\mathrm{GL}_2(\mathbb{A}_{\mathbb{Q}})$ .
- Then  $\mathrm{sym}^3(\pi)$  is a representation of  $\mathrm{GL}_4(\mathbb{A}_{\mathbb{Q}})$ .
- Due to Kim and Shahidi [Theorem 6.1, [KS]],  $\mathrm{sym}^3(\pi)$  is an automorphic representation of  $\mathrm{GL}_4(\mathbb{A}_{\mathbb{Q}})$ . Furthermore, it is cuspidal if the representation  $\pi$  is not obtained by Automorphic induction.

We want to generalize this for symmetric cube transfer.

- Let  $\pi$  be a cuspidal automorphic representation of  $\mathrm{GL}_2(\mathbb{A}_{\mathbb{Q}})$ .
- Then  $\mathrm{sym}^3(\pi)$  is a representation of  $\mathrm{GL}_4(\mathbb{A}_{\mathbb{Q}})$ .
- Due to Kim and Shahidi [Theorem 6.1, [KS]],  $\mathrm{sym}^3(\pi)$  is an automorphic representation of  $\mathrm{GL}_4(\mathbb{A}_{\mathbb{Q}})$ . Furthermore, it is cuspidal if the representation  $\pi$  is not obtained by Automorphic induction.
- Hence we estimate the cuspidal cohomology of  $\mathrm{GL}_4(\mathbb{A}_{\mathbb{Q}})$  which is obtained by symmetric cube transfer from  $\mathrm{GL}_2(\mathbb{A}_{\mathbb{Q}})$  corresponding to a specific level structure.



1 Introduction

2 Notations

3 Cuspidal cohomology of  $GL_4$

4 References

# The level structure

Consider the algebraic group  $GL_m$ . For a finite prime  $p$  of  $\mathbb{Q}$ , let  $\mathbb{Z}_p$  be the ring of  $p$ -adic integers.

# The level structure

Consider the algebraic group  $GL_m$ . For a finite prime  $p$  of  $\mathbb{Q}$ , let  $\mathbb{Z}_p$  be the ring of  $p$ -adic integers. For each integer  $n \geq 0$ , define

$$K_p^m(n) := \{x = (x_{i,j})_{m \times m} \in GL_m(\mathbb{Z}_p) : x_{m,k} \in p^n \mathbb{Z}_p, \ 1 \leq k < m, \\ x_{m,m} - 1 \in p^n \mathbb{Z}_p\}.$$

# The level structure

Consider the algebraic group  $GL_m$ . For a finite prime  $p$  of  $\mathbb{Q}$ , let  $\mathbb{Z}_p$  be the ring of  $p$ -adic integers. For each integer  $n \geq 0$ , define

$$K_p^m(n) := \{x = (x_{i,j})_{m \times m} \in GL_m(\mathbb{Z}_p) : x_{m,k} \in p^n \mathbb{Z}_p, \ 1 \leq k < m, \\ x_{m,m} - 1 \in p^n \mathbb{Z}_p\}.$$

$$\text{Let } N = \prod_{i=1}^r p_i^{n_i}.$$

# The level structure

Consider the algebraic group  $GL_m$ . For a finite prime  $p$  of  $\mathbb{Q}$ , let  $\mathbb{Z}_p$  be the ring of  $p$ -adic integers. For each integer  $n \geq 0$ , define

$$K_p^m(n) := \{x = (x_{i,j})_{m \times m} \in GL_m(\mathbb{Z}_p) : x_{m,k} \in p^n \mathbb{Z}_p, 1 \leq k < m, \\ x_{m,m} - 1 \in p^n \mathbb{Z}_p\}.$$

Let  $N = \prod_{i=1}^r p_i^{n_i}$ . Define a compact open subgroup  $K_f^m(N) = \prod_p K_p$  of  $GL_m(\mathbb{A}_f)$  where

$$K_p = \begin{cases} K_{p_i}^m(n_i) & \text{if } p \mid N \text{ i.e., } p = p_i \\ GL_m(\mathbb{Z}_p) & \text{if } p \nmid N \end{cases}$$

# The level structure

Consider the algebraic group  $GL_m$ . For a finite prime  $p$  of  $\mathbb{Q}$ , let  $\mathbb{Z}_p$  be the ring of  $p$ -adic integers. For each integer  $n \geq 0$ , define

$$K_p^m(n) := \{x = (x_{i,j})_{m \times m} \in GL_m(\mathbb{Z}_p) : x_{m,k} \in p^n \mathbb{Z}_p, \ 1 \leq k < m, \\ x_{m,m} - 1 \in p^n \mathbb{Z}_p\}.$$

Let  $N = \prod_{i=1}^r p_i^{n_i}$ . Define a compact open subgroup  $K_f^m(N) = \prod_p K_p$  of  $GL_m(\mathbb{A}_f)$  where

$$K_p = \begin{cases} K_{p_i}^m(n_i) & \text{if } p \mid N \text{ i.e., } p = p_i \\ GL_m(\mathbb{Z}_p) & \text{if } p \nmid N \end{cases}$$

$K_f^m(N) \subseteq GL_m(\mathbb{A}_f)$  is called the level structure corresponding to  $N$ .

# Conductor of a representation

- Let  $(\rho, \mathcal{H}) = (\bigotimes_{p \leq \infty} \rho_p, \bigotimes_{p \leq \infty} \mathcal{H}_p)$  be an irreducible automorphic representation of  $\mathrm{GL}_m(\mathbb{A}_{\mathbb{Q}})$ .

# Conductor of a representation

- Let  $(\rho, \mathcal{H}) = (\bigotimes_{p \leq \infty} \rho_p, \bigotimes_{p \leq \infty} \mathcal{H}_p)$  be an irreducible automorphic representation of  $\mathrm{GL}_m(\mathbb{A}_{\mathbb{Q}})$ .

For each finite prime  $p$ , the conductor of  $\rho_p$  is defined to be the smallest integer  $c(\rho_p) \geq 0$  such that the set  $\mathcal{H}_p^{K_p^m(c(\rho_p))}$  consisting of all  $K_p^m(c(\rho_p))$ -fixed vectors of  $\mathcal{H}_p$  is non-zero.



# Conductor of a representation

- Let  $(\rho, \mathcal{H}) = (\bigotimes_{p \leq \infty} \rho_p, \bigotimes_{p \leq \infty} \mathcal{H}_p)$  be an irreducible automorphic representation of  $\mathrm{GL}_m(\mathbb{A}_{\mathbb{Q}})$ .

For each finite prime  $p$ , the conductor of  $\rho_p$  is defined to be the smallest integer  $c(\rho_p) \geq 0$  such that the set  $\mathcal{H}_p^{K_p^m(c(\rho_p))}$  consisting of all  $K_p^m(c(\rho_p))$ -fixed vectors of  $\mathcal{H}_p$  is non-zero.

- The conductor of  $\rho$  is defined as

$$N_{\rho} = \prod_{p < \infty} p^{c(\rho_p)}.$$

1 Introduction

2 Notations

3 Cuspidal cohomology of  $GL_4$

4 References

- Let  $\pi$  be a cuspidal automorphic representation of  $\mathrm{GL}_4(\mathbb{A}_{\mathbb{Q}})$  with  $\pi_{\infty}$  and  $\pi_f$  be its infinite and finite parts respectively.

- Let  $\pi$  be a cuspidal automorphic representation of  $\mathrm{GL}_4(\mathbb{A}_{\mathbb{Q}})$  with  $\pi_{\infty}$  and  $\pi_f$  be its infinite and finite parts respectively.
- Let  $\mu$  be a dominant integral weight corresponding to the standard bores subgroup and  $M_{\mu, \mathbb{C}}$  be the underlying vector space.

- Let  $\pi$  be a cuspidal automorphic representation of  $\mathrm{GL}_4(\mathbb{A}_{\mathbb{Q}})$  with  $\pi_{\infty}$  and  $\pi_f$  be its infinite and finite parts respectively.
- Let  $\mu$  be a dominant integral weight corresponding to the standard borel subgroup and  $M_{\mu, \mathbb{C}}$  be the underlying vector space.
- For a level structure  $K_f \subset \mathrm{GL}_4(\mathbb{A}_f)$ , we write  $\pi_f^{K_f}$  to be the  $K_f$ -fixed vectors of  $\pi_f$ .

- Let  $\pi$  be a cuspidal automorphic representation of  $\mathrm{GL}_4(\mathbb{A}_{\mathbb{Q}})$  with  $\pi_{\infty}$  and  $\pi_f$  be its infinite and finite parts respectively.
- Let  $\mu$  be a dominant integral weight corresponding to the standard borel subgroup and  $M_{\mu, \mathbb{C}}$  be the underlying vector space.
- For a level structure  $K_f \subset \mathrm{GL}_4(\mathbb{A}_f)$ , we write  $\pi_f^{K_f}$  to be the  $K_f$ -fixed vectors of  $\pi_f$ .

We write  $\pi \in \mathrm{Coh}(\mathrm{GL}_4(\mathbb{A}_{\mathbb{Q}}), \mu, K_f)$  if for the relative Lie algebra cohomology,

$$H^*(\mathfrak{gl}_4(\mathbb{R}), \mathbb{R}_+^* \cdot \mathrm{SO}_4(\mathbb{R}), \pi_{\infty} \otimes M_{\mu, \mathbb{C}}) \otimes \pi_f^{K_f} \neq 0.$$

## Theorem 2 (Theorem 6.1, [KS])

*Let  $\mathbb{F}$  be a number field and  $\pi$  be a automorphic cuspidal representation of  $\mathrm{GL}_2(\mathbb{A}_{\mathbb{F}})$ . Then  $\mathrm{sym}^3(\pi)$  is automorphic representation of  $\mathrm{GL}_4(\mathbb{A}_{\mathbb{F}})$ .*

## Theorem 2 (Theorem 6.1, [KS])

*Let  $\mathbb{F}$  be a number field and  $\pi$  be a automorphic cuspidal representation of  $\mathrm{GL}_2(\mathbb{A}_{\mathbb{F}})$ . Then  $\mathrm{sym}^3(\pi)$  is automorphic representation of  $\mathrm{GL}_4(\mathbb{A}_{\mathbb{F}})$ .*

*$\mathrm{sym}^3(\pi)$  is cuspidal unless  $\pi$  is dihedral or it is tetrahedral type.*



## Theorem 2 (Theorem 6.1, [KS])

*Let  $\mathbb{F}$  be a number field and  $\pi$  be a automorphic cuspidal representation of  $\mathrm{GL}_2(\mathbb{A}_{\mathbb{F}})$ . Then  $\mathrm{sym}^3(\pi)$  is automorphic representation of  $\mathrm{GL}_4(\mathbb{A}_{\mathbb{F}})$ .*

*$\mathrm{sym}^3(\pi)$  is cuspidal unless  $\pi$  is dihedral or it is tetrahedral type.*

*In particular, if  $\mathbb{F} = \mathbb{Q}$  and  $\pi$  is the automorphic cuspidal representation attached to a non-dihedral holomorphic form of weight  $\geq 2$ , then  $\mathrm{sym}^3(\pi)$  is cuspidal.*

- Let  $E_k(N)$  denote the set of cuspidal automorphic representations of  $\mathrm{GL}_4(\mathbb{A}_{\mathbb{Q}})$  corresponding to the level structure  $K_f^4(N)$  and highest weight  $\nu_k$  which are obtained by symmetric cube transfer of cuspidal automorphic representation of  $\mathrm{GL}_2(\mathbb{A}_{\mathbb{Q}})$  of highest weight

$$\lambda_k = \left( \frac{k}{2} - 1, 1 - \frac{k}{2} \right).$$

- Let  $E_k(N)$  denote the set of cuspidal automorphic representations of  $\mathrm{GL}_4(\mathbb{A}_{\mathbb{Q}})$  corresponding to the level structure  $K_f^4(N)$  and highest weight  $\nu_k$  which are obtained by symmetric cube transfer of cuspidal automorphic representation of  $\mathrm{GL}_2(\mathbb{A}_{\mathbb{Q}})$  of highest weight

$$\lambda_k = \left( \frac{k}{2} - 1, 1 - \frac{k}{2} \right).$$

### Theorem 3 (Bhagwat,...)

Let  $k \geq 2$  be an even integer and  $p \geq 3$  be a prime. Then

$$|E_k(p^n)| \gg_k p^n \quad \text{as } n \rightarrow \infty$$

where the implied constant depends on  $k$ .

## Lemma 4

*Let  $\pi = \bigotimes \pi_p$  be a cohomological cuspidal automorphic representation of  $\mathrm{GL}_2(\mathbb{A}_{\mathbb{Q}})$ .*

## Lemma 4

*Let  $\pi = \bigotimes \pi_p$  be a cohomological cuspidal automorphic representation of  $\mathrm{GL}_2(\mathbb{A}_{\mathbb{Q}})$ . Assume that the highest weight corresponding to  $\pi_{\infty}$  is  $\lambda_k := (k/2 - 1, 1 - k/2)$  where  $k \geq 2$  is an even integer.*

## Lemma 4

Let  $\pi = \otimes \pi_p$  be a cohomological cuspidal automorphic representation of  $GL_2(\mathbb{A}_{\mathbb{Q}})$ . Assume that the highest weight corresponding to  $\pi_{\infty}$  is  $\lambda_k := (k/2 - 1, 1 - k/2)$  where  $k \geq 2$  is an even integer. Let  $\Pi = \text{sym}^3(\pi)$  be the representation of  $GL_4(\mathbb{A}_{\mathbb{Q}})$  obtained by symmetric cube transfer. Let  $c(\Pi)$  be its conductor and  $\nu_k$  be the highest weight corresponding to  $\Pi_{\infty}$ .

## Lemma 4

Let  $\pi = \otimes \pi_p$  be a cohomological cuspidal automorphic representation of  $GL_2(\mathbb{A}_{\mathbb{Q}})$ . Assume that the highest weight corresponding to  $\pi_{\infty}$  is  $\lambda_k := (k/2 - 1, 1 - k/2)$  where  $k \geq 2$  is an even integer. Let  $\Pi = \text{sym}^3(\pi)$  be the representation of  $GL_4(\mathbb{A}_{\mathbb{Q}})$  obtained by symmetric cube transfer. Let  $c(\Pi)$  be its conductor and  $\nu_k$  be the highest weight corresponding to  $\Pi_{\infty}$ .

Then the following holds:

$$\Pi \in \text{Coh}(GL_4(\mathbb{A}_{\mathbb{Q}}), \nu_k, K_f^4(c(\Pi)),$$

$$\nu_k = \left( 3 \left( \frac{k}{2} - 1 \right), \frac{k}{2} - 1, 1 - \frac{k}{2}, 3 \left( 1 - \frac{k}{2} \right) \right)$$

We now give an outline of the proof of theorem 3.

### proof

Let  $\pi = \bigotimes_{q \leq \infty} \pi_q$  be a cuspidal automorphic representation of  $\mathrm{GL}_2(\mathbb{A}_{\mathbb{Q}})$  corresponding to the level structure  $K_f^2(p^n)$ ,  $n \geq 1$ .



We now give an outline of the proof of theorem 3.

### proof

Let  $\pi = \bigotimes_{q \leq \infty} \pi_q$  be an cuspidal automorphic representation of  $\mathrm{GL}_2(\mathbb{A}_{\mathbb{Q}})$  corresponding to the level structure  $K_f^2(p^n)$ ,  $n \geq 1$ .

Step 1: Let  $c(\pi_p), c(\Pi_p)$  be the conductor of the representations of  $\pi_p$  and  $\Pi_p = \mathrm{sym}^3(\pi_p)$  respectively. Then, for odd prime  $p$  and taking  $\pi_p$  to be supercuspidal, we have,  $c(\pi_p) \leq c(\Pi_p) \leq 2c(\pi_p)$ .

## Proof(continued).

Step 2: By Theorem 2, either  $\text{sym}^3(\pi) \in \cup_{(j \geq 1)} E_k(p^j)$  or  $\pi$  is obtained by automorphic induction.

## Proof(continued).

Step 2: By Theorem 2, either  $\text{sym}^3(\pi) \in \cup_{(j \geq 1)} E_k(p^j)$  or  $\pi$  is obtained by automorphic induction. Also we have,  $\pi_p$  correspond to a unique newform in  $S_k^{\text{new}}(\Gamma_1(p^i))$  for some  $1 \leq i \leq n$  with some character.

## Proof(continued).

Step 2: By Theorem 2, either  $\text{sym}^3(\pi) \in \cup_{(j \geq 1)} E_k(p^j)$  or  $\pi$  is obtained by automorphic induction. Also we have,  $\pi_p$  correspond to a unique newform in  $S_k^{\text{new}}(\Gamma_1(p^i))$  for some  $1 \leq i \leq n$  with some character.

We will count the contribution of supercuspidal representations  $\pi_p$  to  $\cup_{(j \geq 1)} E_k(p^j)$ .

## Proof(continued).

Step 2: By Theorem 2, either  $\text{sym}^3(\pi) \in \cup_{(j \geq 1)} E_k(p^j)$  or  $\pi$  is obtained by automorphic induction. Also we have,  $\pi_p$  correspond to a unique newform in  $S_k^{\text{new}}(\Gamma_1(p^i))$  for some  $1 \leq i \leq n$  with some character.

We will count the contribution of supercuspidal representations  $\pi_p$  to  $\cup_{(j \geq 1)} E_k(p^j)$ .

Using this and the relation between the conductors, we get

$$\{ \text{supercuspidal within } \oplus_{1 \leq i \leq n} S_k^{\text{new}}(\Gamma(p^i)) \setminus C_k(p^n) \} \subseteq E_k(p^{2n}).$$

## Proof(continued).

Step 2: By Theorem 2, either  $\text{sym}^3(\pi) \in \cup_{(j \geq 1)} E_k(p^j)$  or  $\pi$  is obtained by automorphic induction. Also we have,  $\pi_p$  correspond to a unique newform in  $S_k^{\text{new}}(\Gamma_1(p^i))$  for some  $1 \leq i \leq n$  with some character.

We will count the contribution of supercuspidal representations  $\pi_p$  to  $\cup_{(j \geq 1)} E_k(p^j)$ .

Using this and the relation between the conductors, we get

$$\{ \text{supercuspidal within } \oplus_{1 \leq i \leq n} S_k^{\text{new}}(\Gamma(p^i)) \setminus C_k(p^n) \} \subseteq E_k(p^{2n}).$$

We now use Theorem 1 and the dimension formula of the space of newforms to get the lower bound.

Using the fact that  $\dim_{\mathbb{C}} S_k(\Gamma_1(N)) \sim_k N^2$ , we have the following upper bound of  $|E_k(p^n)|$  :

Using the fact that  $\dim_{\mathbb{C}} S_k(\Gamma_1(N)) \sim_k N^2$ , we have the following upper bound of  $|E_k(p^n)|$  :

### Corollary 5

For  $p \geq 3$ ,

$$|E_k(p^n)| \ll_k p^{2n} \quad \text{as } n \rightarrow \infty.$$



1 Introduction

2 Notations

3 Cuspidal cohomology of  $GL_4$

4 References

- [KS] Kim, Henry H., and Freydoon Shahidi. *Functorial products for  $GL(2) \times GL(3)$  and the symmetric cube for  $GL(2)$* . *Annals of mathematics* (2002): 837-893.
- [AM] Ambi, Chaitanya. *On the growth of cuspidal cohomology of  $GL(2)$  and  $GL(3)$* . *Journal of Number Theory* (2020).
- [BM] Bhagwat, Chandrasheel, and Sudipa Mondal. *On the growth of cuspidal cohomology of  $GL_4$* . *Journal of Number Theory* (2021).

**Thank you**